

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Chlorido{5-chloro-2-[2-(methylsulfanyl)-phenyldiazenyl]phenyl}platinum(II)

Vivek Bagchi and Debkumar Bandyopadhyay*

 Department of Chemistry, Indian Institute of Technology, New Delhi 110 016, India
 Correspondence e-mail: dkbp@chemistry.iitd.ac.in


Received 3 January 2008; accepted 9 January 2008

 Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å;
 R factor = 0.018; wR factor = 0.044; data-to-parameter ratio = 14.7.

The title compound, $[\text{Pt}(\text{C}_{13}\text{H}_{10}\text{ClN}_2\text{S})\text{Cl}]$, contains a Pt atom tetracoordinated by a benzene C, a diazene N, a Cl and an S atom in an approximately square-planar geometry. The molecules dimerize through a nonbonded $\text{S}\cdots\text{S}$ interaction [$\text{S}\cdots\text{S} = 3.523$ (18) Å]. There are no hydrogen bonds and the crystal packing is stabilized by four intermolecular π - π interactions; the centroid-centroid distances are 3.804 (3), 3.638 (3), 3.804 (3) and 3.638 (3) Å, and the corresponding perpendicular distances are 3.369, 3.448, 3.406 and 3.466 Å.

Related literature

For related literature, see: Bagchi *et al.* (2007); Chattopadhyay *et al.* (1991); Dupont *et al.* (2005); Buraway & Vellins (1954).



Experimental

Crystal data

 $[\text{Pt}(\text{C}_{13}\text{H}_{10}\text{ClN}_2\text{S})\text{Cl}]$
 $M_r = 492.28$
 Triclinic, $P\bar{1}$
 $a = 7.424$ (2) Å
 $b = 8.777$ (3) Å
 $c = 11.069$ (3) Å

 $\alpha = 105.428$ (4)°
 $\beta = 91.798$ (4)°
 $\gamma = 96.641$ (4)°
 $V = 689.1$ (4) Å³
 $Z = 2$

 Mo $K\alpha$ radiation
 $\mu = 10.70$ mm⁻¹
 $T = 273$ (2) K
 $0.42 \times 0.28 \times 0.19$ mm

Data collection

 Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.035$, $T_{\max} = 0.134$
 6783 measured reflections
 2544 independent reflections
 2472 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.018$
 $wR(F^2) = 0.043$
 $S = 1.05$
 2544 reflections
 173 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.89$ e Å⁻³
 $\Delta\rho_{\min} = -0.89$ e Å⁻³
Table 1

Selected geometric parameters (Å, °).

Pt1—N1	1.959 (3)	Pt1—Cl1	2.2911 (11)
Pt1—Cl2	1.986 (3)	Pt1—S1	2.3529 (10)
N1—Pt1—Cl2	79.26 (14)	N1—Pt1—S1	86.25 (9)
Cl2—Pt1—Cl1	96.84 (11)	Cl1—Pt1—S1	97.57 (4)

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We thank the Department of Science and Technology, India (project No. SR/S1/IC-04/2007), for funding and CSIR for a fellowship to VB. The single-crystal X-ray diffractometer facility provided by IITD is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2531).

References

- Bagchi, V., Das, P. & Bandyopadhyay, D. (2007). *Acta Cryst.* **E63**, m1704.
 Bruker (1998). SMART. Version 5.054. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2000). SAINT. Version 6.02a. Bruker AXS Inc., Madison, Wisconsin, USA.
 Buraway, A. & Vellins, C. E. (1954). *J. Chem. Soc.* pp. 90–95.
 Chattopadhyay, S., Sinha, C., Basu, P. & Chakravorty, A. (1991). *Organometallics*, **10**, 1135–1139.
 Dupont, J., Consorti, C. S. & Spencer, J. (2005). *Chem. Rev.* **105**, 2527–2571.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m359 [doi:10.1107/S1600536808000767]

Chlorido{5-chloro-2-[2-(methylsulfanyl)phenyldiazenyl]phenyl}platinum(II)

V. Bagchi and D. Bandyopadhyay

Comment

Cyclometallated compounds have numerous applications (Dupont *et al.*, 2005) in organic synthesis, catalysis and metal-omesogen chemistry. Although a number of cycloplatinated complexes have been reported in the literature, report of platinum complexes with sulfur as auxiliary donor and the existence of C–Pt bond are sparse. Herein we report the crystal structure of (I) having such features.

The molecular structure of the title compound, (I), is shown in Fig. 1, with the atom numbering scheme. The platinum atom along with donor set of four atoms lie almost in one plane. Selected bond lengths, bond angles are listed in Table 1. The packing arrangement of (I) is shown in Fig. 2. The N=N bond length is similar of other cycloplatinated azoarenes (Chattopadhyay *et al.*, 1991).

The metal carbon bond length, 1.986 (3) Å, is slightly lower than the reported values of other *ortho*-metallated azoarenes (Chattopadhyay *et al.*, 1991). The molecules are found to dimerize through non-bonded S...S interaction; having S...Sⁱ [symmetry code: (i) $-x, 2 - y, -z$] distance of 3.523 (18) Å (Bagchi *et al.* 2007) (Fig.3) The crystal packing is stabilized by four inter-molecular π - π interactions; the Cg3—Cg4ⁱⁱ, Cg3—Cg4ⁱⁱⁱ, Cg4—Cg3ⁱⁱ, Cg4—Cg3ⁱⁱⁱ, [symmetry codes: (ii) $-x, 1 - y, -z$, (iii) $1 - x, 1 - y, -z$. Cg3 and Cg4 are the centroids of C1—C6 and C7—C12 rings, respectively] distances are 3.638 (3), 3.804 (3), 3.638 (3) and 3.804 (3) Å, respectively; the corresponding perpendicular distances are 3.448, 3.369, 3.466 and 3.406 Å, respectively (Fig. 4).

Experimental

2-(Methylsulfanyl)diazenyl-4-chlorobenzene was prepared by coupling 2-(methylsulfanyl)aniline with 4-nitroso-chlorobenzene. The ligand thus obtained was reacted with K₂PtCl₄ following a reported method (Buraway & Vellins, 1954). The product was purified by column chromatographic technique using silica gel column and methanol and dichloromethane (1:9 v/v) mixture as eluent. The solvent was evaporated in vacuum to obtain the pure product (78.3%). Suitable crystals of (I) were grown from a dichloromethane–hexane solution by slow evaporation.

Refinement

H atoms were included at calculated positions as riding atoms with C—H set to 0.93 Å for (aromatic) and 0.96 Å for (CH₃) H atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ [1.5 U_{eq} for methyl group].

Figures




Fig. 1. The asymmetric unit of (I), with displacement ellipsoids drawn at the 50% probability level.




Fig. 2. The molecular arrangement of (I) in the unitcell.




Fig. 3. The intermolecular S...S interaction for (I), indicated by dotted line [symmetry code: (i) $-x, 2 - y, -z$].




Fig. 4. The intermolecular π — π interactions for (I), indicated by the dotted lines. [Symmetry codes: (ii) $-x, 1 - y, -z$, (iii) $1 - x, 1 - y, -z$]. *Cg3* and *Cg4* are centroids of C1—C6 and C7—C12 rings, respectively.

Chlorido{5-chloro-2-[2-(methylsulfonyl)phenyldiazenyl]phenyl}platinum(II)

Crystal data

[Pt(C₁₃H₁₀ClN₂S)Cl]

$M_r = 492.28$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.424$ (2) Å

$b = 8.777$ (3) Å

$c = 11.069$ (3) Å

$\alpha = 105.428$ (4)°

$\beta = 91.798$ (4)°

$\gamma = 96.641$ (4)°

$V = 689.1$ (4) Å³

$Z = 2$

$F_{000} = 460.0$

$D_x = 2.372$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2472 reflections

$\theta = 1.0$ – 25.5 °

$\mu = 10.70$ mm⁻¹

$T = 273$ (2) K

Block, pink

$0.42 \times 0.28 \times 0.19$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

2544 independent reflections

Radiation source: fine-focus sealed tube	2472 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.025$
$T = 273(2)$ K	$\theta_{\text{max}} = 25.5^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -8 \rightarrow 8$
$T_{\text{min}} = 0.035$, $T_{\text{max}} = 0.134$	$k = -10 \rightarrow 10$
6783 measured reflections	$l = -13 \rightarrow 13$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.018$	H-atom parameters constrained
$wR(F^2) = 0.043$	$w = 1/[\sigma^2(F_o^2) + (0.0209P)^2 + 0.4377P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
2544 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
173 parameters	$\Delta\rho_{\text{max}} = 0.89 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.89 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C8	0.3562 (5)	0.2238 (4)	-0.1887 (4)	0.0399 (8)
H8	0.3666	0.1521	-0.1411	0.048*
C9	0.3852 (5)	0.1810 (4)	-0.3149 (4)	0.0432 (9)
H9	0.4144	0.0802	-0.3542	0.052*
Pt1	0.216802 (16)	0.685427 (14)	-0.091569 (11)	0.03068 (6)
Cl2	0.40740 (17)	0.23872 (14)	-0.54110 (10)	0.0599 (3)
Cl1	0.19823 (17)	0.81788 (12)	-0.24323 (10)	0.0529 (3)
N1	0.2327 (4)	0.5583 (4)	0.0285 (3)	0.0311 (6)
N2	0.2763 (4)	0.4162 (3)	-0.0062 (3)	0.0356 (6)
C1	0.1910 (4)	0.6209 (4)	0.1559 (3)	0.0322 (7)

supplementary materials

C7	0.3111 (4)	0.3755 (4)	-0.1328 (3)	0.0329 (7)
C10	0.3698 (5)	0.2924 (4)	-0.3816 (3)	0.0390 (8)
C2	0.1378 (5)	0.7727 (4)	0.1893 (3)	0.0347 (7)
C11	0.3267 (5)	0.4437 (4)	-0.3279 (3)	0.0370 (8)
H11	0.3186	0.5149	-0.3763	0.044*
C5	0.1648 (6)	0.6038 (5)	0.3660 (4)	0.0480 (10)
H5	0.1715	0.5467	0.4254	0.058*
S1	0.10822 (13)	0.87598 (11)	0.07262 (8)	0.0362 (2)
C12	0.2953 (5)	0.4887 (4)	-0.2005 (3)	0.0326 (7)
C4	0.1152 (6)	0.7564 (6)	0.4010 (4)	0.0515 (10)
H4	0.0924	0.8021	0.4841	0.062*
C6	0.2041 (5)	0.5359 (5)	0.2441 (4)	0.0412 (8)
H6	0.2389	0.4342	0.2213	0.049*
C3	0.0998 (5)	0.8407 (5)	0.3127 (4)	0.0454 (9)
H3	0.0643	0.9421	0.3358	0.054*
C13	0.2718 (6)	1.0531 (5)	0.1290 (4)	0.0519 (10)
H13A	0.2399	1.1140	0.2093	0.078*
H13B	0.2714	1.1164	0.0704	0.078*
H13C	0.3908	1.0227	0.1371	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C8	0.041 (2)	0.0332 (19)	0.048 (2)	0.0096 (15)	0.0056 (17)	0.0131 (17)
C9	0.043 (2)	0.0313 (19)	0.052 (2)	0.0099 (15)	0.0065 (18)	0.0042 (17)
Pt1	0.03822 (9)	0.02666 (9)	0.02684 (9)	0.00541 (6)	0.00145 (6)	0.00626 (6)
C12	0.0827 (8)	0.0549 (6)	0.0349 (5)	0.0134 (6)	0.0109 (5)	-0.0030 (5)
C11	0.0879 (8)	0.0395 (5)	0.0383 (5)	0.0191 (5)	0.0114 (5)	0.0174 (4)
N1	0.0315 (15)	0.0336 (16)	0.0279 (15)	0.0057 (12)	0.0014 (12)	0.0074 (12)
N2	0.0366 (15)	0.0343 (16)	0.0374 (17)	0.0074 (12)	0.0041 (13)	0.0112 (13)
C1	0.0314 (17)	0.0361 (18)	0.0274 (17)	0.0024 (14)	-0.0006 (13)	0.0069 (14)
C7	0.0318 (17)	0.0315 (18)	0.0350 (19)	0.0042 (14)	0.0018 (14)	0.0082 (15)
C10	0.0388 (19)	0.040 (2)	0.0324 (19)	0.0039 (15)	0.0053 (15)	0.0006 (16)
C2	0.0356 (18)	0.0370 (19)	0.0294 (18)	0.0031 (14)	0.0004 (14)	0.0065 (15)
C11	0.044 (2)	0.0319 (18)	0.0340 (19)	0.0031 (15)	0.0027 (16)	0.0081 (15)
C5	0.052 (2)	0.063 (3)	0.033 (2)	0.006 (2)	-0.0015 (17)	0.0205 (19)
S1	0.0434 (5)	0.0343 (5)	0.0302 (5)	0.0113 (4)	0.0005 (4)	0.0054 (4)
C12	0.0334 (17)	0.0279 (17)	0.0330 (19)	0.0020 (13)	-0.0012 (14)	0.0033 (14)
C4	0.049 (2)	0.073 (3)	0.031 (2)	0.008 (2)	0.0055 (17)	0.011 (2)
C6	0.043 (2)	0.046 (2)	0.037 (2)	0.0049 (16)	-0.0015 (16)	0.0164 (17)
C3	0.048 (2)	0.053 (2)	0.034 (2)	0.0100 (18)	0.0056 (17)	0.0071 (18)
C13	0.070 (3)	0.034 (2)	0.047 (2)	0.0003 (19)	-0.005 (2)	0.0070 (18)

Geometric parameters (\AA , $^\circ$)

C8—C9	1.378 (6)	C10—C11	1.382 (5)
C8—C7	1.394 (5)	C2—C3	1.390 (5)
C8—H8	0.9300	C2—S1	1.785 (4)
C9—C10	1.385 (6)	C11—C12	1.395 (5)

C9—H9	0.9300	C11—H11	0.9300
Pt1—N1	1.959 (3)	C5—C6	1.378 (6)
Pt1—C12	1.986 (3)	C5—C4	1.388 (6)
Pt1—C11	2.2911 (11)	C5—H5	0.9300
Pt1—S1	2.3529 (10)	S1—C13	1.811 (4)
Cl2—C10	1.742 (4)	C4—C3	1.383 (6)
N1—N2	1.286 (4)	C4—H4	0.9300
N1—C1	1.429 (4)	C6—H6	0.9300
N2—C7	1.391 (5)	C3—H3	0.9300
C1—C6	1.385 (5)	C13—H13A	0.9600
C1—C2	1.391 (5)	C13—H13B	0.9600
C7—C12	1.407 (5)	C13—H13C	0.9600
C9—C8—C7	119.5 (4)	C10—C11—C12	119.4 (3)
C9—C8—H8	120.3	C10—C11—H11	120.3
C7—C8—H8	120.3	C12—C11—H11	120.3
C8—C9—C10	118.2 (3)	C6—C5—C4	120.7 (4)
C8—C9—H9	120.9	C6—C5—H5	119.7
C10—C9—H9	120.9	C4—C5—H5	119.7
N1—Pt1—C12	79.26 (14)	C2—S1—C13	102.24 (19)
N1—Pt1—Cl1	175.94 (9)	C2—S1—Pt1	95.61 (12)
C12—Pt1—Cl1	96.84 (11)	C13—S1—Pt1	111.80 (16)
N1—Pt1—S1	86.25 (9)	C11—C12—C7	117.3 (3)
C12—Pt1—S1	165.07 (11)	C11—C12—Pt1	131.9 (3)
Cl1—Pt1—S1	97.57 (4)	C7—C12—Pt1	110.7 (3)
N2—N1—C1	118.9 (3)	C3—C4—C5	120.2 (4)
N2—N1—Pt1	120.9 (2)	C3—C4—H4	119.9
C1—N1—Pt1	120.2 (2)	C5—C4—H4	119.9
N1—N2—C7	110.8 (3)	C5—C6—C1	119.2 (4)
C6—C1—C2	120.7 (3)	C5—C6—H6	120.4
C6—C1—N1	121.9 (3)	C1—C6—H6	120.4
C2—C1—N1	117.4 (3)	C4—C3—C2	119.5 (4)
N2—C7—C8	119.2 (3)	C4—C3—H3	120.2
N2—C7—C12	118.3 (3)	C2—C3—H3	120.2
C8—C7—C12	122.4 (3)	S1—C13—H13A	109.5
C11—C10—C9	123.3 (3)	S1—C13—H13B	109.5
C11—C10—Cl2	118.5 (3)	H13A—C13—H13B	109.5
C9—C10—Cl2	118.2 (3)	S1—C13—H13C	109.5
C3—C2—C1	119.7 (3)	H13A—C13—H13C	109.5
C3—C2—S1	120.1 (3)	H13B—C13—H13C	109.5
C1—C2—S1	120.1 (3)		
C7—C8—C9—C10	0.6 (6)	C1—C2—S1—Pt1	-6.0 (3)
C12—Pt1—N1—N2	-1.8 (3)	N1—Pt1—S1—C2	4.51 (14)
S1—Pt1—N1—N2	174.6 (3)	C12—Pt1—S1—C2	18.4 (4)
C12—Pt1—N1—C1	179.9 (3)	Cl1—Pt1—S1—C2	-176.89 (12)
S1—Pt1—N1—C1	-3.7 (2)	N1—Pt1—S1—C13	110.10 (18)
C1—N1—N2—C7	179.6 (3)	C12—Pt1—S1—C13	124.0 (4)
Pt1—N1—N2—C7	1.4 (4)	Cl1—Pt1—S1—C13	-71.30 (16)
N2—N1—C1—C6	3.0 (5)	C10—C11—C12—C7	0.5 (5)

supplementary materials

Pt1—N1—C1—C6	-178.7 (3)	C10—C11—C12—Pt1	-175.7 (3)
N2—N1—C1—C2	-177.6 (3)	N2—C7—C12—C11	-178.5 (3)
Pt1—N1—C1—C2	0.6 (4)	C8—C7—C12—C11	0.1 (5)
N1—N2—C7—C8	-178.4 (3)	N2—C7—C12—Pt1	-1.6 (4)
N1—N2—C7—C12	0.2 (4)	C8—C7—C12—Pt1	177.0 (3)
C9—C8—C7—N2	178.0 (3)	N1—Pt1—C12—C11	178.0 (4)
C9—C8—C7—C12	-0.6 (5)	C11—Pt1—C12—C11	-0.8 (3)
C8—C9—C10—C11	0.0 (6)	S1—Pt1—C12—C11	163.9 (2)
C8—C9—C10—C12	179.7 (3)	N1—Pt1—C12—C7	1.7 (2)
C6—C1—C2—C3	0.9 (5)	C11—Pt1—C12—C7	-177.2 (2)
N1—C1—C2—C3	-178.5 (3)	S1—Pt1—C12—C7	-12.4 (6)
C6—C1—C2—S1	-176.2 (3)	C6—C5—C4—C3	1.8 (6)
N1—C1—C2—S1	4.4 (4)	C4—C5—C6—C1	-1.0 (6)
C9—C10—C11—C12	-0.5 (6)	C2—C1—C6—C5	-0.4 (5)
C12—C10—C11—C12	179.7 (3)	N1—C1—C6—C5	179.0 (3)
C3—C2—S1—C13	63.1 (4)	C5—C4—C3—C2	-1.3 (6)
C1—C2—S1—C13	-119.8 (3)	C1—C2—C3—C4	-0.1 (6)
C3—C2—S1—Pt1	176.9 (3)	S1—C2—C3—C4	177.0 (3)

Fig. 1




Fig. 2




Fig. 3




Fig. 4

