

Aquabis(2-chloroacetato- κO)(1,10-phenanthroline- $\kappa^2 N,N'$)copper(II)

Rongdong Yang,* Yanfei Li, Junshan Sun, Jikun Li and Changqing Chu

Department of Materials Science, and Chemical Engineering, Taishan University, 271021 Tai'an, Shandong, People's Republic of China
Correspondence e-mail: klsz79@163.com


Received 29 November 2007; accepted 1 January 2008

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.026; wR factor = 0.075; data-to-parameter ratio = 13.0.

In the title complex, $[\text{Cu}(\text{C}_2\text{H}_2\text{ClO}_2)_2(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})]$, the Cu^{II} ion is five-coordinated by two N atoms [$\text{Cu}-\text{N} = 2.005(2)$ and $2.029(2)\text{ \AA}$] from the 1,10-phenanthroline ligand, two O atoms [$\text{Cu}-\text{O} = 1.943(2)$ – $1.966(2)\text{ \AA}$] from two 2-chloroacetate ligands and one water molecule [$\text{Cu}-\text{O} = 2.253(2)\text{ \AA}$] in a distorted square-pyramidal geometry. The crystal structure exhibits intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, short $\text{Cl}\cdots\text{Cl}$ contacts [$3.334(1)\text{ \AA}$] and $\pi-\pi$ interactions [centroid–centroid distance $3.621(11)\text{ \AA}$].

Related literature

For related crystal structures, see: Sieroń (2007); Czylkowska *et al.* (2004); Chen *et al.* (1996); Overgaard *et al.* (2003).



Experimental

Crystal data

$[\text{Cu}(\text{C}_2\text{H}_2\text{ClO}_2)_2(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})]$	$\gamma = 97.9190(10)^\circ$
$M_r = 448.73$	$V = 869.66(11)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 8.7730(6)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.2382(7)\text{ \AA}$	$\mu = 1.59\text{ mm}^{-1}$
$c = 11.4492(8)\text{ \AA}$	$T = 273(2)\text{ K}$
$\alpha = 96.2180(10)^\circ$	$0.38 \times 0.25 \times 0.19\text{ mm}$
$\beta = 106.6760(10)^\circ$	

Data collection

Bruker SMART CCD area detector diffractometer	4610 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	3057 independent reflections
$R_{\text{int}} = 0.015$	2837 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.583$, $T_{\max} = 0.752$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$	3 restraints
$wR(F^2) = 0.074$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.30\text{ e \AA}^{-3}$
3057 reflections	$\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$
235 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5—H15 \cdots O4 ⁱ	0.85	1.96	2.796 (2)	169

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 1997); software used to prepare material for publication: *SHELXTL*.

The authors thank the Postgraduate Foundation of Taishan University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2370).

References

- Chen, X. M., Tong, M. L., Wu, Y. L. & Luo, Y. J. (1996). *J. Chem. Soc. Dalton Trans.*, pp. 2181–2182.
- Czylkowska, A., Kruszynski, R., Czakis-Sulikowska, D. & Bartczak, T. J. (2004). *J. Coord. Chem.* **57**, 239–249.
- Overgaard, J., Larsen, F. K., Schiott, B. & Lversen, B. B. (2003). *J. Am. Chem. Soc.* **125**, 11088–11098.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997). *SHELXTL*. Version 5.1. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Systems Inc., Madison, Wisconsin, USA.
- Sieroń, L. (2007). *Acta Cryst. E* **63**, m1659–m1661.

supporting information

Acta Cryst. (2008). E64, m348 [doi:10.1107/S1600536808000044]

Aquabis(2-chloroacetato- κO)(1,10-phenanthroline- $\kappa^2 N,N'$)copper(II)

Rongdong Yang, Yanfei Li, Junshan Sun, Jikun Li and Changqing Chu

S1. Comment

2-Chloroacetic acid and its derivatives are often used in the synthesis of mononuclear monomeric (Sieroń, 2007; Czylkowska *et al.*, 2004) and polymeric compounds (Chen *et al.*, 1996; Overgaard *et al.*, 2003). In our search for new topological structures, we selected the copper(II) ion with 2-chloroacetic acid in the presence of 1,10-phenanthroline as a co-ligand, and obtained the title compound, (I).

In (I) (Fig. 1), Cu1 exhibits a five-coordinated square-pyramidal environment, formed by two O atoms from two carboxyl ligands (Cu1—O1 1.943 (2) Å, Cu1—O3 1.966 (2) Å), one water molecule (Cu1—O5 2.243 (5) Å) and two N atoms (Cu1—N1 2.029 (2) Å, Cu1—N2 2.005 (2) Å) from 1,10-phenanthroline ligand.

In the crystal structure, there exist short intermolecular Cl···Cl contacts (Table 1), $\pi\cdots\pi$ stacking interactions between the aromatic rings from neighbouring molecules (Table 1), and intermolecular O—H···O hydrogen bonds (Table 2), which link the molecules into centrosymmetric dimers.


S2. Experimental

The reaction was carried out by the solvothermal method. 2-Chloroacetic acid (0.188 g, 2 mmol) and cupric acetate (0.199 g, 1 mmol) and 1,10-phenanthroline (0.180 g, 1 mmol) were added to the airtight vessel with 20 ml water. The resulting green solution was filtered. The filtrate was placed for several days yielding blue block-shaped crystals.

The yield is 81% and elemental analysis: calc. for $C_{16}H_{14}Cl_2CuN_2O_5$: C 42.82, H 3.14, N 6.24; found: C 42.55, H 3.39, N 6.32. The elemental analyses were performed with PERKIN ELMER MODEL 2400 SERIES II.

S3. Refinement


All H atoms were found in Fourier difference map, but placed in idealized positions (C—H 0.93–0.97 Å, O—H 0.85 Å), with $U_{\text{iso}}(\text{H})=1.2U_{\text{eq}}$ of the parent atom.

**Figure 1**

The molecular structure of (I) with atomic numbering and 30% probability displacement ellipsoids.

Aquabis(2-chloroacetato- κ O)(1,10-phenanthroline- κ^2 N,N')copper(II)

Crystal data



$M_r = 448.73$

Triclinic, $P\bar{1}$

$a = 8.7730 (6)$ Å

$b = 9.2382 (7)$ Å

$c = 11.4492 (8)$ Å

$\alpha = 96.218 (1)^\circ$

$\beta = 106.676 (1)^\circ$

$\gamma = 97.919 (1)^\circ$

$V = 869.66 (11)$ Å³

$Z = 2$

$F(000) = 454$

$D_x = 1.714 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3430 reflections

$\theta = 2.5\text{--}28.2^\circ$

$\mu = 1.59 \text{ mm}^{-1}$

$T = 273$ K

Block, blue

$0.38 \times 0.25 \times 0.19$ mm

Data collection

Bruker SMART CCD area detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.583$, $T_{\max} = 0.752$

4610 measured reflections

3057 independent reflections

2837 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\max} = 25.1^\circ$, $\theta_{\min} = 1.9^\circ$

$h = -10 \rightarrow 10$

$k = -8 \rightarrow 10$

$l = -13 \rightarrow 13$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.026$$

$$wR(F^2) = 0.074$$

$$S = 1.00$$

3057 reflections

235 parameters

3 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.043P)^2 + 0.4807P]$$
$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.30 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.29 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.23816 (3)	0.68495 (3)	0.71095 (2)	0.03304 (10)
Cl1	-0.22927 (8)	0.98116 (9)	0.83337 (8)	0.0671 (2)
Cl2	0.40479 (7)	0.93989 (7)	0.62804 (6)	0.04661 (16)
O1	0.14841 (19)	0.84228 (17)	0.77963 (15)	0.0421 (4)
O2	-0.1084 (2)	0.7271 (2)	0.7366 (2)	0.0635 (5)
O3	0.11593 (19)	0.68927 (16)	0.53849 (14)	0.0408 (4)
O4	0.0053 (2)	0.7855 (2)	0.37496 (17)	0.0622 (5)
O5	0.06637 (19)	0.50494 (17)	0.74928 (15)	0.0430 (4)
H15	0.0312	0.4180	0.7091	0.052*
H16	-0.0123	0.5517	0.7385	0.052*
N1	0.4251 (2)	0.70913 (19)	0.86886 (16)	0.0328 (4)
N2	0.3550 (2)	0.53104 (19)	0.65745 (16)	0.0334 (4)
C1	0.0003 (3)	0.8353 (2)	0.77567 (19)	0.0365 (5)
C2	-0.0300 (3)	0.9841 (3)	0.8242 (2)	0.0400 (5)
H2A	-0.0099	1.0543	0.7708	0.048*
H2B	0.0464	1.0189	0.9057	0.048*
C3	0.1023 (3)	0.7947 (2)	0.4784 (2)	0.0366 (5)
C4	0.2075 (3)	0.9464 (3)	0.5315 (2)	0.0488 (6)
H4A	0.1539	1.0031	0.5788	0.059*
H4B	0.2171	0.9981	0.4639	0.059*
C5	0.4552 (3)	0.7987 (3)	0.9743 (2)	0.0404 (5)
H5A	0.3851	0.8640	0.9805	0.049*
C6	0.5887 (3)	0.7988 (3)	1.0767 (2)	0.0486 (6)
H6	0.6071	0.8642	1.1490	0.058*
C7	0.6915 (3)	0.7028 (3)	1.0699 (2)	0.0469 (6)

H7	0.7797	0.7014	1.1380	0.056*
C8	0.6639 (3)	0.6057 (3)	0.9596 (2)	0.0388 (5)
C9	0.7622 (3)	0.4983 (3)	0.9421 (2)	0.0479 (6)
H9	0.8521	0.4911	1.0067	0.057*
C10	0.7268 (3)	0.4079 (3)	0.8339 (3)	0.0477 (6)
H10	0.7934	0.3401	0.8249	0.057*
C11	0.5883 (3)	0.4140 (2)	0.7322 (2)	0.0387 (5)
C12	0.5395 (3)	0.3200 (3)	0.6177 (2)	0.0453 (6)
H12	0.6011	0.2500	0.6028	0.054*
C13	0.4019 (3)	0.3317 (3)	0.5290 (2)	0.0460 (6)
H13	0.3688	0.2690	0.4537	0.055*
C14	0.3111 (3)	0.4378 (2)	0.5512 (2)	0.0390 (5)
H14	0.2168	0.4439	0.4901	0.047*
C15	0.4909 (2)	0.5181 (2)	0.74641 (19)	0.0319 (4)
C16	0.5289 (2)	0.6148 (2)	0.86151 (19)	0.0317 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cu1	0.03296 (16)	0.03095 (16)	0.03298 (16)	0.00803 (11)	0.00661 (11)	0.00203 (10)
Cl1	0.0439 (4)	0.0748 (5)	0.0858 (5)	0.0241 (3)	0.0227 (3)	0.0012 (4)
Cl2	0.0413 (3)	0.0447 (3)	0.0484 (3)	-0.0015 (2)	0.0122 (3)	-0.0006 (3)
O1	0.0360 (8)	0.0378 (8)	0.0516 (9)	0.0094 (7)	0.0135 (7)	-0.0010 (7)
O2	0.0409 (10)	0.0442 (10)	0.0972 (16)	0.0042 (8)	0.0163 (10)	-0.0071 (10)
O3	0.0474 (9)	0.0287 (8)	0.0373 (8)	0.0040 (7)	-0.0002 (7)	0.0058 (6)
O4	0.0747 (13)	0.0462 (10)	0.0453 (10)	0.0019 (9)	-0.0113 (9)	0.0129 (8)
O5	0.0435 (9)	0.0342 (8)	0.0493 (9)	0.0040 (7)	0.0130 (7)	0.0051 (7)
N1	0.0344 (9)	0.0302 (9)	0.0328 (9)	0.0031 (7)	0.0105 (7)	0.0032 (7)
N2	0.0359 (9)	0.0312 (9)	0.0328 (9)	0.0052 (7)	0.0109 (8)	0.0037 (7)
C1	0.0357 (12)	0.0380 (12)	0.0352 (11)	0.0099 (10)	0.0078 (9)	0.0069 (9)
C2	0.0376 (12)	0.0416 (12)	0.0415 (12)	0.0122 (10)	0.0117 (10)	0.0041 (10)
C3	0.0399 (12)	0.0330 (11)	0.0340 (11)	0.0085 (9)	0.0069 (9)	0.0025 (9)
C4	0.0540 (15)	0.0334 (12)	0.0501 (14)	0.0061 (11)	0.0024 (12)	0.0077 (10)
C5	0.0457 (13)	0.0362 (12)	0.0372 (12)	0.0049 (10)	0.0129 (10)	-0.0011 (9)
C6	0.0548 (15)	0.0482 (14)	0.0327 (12)	-0.0027 (12)	0.0063 (11)	-0.0022 (10)
C7	0.0418 (13)	0.0490 (14)	0.0390 (12)	-0.0038 (11)	-0.0005 (10)	0.0114 (11)
C8	0.0344 (11)	0.0398 (12)	0.0409 (12)	0.0010 (9)	0.0090 (9)	0.0142 (10)
C9	0.0337 (12)	0.0546 (15)	0.0570 (15)	0.0110 (11)	0.0099 (11)	0.0230 (12)
C10	0.0400 (13)	0.0467 (14)	0.0657 (16)	0.0188 (11)	0.0217 (12)	0.0192 (12)
C11	0.0396 (12)	0.0326 (11)	0.0516 (13)	0.0075 (9)	0.0231 (10)	0.0129 (10)
C12	0.0530 (15)	0.0319 (12)	0.0609 (15)	0.0109 (10)	0.0316 (13)	0.0060 (10)
C13	0.0582 (15)	0.0329 (12)	0.0474 (14)	0.0002 (11)	0.0247 (12)	-0.0048 (10)
C14	0.0436 (12)	0.0339 (11)	0.0363 (11)	0.0008 (9)	0.0121 (10)	0.0005 (9)
C15	0.0326 (11)	0.0288 (10)	0.0370 (11)	0.0040 (8)	0.0145 (9)	0.0080 (8)
C16	0.0302 (10)	0.0313 (10)	0.0341 (11)	0.0024 (8)	0.0107 (9)	0.0091 (8)

Geometric parameters (\AA , $\text{^{\circ}}$)

Cu1—O1	1.9427 (15)	C4—H4A	0.9700
Cu1—O3	1.9657 (15)	C4—H4B	0.9700
Cu1—N2	2.0052 (18)	C5—C6	1.399 (3)
Cu1—N1	2.0294 (18)	C5—H5A	0.9300
Cu1—O5	2.2531 (16)	C6—C7	1.361 (4)
Cl1—C2	1.778 (2)	C6—H6	0.9300
Cl1—Cl2 ⁱ	3.3340 (10)	C7—C8	1.406 (3)
Cl2—C4	1.779 (3)	C7—H7	0.9300
O1—C1	1.279 (3)	C8—C16	1.398 (3)
O2—C1	1.226 (3)	C8—C9	1.435 (3)
O3—C3	1.251 (3)	C9—C10	1.346 (4)
O4—C3	1.231 (3)	C9—H9	0.9300
O5—H15	0.8498	C10—C11	1.434 (3)
O5—H16	0.8498	C10—H10	0.9300
N1—C5	1.324 (3)	C11—C15	1.398 (3)
N1—C16	1.357 (3)	C11—C12	1.408 (3)
N2—C14	1.336 (3)	C12—C13	1.363 (4)
N2—C15	1.357 (3)	C12—H12	0.9300
C1—C2	1.514 (3)	C13—C14	1.392 (3)
C2—H2A	0.9700	C13—H13	0.9300
C2—H2B	0.9700	C14—H14	0.9300
C3—C4	1.524 (3)	C15—C16	1.434 (3)
Cl1…Cl2 ⁱ	3.334 (1)	Cg1…Cg2 ⁱⁱ	3.621 (11)
O1—Cu1—O3	94.93 (7)	H4A—C4—H4B	107.7
O1—Cu1—N2	173.08 (7)	N1—C5—C6	122.4 (2)
O3—Cu1—N2	91.04 (7)	N1—C5—H5A	118.8
O1—Cu1—N1	91.67 (7)	C6—C5—H5A	118.8
O3—Cu1—N1	160.92 (7)	C7—C6—C5	119.7 (2)
N2—Cu1—N1	81.58 (7)	C7—C6—H6	120.2
O1—Cu1—O5	93.30 (6)	C5—C6—H6	120.2
O3—Cu1—O5	98.18 (6)	C6—C7—C8	119.7 (2)
N2—Cu1—O5	89.32 (7)	C6—C7—H7	120.1
N1—Cu1—O5	99.28 (7)	C8—C7—H7	120.1
C1—O1—Cu1	125.47 (14)	C16—C8—C7	116.6 (2)
C3—O3—Cu1	130.63 (14)	C16—C8—C9	118.6 (2)
Cu1—O5—H15	127.2	C7—C8—C9	124.8 (2)
Cu1—O5—H16	95.7	C10—C9—C8	121.3 (2)
H15—O5—H16	108.2	C10—C9—H9	119.3
C5—N1—C16	117.97 (19)	C8—C9—H9	119.3
C5—N1—Cu1	129.49 (16)	C9—C10—C11	121.2 (2)
C16—N1—Cu1	112.52 (13)	C9—C10—H10	119.4
C14—N2—C15	118.05 (19)	C11—C10—H10	119.4
C14—N2—Cu1	128.50 (16)	C15—C11—C12	116.5 (2)
C15—N2—Cu1	113.32 (13)	C15—C11—C10	118.7 (2)

O2—C1—O1	127.3 (2)	C12—C11—C10	124.8 (2)
O2—C1—C2	121.8 (2)	C13—C12—C11	119.9 (2)
O1—C1—C2	110.88 (19)	C13—C12—H12	120.0
C1—C2—Cl1	113.88 (16)	C11—C12—H12	120.0
C1—C2—H2A	108.8	C12—C13—C14	119.9 (2)
Cl1—C2—H2A	108.8	C12—C13—H13	120.0
C1—C2—H2B	108.8	C14—C13—H13	120.0
Cl1—C2—H2B	108.8	N2—C14—C13	122.0 (2)
H2A—C2—H2B	107.7	N2—C14—H14	119.0
O4—C3—O3	123.8 (2)	C13—C14—H14	119.0
O4—C3—C4	115.3 (2)	N2—C15—C11	123.6 (2)
O3—C3—C4	120.9 (2)	N2—C15—C16	116.28 (18)
C3—C4—Cl2	113.92 (17)	C11—C15—C16	120.1 (2)
C3—C4—H4A	108.8	N1—C16—C8	123.6 (2)
Cl2—C4—H4A	108.8	N1—C16—C15	116.28 (18)
C3—C4—H4B	108.8	C8—C16—C15	120.1 (2)
Cl2—C4—H4B	108.8		
O3—Cu1—O1—C1	65.71 (18)	C5—C6—C7—C8	0.8 (4)
N2—Cu1—O1—C1	-144.9 (5)	C6—C7—C8—C16	0.0 (3)
N1—Cu1—O1—C1	-132.21 (18)	C6—C7—C8—C9	-179.0 (2)
O5—Cu1—O1—C1	-32.81 (18)	C16—C8—C9—C10	0.2 (3)
O1—Cu1—O3—C3	52.0 (2)	C7—C8—C9—C10	179.2 (2)
N2—Cu1—O3—C3	-124.4 (2)	C8—C9—C10—C11	-0.7 (4)
N1—Cu1—O3—C3	-57.7 (3)	C9—C10—C11—C15	1.0 (3)
O5—Cu1—O3—C3	146.1 (2)	C9—C10—C11—C12	-177.6 (2)
O1—Cu1—N1—C5	2.6 (2)	C15—C11—C12—C13	-1.2 (3)
O3—Cu1—N1—C5	112.9 (2)	C10—C11—C12—C13	177.4 (2)
N2—Cu1—N1—C5	-178.9 (2)	C11—C12—C13—C14	0.7 (4)
O5—Cu1—N1—C5	-91.01 (19)	C15—N2—C14—C13	-1.5 (3)
O1—Cu1—N1—C16	-179.12 (14)	Cu1—N2—C14—C13	-177.15 (17)
O3—Cu1—N1—C16	-68.8 (3)	C12—C13—C14—N2	0.7 (4)
N2—Cu1—N1—C16	-0.65 (14)	C14—N2—C15—C11	0.9 (3)
O5—Cu1—N1—C16	87.26 (14)	Cu1—N2—C15—C11	177.24 (16)
O1—Cu1—N2—C14	-170.8 (5)	C14—N2—C15—C16	-176.64 (18)
O3—Cu1—N2—C14	-21.29 (19)	Cu1—N2—C15—C16	-0.3 (2)
N1—Cu1—N2—C14	176.37 (19)	C12—C11—C15—N2	0.4 (3)
O5—Cu1—N2—C14	76.88 (19)	C10—C11—C15—N2	-178.3 (2)
O1—Cu1—N2—C15	13.4 (6)	C12—C11—C15—C16	177.90 (19)
O3—Cu1—N2—C15	162.87 (14)	C10—C11—C15—C16	-0.8 (3)
N1—Cu1—N2—C15	0.53 (14)	C5—N1—C16—C8	1.0 (3)
O5—Cu1—N2—C15	-98.96 (14)	Cu1—N1—C16—C8	-177.50 (16)
Cu1—O1—C1—O2	5.6 (3)	C5—N1—C16—C15	179.16 (18)
Cu1—O1—C1—C2	-172.99 (14)	Cu1—N1—C16—C15	0.7 (2)
O2—C1—C2—Cl1	6.1 (3)	C7—C8—C16—N1	-1.0 (3)
O1—C1—C2—Cl1	-175.31 (16)	C9—C8—C16—N1	178.1 (2)
Cu1—O3—C3—O4	-170.39 (19)	C7—C8—C16—C15	-179.08 (19)
Cu1—O3—C3—C4	9.1 (3)	C9—C8—C16—C15	0.0 (3)

O4—C3—C4—Cl2	−147.4 (2)	N2—C15—C16—N1	−0.2 (3)
O3—C3—C4—Cl2	33.1 (3)	C11—C15—C16—N1	−177.89 (18)
C16—N1—C5—C6	−0.1 (3)	N2—C15—C16—C8	178.01 (18)
Cu1—N1—C5—C6	178.14 (17)	C11—C15—C16—C8	0.3 (3)
N1—C5—C6—C7	−0.8 (4)		

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, -y+1, -z+2$.

Hydrogen-bond geometry (\AA , °)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5—H15···O4 ⁱⁱⁱ	0.85	1.96	2.796 (2)	169

Symmetry code: (iii) $-x, -y+1, -z+1$.