

Monoclinic form of 1,2,4,5-tetracyclohexylbenzene

Joel T. Mague,* Lisa Linhardt, Iliana Medina,
Daniel J. Sattler and Mark J. Fink

Department of Chemistry, Tulane University, New Orleans, LA 70118, USA

Correspondence e-mail: joelt@tulane.edu


Received 13 December 2007; accepted 18 December 2007

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.046; wR factor = 0.125; data-to-parameter ratio = 20.6.

The molecule of the title compound, $C_{30}H_{46}$, has a crystallographically imposed inversion center and the cyclohexyl groups are oriented with their methine H atoms pointing towards one another ($\text{H}\cdots\text{H} = 1.99\text{ \AA}$). The cyclohexyl groups adopt chair conformations. A significant $\text{C}-\text{H}\cdots\pi$ interaction assembles molecules into layers parallel to (100).

Related literature

For related structures, see: Mague *et al.* (2008a,b); Vilardo *et al.* (2000). For related literature, see: Koudelka *et al.* (1985); Saito *et al.* (2004); Schweiger *et al.* (2001).



Experimental

Crystal data

$C_{30}H_{46}$	$V = 1213.99(14)\text{ \AA}^3$
$M_r = 406.67$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 10.3868(7)\text{ \AA}$	$\mu = 0.06\text{ mm}^{-1}$
$b = 10.1434(7)\text{ \AA}$	$T = 100(2)\text{ K}$
$c = 11.5419(8)\text{ \AA}$	$0.24 \times 0.21 \times 0.11\text{ mm}$
$\beta = 93.314(1)^\circ$	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	10421 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2002)	2805 independent reflections
$R_{\text{int}} = 0.019$	2460 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.975$, $T_{\max} = 0.993$	

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$	136 parameters
$wR(F^2) = 0.126$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 0.39\text{ e \AA}^{-3}$
2805 reflections	$\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

Cg is the centroid of the aromatic ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C6-\text{H}6A\cdots Cg^i$	0.99	2.62	3.520 (2)	150

Symmetry code: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT-Plus* (Bruker 2004); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Bruker, 2000); software used to prepare material for publication: *SHELXTL*.

We thank the Chemistry Department of Tulane University for support of the X-ray laboratory, and the Louisiana Board of Regents through the Louisiana Educational Quality Support Fund [grant LEQSF (2003–2003)-ENH-TR-67] for the purchase of the APEX diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2127).

References

- Bruker (2000). *SMART* (Version 5.625) and *SHELXTL* (Version 6.10). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *SAINT-Plus*. Version 7.03. Bruker AXS Inc., Madison, Wisconsin, USA.
- Koudelka, J., Saman, J. & Exner, O. (1985). *Collect. Czech. Chem. Commun.* **50**, 208–214.
- Mague, J. T., Linhardt, L., Medina, I. & Fink, M. J. (2008a). *Acta Cryst. E64*, o376.
- Mague, J. T., Linhardt, L., Medina, I. & Fink, M. J. (2008b). *Acta Cryst. E64*, o335.
- Saito, M., Tokitoh, N. & Okazaki, R. (2004). *J. Am. Chem. Soc.* **126**, 15572–15582.
- Schweiger, S. W., Salberg, M. M., Pulvirenti, A. L., Freeman, E. E., Fanwick, P. E. & Rothwell, I. P. (2001). *J. Chem. Soc. Dalton Trans.* pp. 2020–2031.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2002). *SADABS*, Version 2.05. University of Göttingen, Germany.
- Vilardo, J. S., Salberg, M. M., Parker, J. R., Fanwick, P. E. & Rothwell, I. P. (2000). *Inorg. Chim. Acta*, **299**, 135–141.

supporting information

Acta Cryst. (2008). E64, o375 [doi:10.1107/S160053680706758X]

Monoclinic form of 1,2,4,5-tetracyclohexylbenzene

Joel T. Mague, Lisa Linhardt, Iliana Medina, Daniel J. Sattler and Mark J. Fink

S1. Comment


Poly(cyclohexyl)benzenes have been used in the synthesis of a variety of sterically bulky protecting groups (Koudelka *et al.*, 1985); Saito *et al.*, 2004; Vilardo *et al.*, 2000; Schweiger *et al.*, 2001). Crystallization of 1,2,4,5-tetracyclohexylbenzene ($C_{30}H_{46}$) from hot methylcyclohexane forms colorless needle-shaped crystals together with a smaller quantity having a distinctly different block-shaped morphology. The latter is monoclinic with the molecule having crystallographically imposed centrosymmetry. The cyclohexyl rings adopt the chair conformation and are oriented with their methine hydrogen atoms pointed towards one another ($H\cdots H$ distance 1.99 Å). This contrasts with the structure of 1-bromo-2,4,6-tricyclohexylbenzene (Mague *et al.*, 2008b) where the methine hydrogen atoms of the *meta*-disposed cyclohexyl groups point towards the intervening ring substituent. This is presumably to minimize intramolecular contacts between the *ortho*-disposed cyclohexyl rings. Indeed, there are very few close contacts involving these substituents, the shortest being $H4\cdots H10$ (1.99 Å), $H3\cdots H5a$ (2.18 Å) and $H3\cdots H11a'$ (2.07 Å). The plane defined by the atoms C5, C6, C8, C9 ("seat" of the chair) is inclined to the plane of the aromatic ring by 79.0 (2)° while that for the other cyclohexyl ring (C11, C12, C14, C15) is inclined at an angle of only 61.7 (2)°. A significant C—H $\cdots\pi$ interaction occurs between C6—H6A and the center of gravity (C_g) of the aromatic ring in the molecule at $1 - x, 1/2 + y, 0.5 - z$ where the H— C_g distance is 2.62 Å and the C—H $\cdots C_g$ angle is 150°. This interaction forms layers of molecules parallel to (100) at approximately $x = 0.5$.

S2. Experimental

A mixture of chlorocyclohexane (125 ml, 1.05 mol) and benzene (9 ml, 0.1 mol) in a 250 ml 3-necked flask was cooled to -40° C and mechanically stirred while anhydrous AlCl₃ (6.6 g, 0.05 mol) was added in portions over a 20 min. period. The mixture was allowed to slowly warm to -15° C and the stirring continued for 2 h. The resulting yellow-orange mixture was quenched by pouring it over 800 g of ice and, after thawing, was filtered. The organic layer of the filtrate was separated off, washed several times with water and reduced to a small volume under reduced pressure. After standing for 1 week, a precipitate formed which was filtered off, washed with 10% aqueous HCl and collected by filtration to provide 16.0 g (40%) of white solid (*M.p.* 549–550 K, bulk sample). ¹H NMR (δ , CDCl₃): 1.4 (20*H*, br multiplet), 1.8 (20*H*, br multiplet), 2.7 (4*H*, multiplet), 7.2 (2*H*, s). ¹³C NMR (δ , CDCl₃): 26.59, 27.56, 34.97, 123.25, 141.8.

S3. Refinement

H-atoms were placed in calculated positions (C—H = 0.95 - 0.98 Å) and refined as riding on their carriers with isotropic displacement parameters 1.2 - 1.5 times those of the attached carbon atoms.

**Figure 1**

Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level and H-atoms are represented by spheres of arbitrary radius. Primed atoms are related to unprimed atoms by the symmetry operation $0.5 - x, -y, -z$

1,2,4,5-tetracyclohexylbenzene

Crystal data

$C_{30}H_{46}$
 $M_r = 406.67$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 10.3868 (7)$ Å
 $b = 10.1434 (7)$ Å
 $c = 11.5419 (8)$ Å
 $\beta = 93.314 (1)^\circ$
 $V = 1213.99 (14)$ Å³
 $Z = 2$

$F(000) = 452$
 $D_x = 1.113 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 6595 reflections
 $\theta = 2.7\text{--}28.3^\circ$
 $\mu = 0.06 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
Block, colorless
 $0.24 \times 0.21 \times 0.11 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 0 pixels mm⁻¹

ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 2002)
 $T_{\min} = 0.975$, $T_{\max} = 0.993$
10421 measured reflections

2805 independent reflections
 2460 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$
 $\theta_{\text{max}} = 27.6^\circ, \theta_{\text{min}} = 2.0^\circ$

$h = -13 \rightarrow 13$
 $k = -13 \rightarrow 13$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.126$
 $S = 1.04$
 2805 reflections
 136 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0727P)^2 + 0.3565P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.39 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.21 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The diffraction data were collected in three sets of 606 frames (ω scans, $0.3^\circ/\text{scan}$) at φ settings of 0, 120 and 240° .

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. H-atoms were placed in calculated positions ($C-H = 0.95 - 0.98 \text{ \AA}$) and included as riding contributions with isotropic displacement parameters 1.2 - 1.5 times those of the attached carbon atoms.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.62897 (10)	-0.01942 (10)	0.04453 (8)	0.0136 (2)
C2	0.57336 (10)	0.10675 (10)	0.04764 (9)	0.0144 (2)
C3	0.44681 (10)	0.12214 (10)	0.00191 (9)	0.0152 (2)
H3	0.4101	0.2079	0.0025	0.018*
C4	0.64402 (10)	0.22668 (10)	0.09843 (9)	0.0147 (2)
H4	0.7248	0.1947	0.1408	0.018*
C5	0.56481 (11)	0.30053 (11)	0.18625 (10)	0.0202 (2)
H5A	0.4827	0.3308	0.1472	0.024*
H5B	0.5438	0.2396	0.2494	0.024*
C6	0.63818 (11)	0.41900 (11)	0.23785 (10)	0.0207 (3)
H6A	0.5833	0.4663	0.2915	0.025*
H6B	0.7165	0.3882	0.2829	0.025*
C7	0.67666 (12)	0.51267 (11)	0.14301 (10)	0.0225 (3)
H7A	0.7280	0.5861	0.1783	0.027*
H7B	0.5981	0.5506	0.1034	0.027*
C8	0.75542 (11)	0.44235 (12)	0.05420 (10)	0.0228 (3)
H8A	0.8390	0.4141	0.0918	0.027*
H8B	0.7734	0.5043	-0.0092	0.027*

C9	0.68369 (11)	0.32171 (11)	0.00324 (9)	0.0185 (2)
H9A	0.7399	0.2747	-0.0495	0.022*
H9B	0.6057	0.3512	-0.0429	0.022*
C10	0.76656 (9)	-0.04649 (10)	0.09169 (9)	0.0142 (2)
H10	0.8166	0.0370	0.0840	0.017*
C11	0.83559 (10)	-0.15393 (11)	0.02505 (10)	0.0194 (2)
H11A	0.7864	-0.2374	0.0283	0.023*
H11B	0.8383	-0.1277	-0.0574	0.023*
C12	0.97328 (10)	-0.17627 (12)	0.07605 (10)	0.0225 (3)
H12A	1.0137	-0.2487	0.0335	0.027*
H12B	1.0249	-0.0954	0.0661	0.027*
C13	0.97401 (11)	-0.21099 (11)	0.20452 (11)	0.0223 (3)
H13A	1.0641	-0.2210	0.2361	0.027*
H13B	0.9293	-0.2961	0.2140	0.027*
C14	0.90724 (11)	-0.10416 (12)	0.27195 (10)	0.0211 (3)
H14A	0.9570	-0.0211	0.2689	0.025*
H14B	0.9049	-0.1309	0.3543	0.025*
C15	0.76953 (10)	-0.08069 (11)	0.22168 (9)	0.0176 (2)
H15A	0.7303	-0.0076	0.2643	0.021*
H15B	0.7174	-0.1609	0.2327	0.021*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0136 (5)	0.0149 (5)	0.0123 (4)	-0.0003 (4)	-0.0001 (4)	0.0019 (4)
C2	0.0159 (5)	0.0138 (5)	0.0134 (5)	-0.0014 (4)	-0.0002 (4)	0.0003 (4)
C3	0.0167 (5)	0.0126 (5)	0.0161 (5)	0.0017 (4)	-0.0004 (4)	0.0011 (4)
C4	0.0148 (5)	0.0121 (5)	0.0169 (5)	-0.0006 (4)	-0.0022 (4)	-0.0004 (4)
C5	0.0215 (5)	0.0203 (5)	0.0192 (5)	-0.0063 (4)	0.0045 (4)	-0.0042 (4)
C6	0.0224 (5)	0.0199 (5)	0.0198 (5)	-0.0033 (4)	0.0025 (4)	-0.0067 (4)
C7	0.0268 (6)	0.0135 (5)	0.0267 (6)	-0.0019 (4)	-0.0032 (5)	-0.0019 (4)
C8	0.0267 (6)	0.0217 (6)	0.0203 (5)	-0.0103 (5)	0.0022 (4)	0.0002 (4)
C9	0.0194 (5)	0.0192 (5)	0.0171 (5)	-0.0044 (4)	0.0017 (4)	-0.0018 (4)
C10	0.0127 (5)	0.0132 (5)	0.0164 (5)	0.0001 (4)	-0.0013 (4)	0.0004 (4)
C11	0.0153 (5)	0.0216 (5)	0.0208 (5)	0.0024 (4)	-0.0012 (4)	-0.0041 (4)
C12	0.0139 (5)	0.0243 (6)	0.0291 (6)	0.0034 (4)	0.0008 (4)	-0.0051 (5)
C13	0.0150 (5)	0.0189 (5)	0.0321 (6)	0.0022 (4)	-0.0052 (4)	0.0022 (5)
C14	0.0178 (5)	0.0249 (6)	0.0198 (5)	0.0013 (4)	-0.0045 (4)	0.0022 (4)
C15	0.0148 (5)	0.0209 (5)	0.0167 (5)	0.0023 (4)	-0.0011 (4)	0.0024 (4)

Geometric parameters (\AA , $^\circ$)

C1—C3 ⁱ	1.3941 (14)	C8—H8B	0.9900
C1—C2	1.4053 (14)	C9—H9A	0.9900
C1—C10	1.5245 (13)	C9—H9B	0.9900
C2—C3	1.3967 (14)	C10—C11	1.5353 (14)
C2—C4	1.5207 (14)	C10—C15	1.5384 (14)
C3—C1 ⁱ	1.3941 (14)	C10—H10	1.0000

C3—H3	0.9500	C11—C12	1.5316 (14)
C4—C9	1.5357 (14)	C11—H11A	0.9900
C4—C5	1.5368 (14)	C11—H11B	0.9900
C4—H4	1.0000	C12—C13	1.5237 (17)
C5—C6	1.5255 (15)	C12—H12A	0.9900
C5—H5A	0.9900	C12—H12B	0.9900
C5—H5B	0.9900	C13—C14	1.5243 (16)
C6—C7	1.5203 (16)	C13—H13A	0.9900
C6—H6A	0.9900	C13—H13B	0.9900
C6—H6B	0.9900	C14—C15	1.5308 (14)
C7—C8	1.5250 (17)	C14—H14A	0.9900
C7—H7A	0.9900	C14—H14B	0.9900
C7—H7B	0.9900	C15—H15A	0.9900
C8—C9	1.5322 (15)	C15—H15B	0.9900
C8—H8A	0.9900		
C3 ⁱ —C1—C2	117.82 (9)	C4—C9—H9A	109.3
C3 ⁱ —C1—C10	119.96 (9)	C8—C9—H9B	109.3
C2—C1—C10	122.21 (9)	C4—C9—H9B	109.3
C3—C2—C1	118.12 (9)	H9A—C9—H9B	107.9
C3—C2—C4	118.62 (9)	C1—C10—C11	113.85 (8)
C1—C2—C4	123.26 (9)	C1—C10—C15	110.75 (8)
C1 ⁱ —C3—C2	124.04 (10)	C11—C10—C15	110.17 (9)
C1 ⁱ —C3—H3	118.0	C1—C10—H10	107.3
C2—C3—H3	118.0	C11—C10—H10	107.3
C2—C4—C9	111.71 (8)	C15—C10—H10	107.3
C2—C4—C5	112.29 (8)	C12—C11—C10	111.42 (9)
C9—C4—C5	110.00 (9)	C12—C11—H11A	109.3
C2—C4—H4	107.5	C10—C11—H11A	109.3
C9—C4—H4	107.5	C12—C11—H11B	109.3
C5—C4—H4	107.5	C10—C11—H11B	109.3
C6—C5—C4	111.54 (9)	H11A—C11—H11B	108.0
C6—C5—H5A	109.3	C13—C12—C11	111.12 (9)
C4—C5—H5A	109.3	C13—C12—H12A	109.4
C6—C5—H5B	109.3	C11—C12—H12A	109.4
C4—C5—H5B	109.3	C13—C12—H12B	109.4
H5A—C5—H5B	108.0	C11—C12—H12B	109.4
C7—C6—C5	110.94 (9)	H12A—C12—H12B	108.0
C7—C6—H6A	109.5	C12—C13—C14	110.84 (9)
C5—C6—H6A	109.5	C12—C13—H13A	109.5
C7—C6—H6B	109.5	C14—C13—H13A	109.5
C5—C6—H6B	109.5	C12—C13—H13B	109.5
H6A—C6—H6B	108.0	C14—C13—H13B	109.5
C6—C7—C8	111.42 (9)	H13A—C13—H13B	108.1
C6—C7—H7A	109.3	C13—C14—C15	111.09 (9)
C8—C7—H7A	109.3	C13—C14—H14A	109.4
C6—C7—H7B	109.3	C15—C14—H14A	109.4
C8—C7—H7B	109.3	C13—C14—H14B	109.4

H7A—C7—H7B	108.0	C15—C14—H14B	109.4
C7—C8—C9	111.33 (9)	H14A—C14—H14B	108.0
C7—C8—H8A	109.4	C14—C15—C10	111.74 (9)
C9—C8—H8A	109.4	C14—C15—H15A	109.3
C7—C8—H8B	109.4	C10—C15—H15A	109.3
C9—C8—H8B	109.4	C14—C15—H15B	109.3
H8A—C8—H8B	108.0	C10—C15—H15B	109.3
C8—C9—C4	111.80 (9)	H15A—C15—H15B	107.9
C8—C9—H9A	109.3		
C3 ⁱ —C1—C2—C3	-1.33 (16)	C7—C8—C9—C4	-54.77 (12)
C10—C1—C2—C3	179.74 (9)	C2—C4—C9—C8	-179.76 (9)
C3 ⁱ —C1—C2—C4	178.54 (9)	C5—C4—C9—C8	54.84 (12)
C10—C1—C2—C4	-0.39 (15)	C3 ⁱ —C1—C10—C11	34.20 (13)
C1—C2—C3—C1 ⁱ	1.42 (17)	C2—C1—C10—C11	-146.89 (10)
C4—C2—C3—C1 ⁱ	-178.46 (9)	C3 ⁱ —C1—C10—C15	-90.60 (11)
C3—C2—C4—C9	-73.47 (12)	C2—C1—C10—C15	88.31 (12)
C1—C2—C4—C9	106.67 (11)	C1—C10—C11—C12	179.78 (9)
C3—C2—C4—C5	50.66 (12)	C15—C10—C11—C12	-55.11 (12)
C1—C2—C4—C5	-129.21 (10)	C10—C11—C12—C13	56.53 (13)
C2—C4—C5—C6	178.97 (9)	C11—C12—C13—C14	-56.64 (13)
C9—C4—C5—C6	-55.96 (12)	C12—C13—C14—C15	56.22 (12)
C4—C5—C6—C7	56.89 (12)	C13—C14—C15—C10	-55.83 (12)
C5—C6—C7—C8	-56.00 (12)	C1—C10—C15—C14	-178.25 (9)
C6—C7—C8—C9	54.96 (12)	C11—C10—C15—C14	54.90 (12)

Symmetry code: (i) $-x+1, -y, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
C6—H6A ⁱⁱ —Cg ⁱⁱ	0.99	2.62	3.520 (2)	150

Symmetry code: (ii) $-x+1, y+1/2, -z+1/2$.