

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Triclinic form of 1,2,4,5-tetracyclohexylbenzene


Joel T. Mague,* Lisa Linhardt, Iliana Medina,
Daniel J. Sattler and Mark J. FinkDepartment of Chemistry, Tulane University, New Orleans, LA 70118, USA
Correspondence e-mail: joelt@tulane.edu

Received 13 December 2007; accepted 18 December 2007

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å;
 R factor = 0.056; wR factor = 0.147; data-to-parameter ratio = 17.4.

The molecule of the title compound, $\text{C}_{30}\text{H}_{46}$, has a crystallographically imposed inversion center and the cyclohexyl groups are oriented with their methine H atoms pointing towards one another ($\text{H}\cdots\text{H} = 2.04$ Å).

Related literature

For related structures, see: Mague *et al.* (2008a,b).

Experimental

Crystal data

 $\text{C}_{30}\text{H}_{46}$
 $M_r = 406.67$ Triclinic, $P\bar{1}$
 $a = 6.014$ (1) Å $b = 10.001$ (1) Å
 $c = 10.513$ (2) Å
 $\alpha = 91.164$ (2)°
 $\beta = 94.815$ (2)°
 $\gamma = 106.336$ (3)°
 $V = 604.01$ (16) Å³ $Z = 1$
Mo $K\alpha$ radiation
 $\mu = 0.06$ mm⁻¹
 $T = 100$ (2) K
 $0.20 \times 0.13 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD area-
detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2002)
 $T_{\min} = 0.940$, $T_{\max} = 0.993$ 4730 measured reflections
2363 independent reflections
1862 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.147$
 $S = 1.03$
2363 reflections136 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.44$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus* program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Bruker, 2000); software used to prepare material for publication: *SHELXTL*.

We thank the Chemistry Department of Tulane University for support of the X-ray laboratory, and the Louisiana Board of Regents through the Louisiana Educational Quality Support Fund (Grant LEQSF (2003–2003)–ENH –TR-67) for the purchase of the APEX diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2128).

References

- Bruker (2000). *SMART* (Version 5.625) and *SHELXTL* (Version 6.10). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *SAINT-Plus*. Version 7.03. Bruker AXS Inc., Madison, Wisconsin, USA.
- Mague, J. T., Linhardt, L., Medina, I. & Fink, M. J. (2008a). *Acta Cryst.* **E64**, o375.
- Mague, J. T., Linhardt, L., Medina, I. & Fink, M. J. (2008b). *Acta Cryst.* **E64**, o335.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2002). *SADABS*. Version 2.05. University of Göttingen, Germany.

supplementary materials

Acta Cryst. (2008). E64, o376 [doi:10.1107/S1600536807067578]

Triclinic form of 1,2,4,5-tetracyclohexylbenzene

J. T. Mague, L. Linhardt, I. Medina, D. J. Sattler and M. J. Fink

Comment

Crystallization of 1,2,4,5-tetracyclohexylbenzene (C₃₀H₄₆) from hot methylcyclohexane forms colorless needle-shaped crystals together with a smaller quantity having a distinctly different block-shaped morphology. Many of the needles appear twinned but a fragment cut from a larger needle proved to be single and to be a triclinic modification. The molecule has crystallographically imposed centrosymmetry with the cyclohexyl rings adopting the chair conformation and oriented with their methine hydrogen atoms pointed towards one another (H4···H10 distance 2.04 Å) as is the case in the monoclinic modification (Mague *et al.*, 2008a). Again, there are very few close contacts between the *ortho*-disposed cyclohexyl rings, the shortest being H4···H15*b* (2.30 Å). Additional short contacts are H3···H9*b* (2.28 Å) and H3···H11*a'* (2.14 Å). The plane defined by the atoms C5, C6, C8, C9 ("seat" of the chair) is inclined to the plane of the aromatic ring by 87.3 (2)° while that for the other cyclohexyl ring (C11, C12, C14, C15) is inclined at an angle of only 49.4 (2)°, a much greater disparity in orientation than observed in the monoclinic modification. Another difference is the absence of any significant C—H···π interactions in the triclinic form.

Experimental

The title compound was prepared by the literature method (Mague *et al.*, 2008a)

Refinement

H-atoms were placed in calculated positions (C—H = 0.95 – 0.98 Å) and refined as riding on their carriers with isotropic displacement parameters 1.2 – 1.5 times those of the attached carbon atoms.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level and H-atoms are represented by spheres of arbitrary radius. Primed atoms are related to unprimed atoms by the symmetry operation $-x, 1 - y, -z$.

1,2,4,5-tetracyclohexylbenzene

Crystal data

C₃₀H₄₆

Z = 1

supplementary materials

$M_r = 406.67$	$F_{000} = 226$
Triclinic, $P\bar{1}$	$D_x = 1.118 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 6.014 (1) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 10.001 (1) \text{ \AA}$	Cell parameters from 2522 reflections
$c = 10.513 (2) \text{ \AA}$	$\theta = 2.8\text{--}28.3^\circ$
$\alpha = 91.164 (2)^\circ$	$\mu = 0.06 \text{ mm}^{-1}$
$\beta = 94.815 (2)^\circ$	$T = 100 (2) \text{ K}$
$\gamma = 106.336 (3)^\circ$	Plate, colorless
$V = 604.01 (16) \text{ \AA}^3$	$0.20 \times 0.13 \times 0.10 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer	2363 independent reflections
Radiation source: fine-focus sealed tube	1862 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
Detector resolution: 0 pixels mm^{-1}	$\theta_{\text{max}} = 26.0^\circ$
$T = 100(2) \text{ K}$	$\theta_{\text{min}} = 2.0^\circ$
ω scans	$h = -7 \rightarrow 7$
Absorption correction: multi-scan (SADABS; Sheldrick, 2002)	$k = -12 \rightarrow 12$
$T_{\text{min}} = 0.940$, $T_{\text{max}} = 0.993$	$l = -12 \rightarrow 12$
4730 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H-atom parameters constrained
$wR(F^2) = 0.147$	$w = 1/[\sigma^2(F_o^2) + (0.0804P)^2 + 0.1811P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
2363 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
136 parameters	$\Delta\rho_{\text{max}} = 0.44 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Experimental. The diffraction data were collected in three sets of 606 frames (ω scans, $0.3^\circ/\text{scan}$) at ϕ settings of 0, 120 and 240° .

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. H-atoms were placed in calculated positions (C—H = 0.95 – 0.98 Å) and included as riding contributions with isotropic displacement parameters 1.2 – 1.5 times those of the attached carbon atoms.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.1622 (3)	0.08720 (17)	0.92456 (15)	0.0140 (4)
C2	0.0316 (3)	−0.04621 (17)	0.87681 (15)	0.0141 (4)
C3	−0.1264 (3)	−0.12910 (17)	0.95305 (15)	0.0147 (4)
H3	−0.2148	−0.2192	0.9199	0.018*
C4	0.0526 (3)	−0.10585 (17)	0.74528 (15)	0.0147 (4)
H4	0.1660	−0.0318	0.7022	0.018*
C5	0.1482 (3)	−0.23296 (18)	0.75408 (16)	0.0179 (4)
H5A	0.3031	−0.2056	0.8033	0.021*
H5B	0.0431	−0.3065	0.8002	0.021*
C6	0.1695 (3)	−0.29087 (19)	0.62131 (16)	0.0207 (4)
H6A	0.2875	−0.2207	0.5789	0.025*
H6B	0.2235	−0.3754	0.6301	0.025*
C7	−0.0612 (3)	−0.32752 (18)	0.53844 (16)	0.0211 (4)
H7A	−0.1747	−0.4051	0.5756	0.025*
H7B	−0.0386	−0.3589	0.4518	0.025*
C8	−0.1570 (3)	−0.20179 (19)	0.52917 (16)	0.0201 (4)
H8A	−0.3118	−0.2296	0.4798	0.024*
H8B	−0.0522	−0.1283	0.4829	0.024*
C9	−0.1790 (3)	−0.14328 (18)	0.66175 (15)	0.0178 (4)
H9A	−0.2333	−0.0590	0.6525	0.021*
H9B	−0.2969	−0.2133	0.7043	0.021*
C10	0.3406 (3)	0.18370 (17)	0.84830 (15)	0.0146 (4)
H10	0.2704	0.1748	0.7578	0.018*
C11	0.4030 (3)	0.33832 (18)	0.88976 (16)	0.0178 (4)
H11A	0.4788	0.3524	0.9782	0.021*
H11B	0.2591	0.3680	0.8887	0.021*
C12	0.5673 (3)	0.42760 (18)	0.80049 (16)	0.0190 (4)
H12A	0.4885	0.4178	0.7128	0.023*
H12B	0.6075	0.5270	0.8296	0.023*
C13	0.7888 (3)	0.38188 (18)	0.79954 (17)	0.0197 (4)
H13A	0.8902	0.4373	0.7386	0.024*
H13B	0.8746	0.4000	0.8856	0.024*
C14	0.7330 (3)	0.22785 (18)	0.76158 (16)	0.0184 (4)
H14A	0.8788	0.1998	0.7672	0.022*
H14B	0.6637	0.2117	0.6718	0.022*
C15	0.5639 (3)	0.13813 (18)	0.84782 (16)	0.0166 (4)
H15A	0.6398	0.1461	0.9360	0.020*
H15B	0.5238	0.0392	0.8174	0.020*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0186 (9)	0.0177 (9)	0.0079 (8)	0.0082 (7)	0.0026 (6)	0.0007 (6)
C2	0.0180 (9)	0.0197 (9)	0.0065 (8)	0.0083 (7)	0.0020 (6)	0.0006 (6)
C3	0.0197 (9)	0.0159 (9)	0.0085 (8)	0.0046 (7)	0.0031 (6)	-0.0002 (6)
C4	0.0219 (9)	0.0135 (8)	0.0084 (8)	0.0031 (7)	0.0066 (7)	-0.0004 (6)
C5	0.0230 (9)	0.0215 (9)	0.0100 (8)	0.0071 (7)	0.0044 (7)	-0.0016 (7)
C6	0.0276 (10)	0.0217 (10)	0.0151 (9)	0.0092 (8)	0.0076 (7)	-0.0024 (7)
C7	0.0317 (11)	0.0208 (9)	0.0101 (8)	0.0051 (8)	0.0067 (7)	-0.0047 (7)
C8	0.0274 (10)	0.0239 (10)	0.0077 (8)	0.0047 (8)	0.0023 (7)	-0.0005 (7)
C9	0.0254 (10)	0.0220 (9)	0.0078 (8)	0.0091 (7)	0.0043 (7)	-0.0002 (7)
C10	0.0194 (9)	0.0187 (9)	0.0061 (8)	0.0051 (7)	0.0046 (6)	0.0002 (6)
C11	0.0220 (9)	0.0197 (9)	0.0138 (8)	0.0071 (7)	0.0093 (7)	0.0008 (7)
C12	0.0264 (10)	0.0176 (9)	0.0143 (9)	0.0066 (7)	0.0083 (7)	0.0008 (7)
C13	0.0215 (9)	0.0221 (10)	0.0155 (9)	0.0045 (7)	0.0081 (7)	0.0027 (7)
C14	0.0205 (9)	0.0250 (10)	0.0122 (8)	0.0087 (7)	0.0070 (7)	0.0025 (7)
C15	0.0215 (9)	0.0179 (9)	0.0112 (8)	0.0059 (7)	0.0045 (7)	0.0017 (7)

Geometric parameters (\AA , $^\circ$)

C1—C3 ⁱ	1.397 (2)	C8—H8B	0.9900
C1—C2	1.402 (2)	C9—H9A	0.9900
C1—C10	1.523 (2)	C9—H9B	0.9900
C2—C3	1.395 (2)	C10—C11	1.530 (2)
C2—C4	1.524 (2)	C10—C15	1.535 (2)
C3—C1 ⁱ	1.397 (2)	C10—H10	1.0000
C3—H3	0.9500	C11—C12	1.532 (2)
C4—C9	1.529 (2)	C11—H11A	0.9900
C4—C5	1.537 (2)	C11—H11B	0.9900
C4—H4	1.0000	C12—C13	1.526 (2)
C5—C6	1.529 (2)	C12—H12A	0.9900
C5—H5A	0.9900	C12—H12B	0.9900
C5—H5B	0.9900	C13—C14	1.519 (2)
C6—C7	1.522 (3)	C13—H13A	0.9900
C6—H6A	0.9900	C13—H13B	0.9900
C6—H6B	0.9900	C14—C15	1.529 (2)
C7—C8	1.525 (2)	C14—H14A	0.9900
C7—H7A	0.9900	C14—H14B	0.9900
C7—H7B	0.9900	C15—H15A	0.9900
C8—C9	1.531 (2)	C15—H15B	0.9900
C8—H8A	0.9900		
C3 ⁱ —C1—C2	117.64 (15)	C8—C9—H9A	109.3
C3 ⁱ —C1—C10	120.34 (15)	C4—C9—H9B	109.3
C2—C1—C10	122.00 (14)	C8—C9—H9B	109.3
C3—C2—C1	118.62 (14)	H9A—C9—H9B	107.9
C3—C2—C4	117.96 (14)	C1—C10—C11	114.86 (13)

C1—C2—C4	123.43 (14)	C1—C10—C15	111.62 (13)
C2—C3—C1 ⁱ	123.74 (16)	C11—C10—C15	109.51 (14)
C2—C3—H3	118.1	C1—C10—H10	106.8
C1 ⁱ —C3—H3	118.1	C11—C10—H10	106.8
C2—C4—C9	111.87 (13)	C15—C10—H10	106.8
C2—C4—C5	111.95 (13)	C10—C11—C12	110.82 (13)
C9—C4—C5	110.22 (14)	C10—C11—H11A	109.5
C2—C4—H4	107.5	C12—C11—H11A	109.5
C9—C4—H4	107.5	C10—C11—H11B	109.5
C5—C4—H4	107.5	C12—C11—H11B	109.5
C6—C5—C4	111.24 (14)	H11A—C11—H11B	108.1
C6—C5—H5A	109.4	C13—C12—C11	110.19 (14)
C4—C5—H5A	109.4	C13—C12—H12A	109.6
C6—C5—H5B	109.4	C11—C12—H12A	109.6
C4—C5—H5B	109.4	C13—C12—H12B	109.6
H5A—C5—H5B	108.0	C11—C12—H12B	109.6
C7—C6—C5	111.78 (14)	H12A—C12—H12B	108.1
C7—C6—H6A	109.3	C14—C13—C12	111.15 (15)
C5—C6—H6A	109.3	C14—C13—H13A	109.4
C7—C6—H6B	109.3	C12—C13—H13A	109.4
C5—C6—H6B	109.3	C14—C13—H13B	109.4
H6A—C6—H6B	107.9	C12—C13—H13B	109.4
C6—C7—C8	110.76 (14)	H13A—C13—H13B	108.0
C6—C7—H7A	109.5	C13—C14—C15	111.44 (14)
C8—C7—H7A	109.5	C13—C14—H14A	109.3
C6—C7—H7B	109.5	C15—C14—H14A	109.3
C8—C7—H7B	109.5	C13—C14—H14B	109.3
H7A—C7—H7B	108.1	C15—C14—H14B	109.3
C7—C8—C9	111.34 (14)	H14A—C14—H14B	108.0
C7—C8—H8A	109.4	C14—C15—C10	111.17 (14)
C9—C8—H8A	109.4	C14—C15—H15A	109.4
C7—C8—H8B	109.4	C10—C15—H15A	109.4
C9—C8—H8B	109.4	C14—C15—H15B	109.4
H8A—C8—H8B	108.0	C10—C15—H15B	109.4
C4—C9—C8	111.79 (14)	H15A—C15—H15B	108.0
C4—C9—H9A	109.3		
C3 ⁱ —C1—C2—C3	0.3 (3)	C2—C4—C9—C8	179.42 (14)
C10—C1—C2—C3	179.24 (15)	C5—C4—C9—C8	-55.33 (18)
C3 ⁱ —C1—C2—C4	-179.88 (15)	C7—C8—C9—C4	55.87 (19)
C10—C1—C2—C4	-0.9 (2)	C3 ⁱ —C1—C10—C11	-21.8 (2)
C1—C2—C3—C1 ⁱ	-0.3 (3)	C2—C1—C10—C11	159.29 (15)
C4—C2—C3—C1 ⁱ	179.86 (15)	C3 ⁱ —C1—C10—C15	103.66 (18)
C3—C2—C4—C9	59.29 (19)	C2—C1—C10—C15	-75.29 (19)
C1—C2—C4—C9	-120.57 (17)	C1—C10—C11—C12	-175.25 (14)
C3—C2—C4—C5	-65.00 (19)	C15—C10—C11—C12	58.25 (18)
C1—C2—C4—C5	115.15 (17)	C10—C11—C12—C13	-58.39 (19)
C2—C4—C5—C6	-179.65 (14)	C11—C12—C13—C14	56.47 (18)

supplementary materials

C9—C4—C5—C6	55.15 (18)	C12—C13—C14—C15	-55.38 (19)
C4—C5—C6—C7	-55.97 (19)	C13—C14—C15—C10	55.66 (19)
C5—C6—C7—C8	55.61 (19)	C1—C10—C15—C14	175.04 (13)
C6—C7—C8—C9	-55.23 (19)	C11—C10—C15—C14	-56.63 (17)

Symmetry codes: (i) $-x, -y, -z+2$.

Fig. 1

