

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Diaquabis[6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinato- $\kappa^3 N, N', O$](nitrate- $\kappa^2 O, O'$)lanthanum(III) monohydrate

Zhao Kai,^{a,b} Xian-Hong Yin,^{a*} Feng Yu,^a Zhu Jie^b and Cui-Wu Lin^b

^aCollege of Chemistry and Ecological Engineering, Guangxi University for Nationalities, Nanning 530006, People's Republic of China, and ^bCollege of Chemistry and Chemical Engineering, Guangxi University, Nanning 530004, People's Republic of China

Correspondence e-mail: yxhphd@163.com


Received 4 January 2008; accepted 18 January 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.025; wR factor = 0.077; data-to-parameter ratio = 12.9.

In the title complex, $[La(C_{11}H_{10}N_3O_2)_2(NO_3)(H_2O)_2] \cdot H_2O$, the La atom is coordinated by four N atoms and six O atoms derived from two 6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinate ligands, one nitrate anion and two water molecules. The molecules are linked together *via* hydrogen bonds involving the water molecules, forming a three-dimensional network.

Related literature

For related literature, see: Zhao *et al.* (2007); Yin *et al.* (2007).



Experimental

Crystal data

$[La(C_{11}H_{10}N_3O_2)_2(NO_3)(H_2O)_2] \cdot H_2O$
 $M_r = 687.41$
 Monoclinic, $P2_1/c$

$a = 17.396$ (2) Å
 $b = 15.0270$ (18) Å
 $c = 10.1607$ (13) Å
 $\beta = 94.737$ (2)°

$V = 2647.0$ (6) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 1.68$ mm⁻¹
 $T = 298$ (2) K
 $0.46 \times 0.45 \times 0.40$ mm

Data collection

Siemens SMART CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 1996)
 $T_{min} = 0.512$, $T_{max} = 0.553$
 (expected range = 0.473–0.510)

13524 measured reflections
 4656 independent reflections
 3941 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.076$
 $S = 1.01$
 4656 reflections

361 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 0.68$ e Å⁻³
 $\Delta\rho_{min} = -0.55$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> — <i>H</i> ··· <i>A</i>	<i>D</i> — <i>H</i>	<i>H</i> ··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> — <i>H</i> ··· <i>A</i>
O8—H8B···O10 ⁱ	0.85	2.13	2.918 (5)	154
O8—H8C···O4 ⁱⁱ	0.85	1.91	2.713 (4)	158
O9—H9B···O2 ⁱⁱⁱ	0.85	1.96	2.731 (4)	151
O9—H9B···N1	0.85	2.46	2.878 (4)	112
O10—H10C···N6 ^{iv}	0.85	2.49	3.156 (5)	136
O10—H10D···O9 ^v	0.85	2.24	2.977 (5)	146
O10—H10D···O1 ^{iv}	0.85	2.46	2.913 (4)	114

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x + 1, -y, -z + 1$; (iii) $x, -y - \frac{1}{2}, z + \frac{1}{2}$; (iv) $x, y + 1, z$; (v) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (grant No. 20761002). This research was sponsored by the Fund of the Talented Highland Research Programme of Guangxi University (grant No. 205121), the Science Foundation of the State Ethnic Affairs Commission (grant No. 07GX05), the Ministry of Education, Science and Technology Key Projects (grant No. 205121), the Development Foundation of Guangxi Research Institute of Chemical Industry, and the Science Foundation of Guangxi University for Nationalities (grant Nos. 0409032, 0409012, 0509ZD047), People's Republic of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2369).

References

- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Yin, X.-H., Zhao, K., Feng, Y. & Zhu, J. (2007). *Acta Cryst.* **E63**, m2926.
 Zhao, K., Yin, X.-H., Feng, Y. & Zhu, J. (2007). *Acta Cryst.* **E63**, m3024.

supplementary materials

Acta Cryst. (2008). E64, m410 [doi:10.1107/S1600536808001955]

Diaquabis[6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinato- $\kappa^3 N, N', O$](nitrate- $\kappa^2 O, O'$)lanthanum(III) monohydrate

Z. Kai, X.-H. Yin, F. Yu, Z. Jie and C.-W. Lin

Comment

Recently we reported the crystal structures of bis(6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinato)zinc(II) trihydrate (Yin *et al.*, 2007) and bis[3-chloro-6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinato]cobalt(II) 2.5-hydrate (Zhao *et al.*, 2007). As a continuation of these investigations, we report in this paper the crystal structure of Nitrate-diaqua-bis(6-(3,5-dimethyl-1*H*-pyrazol-1-yl)) picolinato)lanthanum(III) monohydrate.

The asymmetric unit of the title structure consists of the central mononuclear lanthanum(III) complex and one uncoordinated water molecule. The La atom is ten-coordinated by four N atoms and six O atoms derived from two 6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinate ligands (DPP), one bidentate nitrate anion and two water molecules that define a pseudotricapped trigonal environment for the lanthanum atom. The angles around La(III) atom range from 47.99 (8) to 144.42 (10)°, the La—O distances range from 2.452 (2) to 2.676 (3) Å, the La—N distances range is from 2.688 (3) to 2.811 (3) Å.

In the crystal structure, the oxygen atoms contribute to the formation of intermolecular hydrogen bonds involving the water molecules; three water molecules and three DPP O atoms form a rings *via* intermolecular H—O···H hydrogen bonds. A great number of hydrogen contacts link the complex into a three-dimensional network. (Fig. 2; for symmetry codes see Table 2).

Experimental

6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinic acid, and La(NO₃)₃·6H₂O were available commercially and were used without further purification. Equimolar 6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinic acid (1 mmol, 217 mg) was dissolved in anhydrous ethyl alcohol (AR,99.9%) (15 ml). The mixture was stirred to give a clear solution, To this solution was added La(NO₃)₃·6H₂O (0.33 mmol, 144 mg) in anhydrous alcohol (10 ml). After keeping the resulting solution in air to evaporate about half of the solvents, colorless blocks of the title complex were formed. The crystals were isolated, washed with alcohol three times (Yield 75%). Elemental analysis: found: C, 38.24; H, 3.91; N, 14.16%; calc. for C₂₂H₂₆LaN₇O₁₀: C, 38.44; H, 3.81; N, 14.26%.

Refinement

H atoms on C atoms were positioned geometrically and refined using a riding model with C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The water H atoms were located in difference Fourier maps and the O—H distances were constrained 0.85 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The structure of the title compound (I) showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. Crystal packing of (I) showing the hydrogen bonded interactions as dashed lines.

Diaquabis[6-(3,5-dimethyl-1*H*-pyrazol-1-yl)picolinato- κ^3N,N',O](nitrate- κ^2O,O')lanthanum(III) monohydrate

Crystal data

[La(C₁₁H₁₀N₃O₂)₂(NO₃)(H₂O)₂]·H₂O

M_r = 687.41

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 17.396 (2) Å

b = 15.0270 (18) Å

c = 10.1607 (13) Å

β = 94.737 (2)°

V = 2647.0 (6) Å³

Z = 4

*F*₀₀₀ = 1376

D_x = 1.725 Mg m⁻³

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 8663 reflections

θ = 2.4–28.3°

μ = 1.68 mm⁻¹

T = 298 (2) K

Block, colorless

0.46 × 0.45 × 0.40 mm

Data collection

Siemens SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 298(2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

4656 independent reflections

3941 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.022

θ_{max} = 25.0°

θ_{min} = 1.8°

h = -19→20

$T_{\min} = 0.512$, $T_{\max} = 0.553$
13524 measured reflections

$k = -17 \rightarrow 13$
 $l = -12 \rightarrow 12$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.025$	H-atom parameters constrained
$wR(F^2) = 0.077$	$w = 1/[\sigma^2(F_o^2) + (0.0425P)^2 + 3.0256P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
4656 reflections	$(\Delta/\sigma)_{\max} = 0.001$
361 parameters	$\Delta\rho_{\max} = 0.68 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.55 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
La1	0.725615 (10)	-0.033374 (12)	0.360394 (18)	0.02756 (8)
N1	0.86114 (15)	-0.10885 (18)	0.4512 (3)	0.0267 (6)
N2	0.88283 (16)	0.0047 (2)	0.6034 (3)	0.0325 (6)
N3	0.82454 (16)	0.05236 (19)	0.5339 (3)	0.0342 (7)
N4	0.59938 (17)	0.0651 (2)	0.2758 (3)	0.0353 (7)
N5	0.6301 (2)	0.0557 (2)	0.0593 (3)	0.0460 (8)
N6	0.6814 (2)	-0.0130 (3)	0.0897 (3)	0.0489 (9)
N7	0.83143 (19)	0.0967 (2)	0.2242 (4)	0.0497 (9)
O1	0.75996 (14)	-0.17799 (17)	0.2701 (3)	0.0428 (6)
O2	0.83910 (15)	-0.28912 (17)	0.2271 (3)	0.0442 (6)
O3	0.64201 (14)	0.02652 (17)	0.5260 (3)	0.0391 (6)
O4	0.53914 (16)	0.0940 (2)	0.5969 (3)	0.0562 (8)
O5	0.76750 (16)	0.12248 (19)	0.2609 (3)	0.0500 (7)
O6	0.84704 (16)	0.01423 (19)	0.2392 (3)	0.0479 (7)
O7	0.87668 (19)	0.1474 (3)	0.1778 (5)	0.0911 (14)
O8	0.60430 (15)	-0.13605 (19)	0.3366 (3)	0.0529 (8)

supplementary materials

H8B	0.5958	-0.1549	0.2578	0.064*
H8C	0.5645	-0.1090	0.3590	0.064*
O9	0.72302 (15)	-0.12026 (18)	0.5879 (3)	0.0466 (7)
H9B	0.7684	-0.1352	0.6167	0.056*
H9C	0.6957	-0.1670	0.5773	0.056*
O10	0.6289 (2)	0.7859 (2)	0.0804 (4)	0.0771 (10)
H10C	0.6333	0.8333	0.1260	0.093*
H10D	0.6685	0.7544	0.1017	0.093*
C1	0.8231 (2)	-0.2207 (2)	0.2862 (3)	0.0329 (8)
C2	0.88316 (19)	-0.1827 (2)	0.3894 (3)	0.0303 (7)
C3	0.9541 (2)	-0.2219 (3)	0.4173 (4)	0.0423 (9)
H3	0.9677	-0.2731	0.3733	0.051*
C4	1.0046 (2)	-0.1830 (2)	0.5129 (4)	0.0458 (10)
H4	1.0528	-0.2082	0.5340	0.055*
C5	0.9835 (2)	-0.1071 (3)	0.5771 (4)	0.0394 (9)
H5	1.0172	-0.0798	0.6406	0.047*
C6	0.91020 (18)	-0.0727 (2)	0.5436 (3)	0.0293 (7)
C7	0.9641 (3)	0.0119 (4)	0.8243 (5)	0.0706 (15)
H7A	1.0150	0.0209	0.7971	0.106*
H7B	0.9554	-0.0505	0.8367	0.106*
H7C	0.9590	0.0428	0.9058	0.106*
C8	0.9060 (2)	0.0469 (3)	0.7204 (4)	0.0434 (9)
C9	0.8629 (2)	0.1226 (3)	0.7219 (4)	0.0497 (10)
H9	0.8657	0.1654	0.7882	0.060*
C10	0.8133 (2)	0.1242 (3)	0.6055 (4)	0.0401 (9)
C11	0.7556 (3)	0.1936 (3)	0.5615 (5)	0.0584 (12)
H11A	0.7376	0.1834	0.4709	0.088*
H11B	0.7792	0.2513	0.5699	0.088*
H11C	0.7129	0.1909	0.6154	0.088*
C12	0.5787 (2)	0.0682 (3)	0.5086 (4)	0.0376 (8)
C13	0.5506 (2)	0.0867 (2)	0.3666 (4)	0.0387 (9)
C14	0.4786 (2)	0.1229 (3)	0.3322 (5)	0.0543 (11)
H14	0.4470	0.1401	0.3971	0.065*
C15	0.4546 (3)	0.1331 (3)	0.2014 (5)	0.0624 (13)
H15	0.4058	0.1558	0.1766	0.075*
C16	0.5029 (2)	0.1095 (3)	0.1070 (5)	0.0558 (11)
H16	0.4871	0.1143	0.0175	0.067*
C17	0.5761 (2)	0.0784 (3)	0.1492 (4)	0.0417 (9)
C18	0.6004 (3)	0.1738 (3)	-0.1168 (5)	0.0749 (16)
H18A	0.6331	0.2046	-0.1733	0.112*
H18B	0.5884	0.2124	-0.0460	0.112*
H18C	0.5536	0.1564	-0.1668	0.112*
C19	0.6413 (3)	0.0926 (3)	-0.0609 (4)	0.0540 (11)
C20	0.6981 (3)	0.0444 (3)	-0.1085 (5)	0.0657 (14)
H20	0.7179	0.0526	-0.1898	0.079*
C21	0.7219 (3)	-0.0198 (3)	-0.0151 (4)	0.0567 (12)
C22	0.7836 (3)	-0.0884 (4)	-0.0240 (5)	0.0836 (17)
H22A	0.8261	-0.0752	0.0394	0.125*
H22B	0.8010	-0.0879	-0.1112	0.125*

supplementary materials

La1—O3	2.482 (2)	C3—H3	0.9300
La1—O8	2.609 (3)	C4—C5	1.379 (5)
La1—O6	2.630 (3)	C4—H4	0.9300
La1—O9	2.659 (3)	C5—C6	1.392 (5)
La1—O5	2.676 (3)	C5—H5	0.9300
La1—N3	2.688 (3)	C7—C8	1.495 (6)
La1—N1	2.709 (3)	C7—H7A	0.9600
La1—N4	2.727 (3)	C7—H7B	0.9600
La1—N6	2.811 (3)	C7—H7C	0.9600
N1—C6	1.331 (4)	C8—C9	1.363 (6)
N1—C2	1.347 (4)	C9—C10	1.405 (5)
N2—C8	1.378 (5)	C9—H9	0.9300
N2—N3	1.386 (4)	C10—C11	1.491 (6)
N2—C6	1.413 (5)	C11—H11A	0.9600
N3—C10	1.325 (5)	C11—H11B	0.9600
N4—C17	1.331 (5)	C11—H11C	0.9600
N4—C13	1.344 (5)	C12—C13	1.510 (5)
N5—C19	1.369 (5)	C13—C14	1.383 (5)
N5—N6	1.382 (5)	C14—C15	1.369 (7)
N5—C17	1.406 (5)	C14—H14	0.9300
N6—C21	1.329 (6)	C15—C16	1.373 (7)
N7—O7	1.218 (4)	C15—H15	0.9300
N7—O5	1.263 (4)	C16—C17	1.392 (5)
N7—O6	1.275 (4)	C16—H16	0.9300
O1—C1	1.271 (4)	C18—C19	1.501 (7)
O2—C1	1.234 (4)	C18—H18A	0.9600
O3—C12	1.266 (4)	C18—H18B	0.9600
O4—C12	1.238 (4)	C18—H18C	0.9600
O8—H8B	0.8499	C19—C20	1.346 (7)
O8—H8C	0.8500	C20—C21	1.393 (7)
O9—H9B	0.8500	C20—H20	0.9300
O9—H9C	0.8499	C21—C22	1.496 (7)
O10—H10C	0.8501	C22—H22A	0.9600
O10—H10D	0.8499	C22—H22B	0.9600
C1—C2	1.528 (5)	C22—H22C	0.9600
C2—C3	1.375 (5)		
O1—La1—O3	138.56 (9)	O2—C1—C2	118.4 (3)
O1—La1—O8	70.21 (8)	O1—C1—C2	115.8 (3)
O3—La1—O8	76.27 (9)	N1—C2—C3	122.8 (3)
O1—La1—O6	80.56 (9)	N1—C2—C1	115.0 (3)
O3—La1—O6	139.19 (9)	C3—C2—C1	122.2 (3)
O8—La1—O6	142.37 (9)	C2—C3—C4	118.1 (4)
O1—La1—O9	84.97 (9)	C2—C3—H3	121.0
O3—La1—O9	62.33 (8)	C4—C3—H3	121.0
O8—La1—O9	73.66 (9)	C5—C4—C3	120.3 (3)
O6—La1—O9	127.78 (8)	C5—C4—H4	119.9
O1—La1—O5	123.50 (10)	C3—C4—H4	119.9
O3—La1—O5	97.58 (9)	C4—C5—C6	117.7 (3)
O8—La1—O5	136.54 (9)	C4—C5—H5	121.1

O6—La1—O5	47.99 (8)	C6—C5—H5	121.1
O9—La1—O5	141.55 (9)	N1—C6—C5	122.8 (3)
O1—La1—N3	120.50 (8)	N1—C6—N2	114.8 (3)
O3—La1—N3	76.17 (9)	C5—C6—N2	122.3 (3)
O8—La1—N3	144.42 (10)	C8—C7—H7A	109.5
O6—La1—N3	71.41 (10)	C8—C7—H7B	109.5
O9—La1—N3	73.81 (9)	H7A—C7—H7B	109.5
O5—La1—N3	69.44 (9)	C8—C7—H7C	109.5
O1—La1—N1	61.59 (8)	H7A—C7—H7C	109.5
O3—La1—N1	117.58 (8)	H7B—C7—H7C	109.5
O8—La1—N1	117.29 (9)	C9—C8—N2	105.7 (3)
O6—La1—N1	64.12 (8)	C9—C8—C7	128.8 (4)
O9—La1—N1	64.84 (8)	N2—C8—C7	125.4 (4)
O5—La1—N1	103.81 (8)	C8—C9—C10	107.8 (3)
N3—La1—N1	59.02 (8)	C8—C9—H9	126.1
O1—La1—N4	125.28 (8)	C10—C9—H9	126.1
O3—La1—N4	61.02 (9)	N3—C10—C9	109.9 (3)
O8—La1—N4	70.58 (9)	N3—C10—C11	122.1 (3)
O6—La1—N4	111.20 (9)	C9—C10—C11	128.0 (4)
O9—La1—N4	118.00 (9)	C10—C11—H11A	109.5
O5—La1—N4	69.10 (9)	C10—C11—H11B	109.5
N3—La1—N4	113.69 (9)	H11A—C11—H11B	109.5
N1—La1—N4	171.90 (9)	C10—C11—H11C	109.5
O1—La1—N6	77.64 (10)	H11A—C11—H11C	109.5
O3—La1—N6	119.69 (9)	H11B—C11—H11C	109.5
O8—La1—N6	79.60 (11)	O4—C12—O3	125.6 (4)
O6—La1—N6	71.22 (10)	O4—C12—C13	118.7 (3)
O9—La1—N6	151.80 (10)	O3—C12—C13	115.7 (3)
O5—La1—N6	66.17 (10)	N4—C13—C14	122.1 (4)
N3—La1—N6	134.28 (10)	N4—C13—C12	116.0 (3)
N1—La1—N6	122.64 (9)	C14—C13—C12	121.9 (4)
N4—La1—N6	58.94 (9)	C15—C14—C13	119.1 (4)
C6—N1—C2	118.3 (3)	C15—C14—H14	120.4
C6—N1—La1	124.0 (2)	C13—C14—H14	120.4
C2—N1—La1	117.3 (2)	C14—C15—C16	119.6 (4)
C8—N2—N3	110.6 (3)	C14—C15—H15	120.2
C8—N2—C6	131.7 (3)	C16—C15—H15	120.2
N3—N2—C6	117.7 (3)	C15—C16—C17	118.0 (4)
C10—N3—N2	106.0 (3)	C15—C16—H16	121.0
C10—N3—La1	129.4 (2)	C17—C16—H16	121.0
N2—N3—La1	119.2 (2)	N4—C17—C16	123.1 (4)
C17—N4—C13	117.9 (3)	N4—C17—N5	115.2 (3)
C17—N4—La1	124.0 (2)	C16—C17—N5	121.7 (4)
C13—N4—La1	116.9 (2)	C19—C18—H18A	109.5
C19—N5—N6	111.5 (4)	C19—C18—H18B	109.5
C19—N5—C17	129.2 (4)	H18A—C18—H18B	109.5
N6—N5—C17	119.3 (3)	C19—C18—H18C	109.5
C21—N6—N5	104.7 (3)	H18A—C18—H18C	109.5
C21—N6—La1	130.9 (3)	H18B—C18—H18C	109.5

supplementary materials

N5—N6—La1	114.8 (2)	C20—C19—N5	105.4 (4)
O7—N7—O5	122.4 (4)	C20—C19—C18	129.8 (4)
O7—N7—O6	121.1 (4)	N5—C19—C18	124.7 (4)
O5—N7—O6	116.5 (3)	C19—C20—C21	108.3 (4)
C1—O1—La1	129.3 (2)	C19—C20—H20	125.9
C12—O3—La1	129.5 (2)	C21—C20—H20	125.9
N7—O5—La1	96.7 (2)	N6—C21—C20	110.2 (4)
N7—O6—La1	98.6 (2)	N6—C21—C22	122.2 (4)
La1—O8—H8B	111.1	C20—C21—C22	127.7 (5)
La1—O8—H8C	111.3	C21—C22—H22A	109.5
H8B—O8—H8C	109.2	C21—C22—H22B	109.5
La1—O9—H9B	110.3	H22A—C22—H22B	109.5
La1—O9—H9C	110.2	C21—C22—H22C	109.5
H9B—O9—H9C	108.6	H22A—C22—H22C	109.5
H10C—O10—H10D	107.0	H22B—C22—H22C	109.5
O2—C1—O1	125.8 (3)		
O1—La1—N1—C6	-179.1 (3)	N4—La1—O3—C12	5.2 (3)
O3—La1—N1—C6	-46.4 (3)	N6—La1—O3—C12	-0.6 (3)
O8—La1—N1—C6	-134.5 (2)	O7—N7—O5—La1	-175.3 (4)
O6—La1—N1—C6	87.7 (3)	O6—N7—O5—La1	4.1 (3)
O9—La1—N1—C6	-80.9 (2)	O1—La1—O5—N7	-33.0 (3)
O5—La1—N1—C6	60.0 (3)	O3—La1—O5—N7	152.8 (2)
N3—La1—N1—C6	4.7 (2)	O8—La1—O5—N7	-129.2 (2)
N4—La1—N1—C6	31.5 (7)	O6—La1—O5—N7	-2.4 (2)
N6—La1—N1—C6	130.3 (2)	O9—La1—O5—N7	98.7 (2)
O1—La1—N1—C2	7.7 (2)	N3—La1—O5—N7	80.7 (2)
O3—La1—N1—C2	140.5 (2)	N1—La1—O5—N7	31.9 (2)
O8—La1—N1—C2	52.3 (2)	N4—La1—O5—N7	-152.2 (2)
O6—La1—N1—C2	-85.4 (2)	N6—La1—O5—N7	-88.0 (2)
O9—La1—N1—C2	106.0 (2)	O7—N7—O6—La1	175.2 (4)
O5—La1—N1—C2	-113.1 (2)	O5—N7—O6—La1	-4.2 (4)
N3—La1—N1—C2	-168.4 (3)	O1—La1—O6—N7	156.9 (2)
N4—La1—N1—C2	-141.6 (6)	O3—La1—O6—N7	-37.2 (3)
N6—La1—N1—C2	-42.8 (3)	O8—La1—O6—N7	117.9 (2)
C8—N2—N3—C10	1.6 (4)	O9—La1—O6—N7	-127.1 (2)
C6—N2—N3—C10	-175.5 (3)	O5—La1—O6—N7	2.4 (2)
C8—N2—N3—La1	-155.0 (3)	N3—La1—O6—N7	-76.3 (2)
C6—N2—N3—La1	27.9 (4)	N1—La1—O6—N7	-140.2 (2)
O1—La1—N3—C10	-170.6 (3)	N4—La1—O6—N7	32.6 (2)
O3—La1—N3—C10	-31.9 (3)	N6—La1—O6—N7	76.8 (2)
O8—La1—N3—C10	-72.1 (4)	La1—O1—C1—O2	-172.2 (3)
O6—La1—N3—C10	122.9 (3)	La1—O1—C1—C2	8.3 (5)
O9—La1—N3—C10	-96.7 (3)	C6—N1—C2—C3	-0.3 (5)
O5—La1—N3—C10	71.8 (3)	La1—N1—C2—C3	173.2 (3)
N1—La1—N3—C10	-166.7 (4)	C6—N1—C2—C1	179.2 (3)
N4—La1—N3—C10	17.3 (3)	La1—N1—C2—C1	-7.3 (4)
N6—La1—N3—C10	86.3 (3)	O2—C1—C2—N1	-179.0 (3)
O1—La1—N3—N2	-20.2 (3)	O1—C1—C2—N1	0.6 (4)
O3—La1—N3—N2	118.4 (3)	O2—C1—C2—C3	0.4 (5)

O8—La1—N3—N2	78.3 (3)	O1—C1—C2—C3	-180.0 (3)
O6—La1—N3—N2	-86.7 (2)	N1—C2—C3—C4	-0.2 (6)
O9—La1—N3—N2	53.7 (2)	C1—C2—C3—C4	-179.6 (3)
O5—La1—N3—N2	-137.8 (3)	C2—C3—C4—C5	-0.2 (6)
N1—La1—N3—N2	-16.3 (2)	C3—C4—C5—C6	1.0 (6)
N4—La1—N3—N2	167.7 (2)	C2—N1—C6—C5	1.2 (5)
N6—La1—N3—N2	-123.4 (2)	La1—N1—C6—C5	-171.9 (3)
O1—La1—N4—C17	-44.0 (3)	C2—N1—C6—N2	179.8 (3)
O3—La1—N4—C17	-175.1 (3)	La1—N1—C6—N2	6.8 (4)
O8—La1—N4—C17	-90.4 (3)	C4—C5—C6—N1	-1.5 (5)
O6—La1—N4—C17	49.4 (3)	C4—C5—C6—N2	179.9 (3)
O9—La1—N4—C17	-148.8 (3)	C8—N2—C6—N1	161.2 (4)
O5—La1—N4—C17	73.0 (3)	N3—N2—C6—N1	-22.4 (4)
N3—La1—N4—C17	127.6 (3)	C8—N2—C6—C5	-20.2 (6)
N1—La1—N4—C17	102.6 (6)	N3—N2—C6—C5	156.2 (3)
N6—La1—N4—C17	-1.1 (3)	N3—N2—C8—C9	-1.4 (4)
O1—La1—N4—C13	123.1 (3)	C6—N2—C8—C9	175.2 (4)
O3—La1—N4—C13	-8.1 (2)	N3—N2—C8—C7	175.9 (4)
O8—La1—N4—C13	76.6 (3)	C6—N2—C8—C7	-7.5 (7)
O6—La1—N4—C13	-143.6 (3)	N2—C8—C9—C10	0.7 (5)
O9—La1—N4—C13	18.3 (3)	C7—C8—C9—C10	-176.5 (5)
O5—La1—N4—C13	-120.0 (3)	N2—N3—C10—C9	-1.1 (4)
N3—La1—N4—C13	-65.3 (3)	La1—N3—C10—C9	152.2 (3)
N1—La1—N4—C13	-90.3 (6)	N2—N3—C10—C11	178.7 (4)
N6—La1—N4—C13	166.0 (3)	La1—N3—C10—C11	-28.0 (6)
C19—N5—N6—C21	-2.3 (5)	C8—C9—C10—N3	0.3 (5)
C17—N5—N6—C21	177.8 (4)	C8—C9—C10—C11	-179.5 (4)
C19—N5—N6—La1	147.9 (3)	La1—O3—C12—O4	177.9 (3)
C17—N5—N6—La1	-32.0 (4)	La1—O3—C12—C13	-1.9 (5)
O1—La1—N6—C21	-58.1 (4)	C17—N4—C13—C14	-0.9 (6)
O3—La1—N6—C21	162.6 (4)	La1—N4—C13—C14	-168.8 (3)
O8—La1—N6—C21	-129.9 (4)	C17—N4—C13—C12	178.5 (3)
O6—La1—N6—C21	26.0 (4)	La1—N4—C13—C12	10.6 (4)
O9—La1—N6—C21	-111.3 (4)	O4—C12—C13—N4	173.7 (4)
O5—La1—N6—C21	77.5 (4)	O3—C12—C13—N4	-6.5 (5)
N3—La1—N6—C21	62.7 (4)	O4—C12—C13—C14	-7.0 (6)
N1—La1—N6—C21	-14.1 (5)	O3—C12—C13—C14	172.9 (4)
N4—La1—N6—C21	156.6 (4)	N4—C13—C14—C15	3.3 (7)
O1—La1—N6—N5	161.5 (3)	C12—C13—C14—C15	-176.0 (4)
O3—La1—N6—N5	22.2 (3)	C13—C14—C15—C16	-1.8 (7)
O8—La1—N6—N5	89.7 (3)	C14—C15—C16—C17	-1.9 (7)
O6—La1—N6—N5	-114.4 (3)	C13—N4—C17—C16	-3.1 (6)
O9—La1—N6—N5	108.3 (3)	La1—N4—C17—C16	163.9 (3)
O5—La1—N6—N5	-62.9 (3)	C13—N4—C17—N5	179.1 (3)
N3—La1—N6—N5	-77.7 (3)	La1—N4—C17—N5	-13.9 (5)
N1—La1—N6—N5	-154.5 (2)	C15—C16—C17—N4	4.5 (7)
N4—La1—N6—N5	16.2 (3)	C15—C16—C17—N5	-177.9 (4)
O3—La1—O1—C1	-109.3 (3)	C19—N5—C17—N4	-148.7 (4)
O8—La1—O1—C1	-147.2 (3)	N6—N5—C17—N4	31.1 (5)

supplementary materials

O6—La1—O1—C1	56.9 (3)	C19—N5—C17—C16	33.4 (7)
O9—La1—O1—C1	-72.8 (3)	N6—N5—C17—C16	-146.7 (4)
O5—La1—O1—C1	79.4 (3)	N6—N5—C19—C20	2.3 (5)
N3—La1—O1—C1	-4.9 (3)	C17—N5—C19—C20	-177.8 (4)
N1—La1—O1—C1	-8.7 (3)	N6—N5—C19—C18	-173.9 (4)
N4—La1—O1—C1	166.2 (3)	C17—N5—C19—C18	6.0 (7)
N6—La1—O1—C1	129.6 (3)	N5—C19—C20—C21	-1.4 (5)
O1—La1—O3—C12	-106.5 (3)	C18—C19—C20—C21	174.5 (5)
O8—La1—O3—C12	-70.0 (3)	N5—N6—C21—C20	1.3 (5)
O6—La1—O3—C12	94.7 (3)	La1—N6—C21—C20	-141.9 (4)
O9—La1—O3—C12	-148.5 (3)	N5—N6—C21—C22	-178.8 (5)
O5—La1—O3—C12	66.2 (3)	La1—N6—C21—C22	37.9 (7)
N3—La1—O3—C12	132.8 (3)	C19—C20—C21—N6	0.0 (6)
N1—La1—O3—C12	176.2 (3)	C19—C20—C21—C22	-179.8 (5)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O8—H8B \cdots O10 ⁱ	0.85	2.13	2.918 (5)	154
O8—H8C \cdots O4 ⁱⁱ	0.85	1.91	2.713 (4)	158
O9—H9B \cdots O2 ⁱⁱⁱ	0.85	1.96	2.731 (4)	151
O9—H9B \cdots N1	0.85	2.46	2.878 (4)	112
O10—H10C \cdots N6 ^{iv}	0.85	2.49	3.156 (5)	136
O10—H10D \cdots O9 ^v	0.85	2.24	2.977 (5)	146
O10—H10D \cdots O1 ^{iv}	0.85	2.46	2.913 (4)	114

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, -y, -z+1$; (iii) $x, -y-1/2, z+1/2$; (iv) $x, y+1, z$; (v) $x, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

