

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

cis-Bis(azido- κ N)bis(pyridine-2-carboxamide- κ^2 N¹,O)nickel(II)

Marijana Đaković and Zora Popović*

 Department of Chemistry, Laboratory of General and Inorganic Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, HR-10000 Zagreb, Croatia
 Correspondence e-mail: zpopovic@chem.pmf.hr


Received 14 December 2007; accepted 4 January 2008

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.030; wR factor = 0.068; data-to-parameter ratio = 18.7.

The title compound, $[\text{Ni}(\text{N}_3)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2]$, was obtained as the first crystalline product from the reaction of $\text{Ni}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$, picolinamide and NaN_3 in aqueous media. After a few days in the mother liquor, crystals of the *cis* isomer transformed into the *trans* isomer [Đaković & Popović (2007). *Acta Cryst.* **C63**, m507–m509]. The Ni atom exhibits a distorted octahedral environment and contains two azide ions and two planar *N,O*-chelating picolinamide ligands, all *cis* related. The dihedral angle between the two chelate rings is $82.43(7)^\circ$. Pairs of molecules are linked by $\text{N}-\text{H} \cdots \text{N}$ hydrogen bonds into cyclic $R_2^2(16)$ dimers, which are further packed into a three-dimensional framework by $C(6)$ and $C(8)$ chains by $\text{N}-\text{H} \cdots \text{N}$ hydrogen bonds.

Related literature

For information on the importance of azides in complexation, see Yuwen *et al.* (2000). A *trans* isomer of the title compound $[\text{Ni}(\text{N}_3)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2]$ has been reported by Đaković & Popović (2007). For related literature, see: Allen *et al.* (1987); Bernstein *et al.* (1995); Etter (1990).



Experimental

Crystal data

 $[\text{Ni}(\text{N}_3)_2(\text{C}_6\text{H}_6\text{N}_2\text{O})_2]$
 $M_r = 387.01$
 Monoclinic, $P2_1/c$
 $a = 14.3438(5)$ Å
 $b = 6.6986(2)$ Å
 $c = 18.7969(10)$ Å
 $\beta = 120.738(3)^\circ$
 $V = 1552.34(12)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.28$ mm⁻¹
 $T = 296$ K
 $0.22 \times 0.18 \times 0.05$ mm

Data collection

 Oxford Diffraction Xcalibur
 diffractometer with Sapphire3
 detector
 Absorption correction: multi-scan
 (*CrysAlisPro*; Oxford)

 Diffraction, 2007
 $T_{\min} = 0.815$, $T_{\max} = 0.938$
 15901 measured reflections
 4516 independent reflections
 2692 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.034$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.068$
 $S = 0.89$
 4516 reflections
 242 parameters

 H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\max} = 0.43$ e Å⁻³
 $\Delta\rho_{\min} = -0.30$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Ni1—O1	2.0701 (13)	Ni1—N3	2.0559 (17)
Ni1—O2	2.0941 (12)	Ni1—N5	2.0652 (14)
Ni1—N1	2.0685 (17)	Ni1—N8	2.0863 (17)
O1—Ni1—O2	93.30 (5)	O2—Ni1—N8	89.94 (5)
O1—Ni1—N1	78.45 (6)	N1—Ni1—N3	165.67 (6)
O1—Ni1—N3	89.76 (6)	N1—Ni1—N5	92.18 (6)
O1—Ni1—N5	88.69 (6)	N1—Ni1—N8	98.83 (7)
O1—Ni1—N8	175.91 (6)	N3—Ni1—N5	95.69 (7)
O2—Ni1—N1	94.34 (6)	N3—Ni1—N8	93.36 (7)
O2—Ni1—N3	78.08 (6)	N5—Ni1—N8	88.36 (6)
O2—Ni1—N5	173.44 (7)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N2—H12 \cdots N7 ⁱ	0.86 (3)	2.12 (2)	2.967 (3)	165 (3)
N2—H13 \cdots N10 ⁱⁱ	0.85 (3)	2.31 (3)	3.154 (4)	172 (2)
N4—H14 \cdots N8 ⁱⁱⁱ	0.80 (2)	2.36 (2)	3.136 (3)	164 (2)
N4—H15 \cdots N10 ^{iv}	0.94 (3)	2.50 (3)	3.442 (4)	179 (3)

 Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $x, y - 1, z$; (iv) $-x, -y, -z$.

Data collection: *CrysAlisPro* (Oxford Diffraction, 2007); cell refinement: *CrysAlisPro*; data reduction: *CrysAlisPro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *PLATON* (Spek, 2003).

This research was supported by the Ministry of Science, Education and Sports of the Republic of Croatia, Zagreb (grant No. 119-1193079-1332).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2158).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Đaković, M. & Popović, Z. (2007). *Acta Cryst.* **C63**, m507–m509.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Oxford Diffraction (2007). *CrysAlisPro*. Version 171.32. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Yuwen, L., Yi, L., Songshen, Q. & Fengjiao, D. (2000). *Thermochim. Acta*, **351**, 51–54.

supplementary materials

Acta Cryst. (2008). E64, m311-m312 [doi:10.1107/S1600536808000299]

***cis*-Bis(azido- κN)bis(pyridine-2-carboxamide- $\kappa^2 N^1, O$)nickel(II)**

M. Đakovic and Z. Popovic

Comment

This research is a part of our wider interest of the structural role of azide ions and of its metal complexes in metabolic processes of mitochondria (Yuwen *et al.*, 2000).

In the title compound Ni^{II} atom lies in a general position and exhibits distorted octahedral environment (Fig. 1). The coordination sphere is composed by two *cis*-related *N,O*-chelating picolinamide and two azide ligands. The picolinamide ligands are bonded more tightly (Table 1) than in its *trans*-isomer (Đaković & Popović, 2007). All other bond lengths are comparable to the values reported for similar compounds (Allen *et al.*, 1987). The azide ligands are coordinated to the central metal ion in non-linear mode (123.7 (1) and 123.0 (1)°) with the azide bond angles being 178.2 (2) and 176.9 (3)°.

The crystals of the title compound (I) are turquoise-green apart from the crystals of its *trans*-isomer which are olive-green.

The crystal structure (Fig 2) is stabilized by N—H \cdots N hydrogen bond network between carboxamide groups and azide ligands. Typical amide N—H \cdots O carboxamide dimers of $R_2^2(8)$ found in *trans*-isomer are not observed in the *cis*-isomer. Instead, the amide N atoms, N2 and N4, are involved in two hydrogen bonds, forming $R_2^2(16)$ rings, between two neighbouring molecules whereas C(8) chains along the axis *c* and C(6) chains along the axis *b* complete the network (Bernstein *et al.*, 1995; Etter, 1990).

The slightly smaller density of (I), and the fact that it is formed first and then transformed into its *trans*-isomer, suggests that (I) is the thermodynamically less stable isomer.

Experimental

The title compound was obtained by *in situ* reaction from Ni^{II} nitrate hexahydrate, sodium azide and picolinamide in a 1: 2: 2 molar ratio. All starting substances were dissolved in water. The sodium azide solution was added in small portions with stirring into the solution mixture of the picolinamide and Ni^{II} nitrate. In a few h the dark-green crystals of (I) were isolated. If the crystals are left in a mother liquor for a few days the dark-green crystals of (I) were transformed into the olive-green *trans*-isomer.

Refinement

Aromatic H atoms were fixed in geometrically idealized positions and refined using a riding model with [C—H = 0.93 Å and $U_{iso}(H) = 1.2U_{eq}(C)$]. The amide H atoms were placed in the positions indicated by difference electron-density maps and their positions were allowed to refine together with individual isotropic displacement parameters.

Figures




Fig. 1. The ORTEP-3 drawing of (I) with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 30% probability level.




Fig. 2. Crystal packing of (I) showing the hydrogen bonds as dashed lines.

cis-Bis(azido- κ N)bis(pyridine-2-carboxamide- κ^2 N¹,O)nickel(II)

Crystal data

[Ni(N₃)₂(C₆H₆N₂O)₂]

$M_r = 387.01$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.3438$ (5) Å

$b = 6.6986$ (2) Å

$c = 18.7969$ (10) Å

$\beta = 120.738$ (3)°

$V = 1552.34$ (12) Å³

$Z = 4$

$F_{000} = 792$

$D_x = 1.656$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 5015 reflections

$\theta = 3.8$ – 32.4 °

$\mu = 1.28$ mm⁻¹

$T = 296$ K

Plate, blue

$0.22 \times 0.18 \times 0.05$ mm

Data collection

Oxford Diffraction Xcalibur
diffractometer with Sapphire3 detector

Radiation source: Enhance (Mo) X-ray Source

Monochromator: graphite

Detector resolution: 16.3426 pixels mm⁻¹

$T = 296$ K

CCD scans

Absorption correction: multi-scan
(CrysAlisPro; Oxford Diffraction, 2007)

$T_{\min} = 0.815$, $T_{\max} = 0.938$

15901 measured reflections

4516 independent reflections

2692 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.034$

$\theta_{\text{max}} = 30.0$ °

$\theta_{\text{min}} = 3.8$ °

$h = -20 \rightarrow 20$

$k = -9 \rightarrow 9$

$l = -26 \rightarrow 26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.030$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.068$	$w = 1/[\sigma^2(F_o^2) + (0.0323P)^2]$
$S = 0.89$	where $P = (F_o^2 + 2F_c^2)/3$
4516 reflections	$(\Delta/\sigma)_{\max} < 0.001$
242 parameters	$\Delta\rho_{\max} = 0.43 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement on F^2 for ALL reflections except those flagged by the user for potential systematic errors. Weighted R -factors wR and all goodnesses of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > \sigma(F^2)$ is used only for calculating $-R$ -factor-obs *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.23220 (2)	0.37068 (3)	0.22317 (1)	0.0317 (1)
O1	0.26131 (9)	0.2742 (2)	0.33739 (8)	0.0434 (4)
O2	0.20315 (9)	0.08148 (18)	0.17407 (8)	0.0398 (4)
N1	0.39913 (11)	0.3396 (2)	0.28785 (9)	0.0334 (4)
N2	0.38916 (17)	0.2368 (3)	0.47094 (11)	0.0477 (6)
N3	0.06825 (11)	0.3406 (2)	0.17609 (8)	0.0323 (4)
N4	0.07207 (16)	-0.1400 (3)	0.09546 (11)	0.0461 (6)
N5	0.24200 (12)	0.6613 (2)	0.26310 (10)	0.0428 (5)
N6	0.31950 (13)	0.7291 (2)	0.32014 (10)	0.0385 (5)
N7	0.39487 (16)	0.8009 (3)	0.37649 (12)	0.0648 (7)
N8	0.21165 (13)	0.4829 (2)	0.11250 (10)	0.0447 (6)
N9	0.20157 (13)	0.3801 (2)	0.05837 (10)	0.0423 (5)
N10	0.1963 (2)	0.2815 (3)	0.00574 (13)	0.0815 (9)
C1	0.44295 (14)	0.2970 (3)	0.36860 (11)	0.0342 (6)
C2	0.55345 (15)	0.2892 (3)	0.42221 (13)	0.0468 (7)
C3	0.62099 (16)	0.3289 (3)	0.39169 (14)	0.0531 (8)
C4	0.57737 (15)	0.3711 (3)	0.30972 (13)	0.0482 (7)

supplementary materials

C5	0.46602 (15)	0.3750 (3)	0.25924 (12)	0.0403 (6)
C6	0.35865 (14)	0.2657 (3)	0.39230 (11)	0.0352 (6)
C7	0.02452 (13)	0.1678 (3)	0.13675 (10)	0.0341 (5)
C8	-0.08351 (14)	0.1241 (3)	0.10454 (12)	0.0467 (6)
C9	-0.14833 (16)	0.2628 (4)	0.11367 (13)	0.0573 (8)
C10	-0.10408 (16)	0.4361 (4)	0.15429 (12)	0.0512 (7)
C11	0.00463 (15)	0.4720 (3)	0.18414 (11)	0.0426 (6)
C12	0.10687 (14)	0.0314 (3)	0.13626 (10)	0.0340 (6)
H2	0.58190	0.25790	0.47780	0.0560*
H3	0.69590	0.32680	0.42690	0.0640*
H4	0.62190	0.39680	0.28820	0.0580*
H5	0.43630	0.40330	0.20330	0.0480*
H8	-0.11240	0.00400	0.07720	0.0560*
H9	-0.22170	0.23720	0.09210	0.0690*
H10	-0.14620	0.52930	0.16190	0.0610*
H11	0.03440	0.59240	0.21080	0.0510*
H12	0.4567 (19)	0.229 (3)	0.5093 (14)	0.060 (7)*
H13	0.3411 (17)	0.224 (3)	0.4844 (12)	0.049 (6)*
H14	0.1179 (17)	-0.220 (3)	0.1030 (13)	0.048 (7)*
H15	-0.001 (2)	-0.180 (3)	0.0684 (15)	0.076 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0266 (1)	0.0335 (1)	0.0333 (1)	-0.0016 (1)	0.0141 (1)	-0.0003 (1)
O1	0.0295 (7)	0.0589 (9)	0.0385 (7)	-0.0032 (6)	0.0149 (6)	0.0086 (6)
O2	0.0307 (7)	0.0327 (7)	0.0541 (8)	0.0009 (5)	0.0203 (6)	-0.0001 (6)
N1	0.0283 (7)	0.0316 (8)	0.0397 (8)	-0.0026 (6)	0.0170 (6)	-0.0001 (7)
N2	0.0427 (11)	0.0594 (12)	0.0359 (10)	-0.0036 (9)	0.0164 (9)	0.0035 (9)
N3	0.0282 (7)	0.0349 (9)	0.0314 (7)	0.0022 (6)	0.0136 (6)	0.0001 (7)
N4	0.0430 (10)	0.0347 (10)	0.0493 (10)	0.0040 (9)	0.0154 (8)	-0.0012 (8)
N5	0.0387 (9)	0.0392 (10)	0.0429 (9)	-0.0012 (7)	0.0154 (8)	-0.0075 (8)
N6	0.0427 (9)	0.0333 (9)	0.0391 (9)	0.0023 (8)	0.0206 (8)	-0.0010 (8)
N7	0.0520 (12)	0.0594 (12)	0.0527 (11)	-0.0042 (9)	0.0048 (9)	-0.0122 (10)
N8	0.0602 (10)	0.0388 (10)	0.0398 (9)	-0.0011 (8)	0.0290 (8)	-0.0004 (8)
N9	0.0484 (9)	0.0424 (10)	0.0381 (9)	-0.0028 (8)	0.0236 (8)	0.0049 (9)
N10	0.138 (2)	0.0638 (14)	0.0597 (13)	-0.0046 (14)	0.0629 (15)	-0.0130 (11)
C1	0.0309 (9)	0.0304 (10)	0.0373 (10)	-0.0016 (8)	0.0145 (8)	-0.0022 (8)
C2	0.0319 (10)	0.0545 (13)	0.0433 (11)	-0.0021 (9)	0.0114 (9)	-0.0047 (10)
C3	0.0271 (10)	0.0607 (15)	0.0608 (14)	0.0003 (9)	0.0147 (10)	-0.0090 (11)
C4	0.0366 (10)	0.0476 (12)	0.0687 (14)	-0.0008 (10)	0.0329 (10)	-0.0020 (11)
C5	0.0390 (10)	0.0381 (11)	0.0501 (11)	-0.0008 (9)	0.0274 (9)	0.0006 (10)
C6	0.0341 (10)	0.0321 (11)	0.0351 (10)	-0.0031 (8)	0.0146 (8)	0.0016 (8)
C7	0.0283 (8)	0.0435 (11)	0.0275 (8)	0.0007 (8)	0.0121 (7)	0.0025 (8)
C8	0.0320 (10)	0.0598 (13)	0.0411 (10)	-0.0076 (10)	0.0134 (8)	-0.0059 (10)
C9	0.0258 (10)	0.097 (2)	0.0460 (12)	0.0026 (11)	0.0162 (9)	-0.0008 (12)
C10	0.0369 (11)	0.0722 (16)	0.0451 (11)	0.0160 (11)	0.0214 (10)	0.0006 (11)
C11	0.0411 (10)	0.0473 (12)	0.0392 (10)	0.0116 (9)	0.0204 (9)	0.0022 (9)

C12 0.0360 (10) 0.0320 (10) 0.0317 (9) 0.0003 (8) 0.0157 (8) 0.0024 (8)

Geometric parameters (Å, °)

Ni1—O1	2.0701 (13)	N4—H14	0.80 (2)
Ni1—O2	2.0941 (12)	N4—H15	0.94 (3)
Ni1—N1	2.0685 (17)	C1—C6	1.502 (3)
Ni1—N3	2.0559 (17)	C1—C2	1.378 (3)
Ni1—N5	2.0652 (14)	C2—C3	1.381 (4)
Ni1—N8	2.0863 (17)	C3—C4	1.364 (3)
O1—C6	1.243 (2)	C4—C5	1.379 (3)
O2—C12	1.234 (3)	C7—C8	1.376 (3)
N1—C1	1.344 (2)	C7—C12	1.497 (3)
N1—C5	1.339 (3)	C8—C9	1.386 (3)
N2—C6	1.324 (3)	C9—C10	1.356 (4)
N3—C7	1.344 (2)	C10—C11	1.382 (3)
N3—C11	1.331 (3)	C2—H2	0.9300
N4—C12	1.328 (3)	C3—H3	0.9300
N5—N6	1.172 (2)	C4—H4	0.9300
N6—N7	1.163 (3)	C5—H5	0.9300
N8—N9	1.175 (2)	C8—H8	0.9300
N9—N10	1.159 (3)	C9—H9	0.9300
N2—H12	0.86 (3)	C10—H10	0.9300
N2—H13	0.85 (3)	C11—H11	0.9300
O1—Ni1—O2	93.30 (5)	C2—C1—C6	125.28 (17)
O1—Ni1—N1	78.45 (6)	C1—C2—C3	118.6 (2)
O1—Ni1—N3	89.76 (6)	C2—C3—C4	119.7 (2)
O1—Ni1—N5	88.69 (6)	C3—C4—C5	118.8 (2)
O1—Ni1—N8	175.91 (6)	N1—C5—C4	122.39 (18)
O2—Ni1—N1	94.34 (6)	O1—C6—N2	121.6 (2)
O2—Ni1—N3	78.08 (6)	N2—C6—C1	119.7 (2)
O2—Ni1—N5	173.44 (7)	O1—C6—C1	118.71 (16)
O2—Ni1—N8	89.94 (5)	C8—C7—C12	125.59 (18)
N1—Ni1—N3	165.67 (6)	N3—C7—C12	112.41 (17)
N1—Ni1—N5	92.18 (6)	N3—C7—C8	121.95 (19)
N1—Ni1—N8	98.83 (7)	C7—C8—C9	118.6 (2)
N3—Ni1—N5	95.69 (7)	C8—C9—C10	119.6 (2)
N3—Ni1—N8	93.36 (7)	C9—C10—C11	118.9 (2)
N5—Ni1—N8	88.36 (6)	N3—C11—C10	122.39 (19)
Ni1—O1—C6	114.90 (13)	N4—C12—C7	117.8 (2)
Ni1—O2—C12	114.68 (13)	O2—C12—N4	123.2 (2)
Ni1—N1—C1	114.71 (14)	O2—C12—C7	119.01 (17)
Ni1—N1—C5	126.71 (13)	C1—C2—H2	121.00
C1—N1—C5	118.28 (18)	C3—C2—H2	121.00
Ni1—N3—C7	115.52 (13)	C2—C3—H3	120.00
Ni1—N3—C11	125.88 (13)	C4—C3—H3	120.00
C7—N3—C11	118.55 (18)	C3—C4—H4	121.00
Ni1—N5—N6	123.67 (13)	C5—C4—H4	121.00
N5—N6—N7	178.2 (2)	N1—C5—H5	119.00

supplementary materials

Ni1—N8—N9	122.99 (12)	C4—C5—H5	119.00
N8—N9—N10	176.9 (3)	C7—C8—H8	121.00
H12—N2—H13	119 (2)	C9—C8—H8	121.00
C6—N2—H12	122.0 (18)	C8—C9—H9	120.00
C6—N2—H13	119.4 (14)	C10—C9—H9	120.00
C12—N4—H14	116.3 (16)	C9—C10—H10	121.00
C12—N4—H15	123.0 (15)	C11—C10—H10	121.00
H14—N4—H15	119 (2)	N3—C11—H11	119.00
N1—C1—C2	122.2 (2)	C10—C11—H11	119.00
N1—C1—C6	112.46 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H12 \cdots N7 ⁱ	0.86 (3)	2.12 (2)	2.967 (3)	165 (3)
N2—H13 \cdots N10 ⁱⁱ	0.85 (3)	2.31 (3)	3.154 (4)	172 (2)
N4—H14 \cdots N8 ⁱⁱⁱ	0.80 (2)	2.36 (2)	3.136 (3)	164 (2)
N4—H15 \cdots N10 ^{iv}	0.94 (3)	2.50 (3)	3.442 (4)	179 (3)
C2—H2 \cdots N7 ⁱ	0.93	2.61	3.516 (3)	163
C4—H4 \cdots O2 ^v	0.93	2.55	3.318 (3)	140
C8—H8 \cdots N10 ^{iv}	0.93	2.37	3.297 (3)	174
C10—H10 \cdots O1 ^{vi}	0.93	2.33	3.256 (3)	172

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, -y+1/2, z+1/2$; (iii) $x, y-1, z$; (iv) $-x, -y, -z$; (v) $-x+1, y+1/2, -z+1/2$; (vi) $-x, y+1/2, -z+1/2$.

Fig. 1




Fig. 2

