

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Norbornane-*exo-cis*-2,3-diyl 1',2'-phenylene orthocarbonate

Richard Betz and Peter Klüfers*

 Ludwig-Maximilians-Universität, Department Chemie und Biochemie, Butenandtstrasse 5–13, 81377 München, Germany
 Correspondence e-mail: kluef@cup.uni-muenchen.de


Received 3 October 2007; accepted 27 November 2007

 Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}—\text{C}) = 0.002$ Å; R factor = 0.045; wR factor = 0.119; data-to-parameter ratio = 15.7.

The title compound (systematic name: 4,7-methano-2,2'-spirobi[1,3-benzodioxole]), $\text{C}_{14}\text{H}_{14}\text{O}_4$, is an asymmetric spiro ester of orthocarbonic acid and two diols, *viz.* the aromatic benzene-1,2-diol and the aliphatic vicinal norbornane-*exo-cis*-2,3-diol. The orthocarbonate molecule is close to having non-crystallographic C_s symmetry. The five-membered ring stemming from the aliphatic diol has an envelope conformation. C—O bonds including the spiro-C atom span an approximately 0.07 Å range, but are within 0.02 Å of the respective distances in a density functional theory calculation, *i.e.* the distance difference is not caused by packing forces. Accordingly, the crystal packing is characterized by weak C—H \cdots O and C—H $\cdots\pi$ interactions.

Related literature

For the synthesis of the title compound, see: Komatsu *et al.* (1992). For related compounds, see: Betz & Klüfers (2007*a,b,c*); Betz *et al.* (2007). Density functional theory calculations were performed by Betz & Klüfers (2008).



Experimental

Crystal data

 $\text{C}_{14}\text{H}_{14}\text{O}_4$
 $M_r = 246.25$
 Monoclinic, $P2_1/n$
 $a = 7.9125$ (3) Å
 $b = 9.5545$ (5) Å
 $c = 15.2813$ (6) Å
 $\beta = 101.490$ (3) $^\circ$
 $V = 1132.11$ (9) Å 3
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm $^{-1}$
 $T = 200$ (2) K
 $0.30 \times 0.25 \times 0.16$ mm

Data collection

 Nonius KappaCCD diffractometer
 Absorption correction: none
 8399 measured reflections
 2582 independent reflections
 1716 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.119$
 $S = 1.05$
 2582 reflections
 164 parameters

 Only H-atom displacement parameters refined
 $\Delta\rho_{\text{max}} = 0.21$ e Å $^{-3}$
 $\Delta\rho_{\text{min}} = -0.19$ e Å $^{-3}$

Table 1

Selected bond lengths (Å).

O1—C1	1.367 (2)	O3—C1	1.412 (2)
O2—C1	1.370 (2)	O4—C1	1.435 (2)

Table 2

 Hydrogen-bond geometry (Å, $^\circ$).

 C_g is the centroid of the C9–C14 phenylene ring.

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C3—H3 \cdots O4 ⁱ	1.00	2.71	3.623 (2)	151
C7—H7A \cdots O4 ⁱ	0.99	2.76	3.663 (2)	151
C14—H14 \cdots O2 ⁱⁱ	0.95	2.61	3.501 (2)	156
C14—H14 \cdots O4 ⁱⁱ	0.95	2.88	3.514 (2)	125
C2—H2 \cdots C _g ⁱⁱⁱ	1.00	2.86	3.563 (2)	128
C5—H5 \cdots C _g ^{iv}	1.00	2.66	3.568 (2)	152

 Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x - \frac{1}{2}, -y - \frac{1}{2}, z - \frac{1}{2}$; (iv) $x + 1, y, z$.

Data collection: *COLLECT* (Nonius, 2004); cell refinement: *SCALEPACK* (Otwinowski & Minor 1997); data reduction: *DENZO* (Otwinowski & Minor 1997) and *SCALEPACK*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2003; 2006 version).

The authors thank Moritz Reichvilser for professional support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2027).

References

- Betz, R., Jahn, N. & Klüfers, P. (2007). *Acta Cryst.* **E63**, o4152.
 Betz, R. & Klüfers, P. (2007*a*). *Acta Cryst.* **E63**, o3933.
 Betz, R. & Klüfers, P. (2007*b*). *Acta Cryst.* **E63**, o4132.
 Betz, R. & Klüfers, P. (2007*c*). *Acta Cryst.* **E63**, o4300.
 Betz, R. & Klüfers, P. (2008). Unpublished results.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Komatsu, S., Takata, T. & Endo, T. (1992). *Macromolecules*, **25**, 7286–7293.
 Nonius (2004). *COLLECT*. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o399 [doi:10.1107/S1600536807063660]

Norbornane-*exo-cis*-2,3-diyl 1',2'-phenylene orthocarbonate

R. Betz and P. Klüfers

Comment

The title compound was prepared in order to compare its NMR-spectroscopic data with those of related silicon compounds.

In the molecule, a central carbon atom is chelated by a phenylene-1,2-dioxy and a norbornylene-*exo-cis*-2,3-dioxy moiety. The C—O bond lengths differ markedly (1.37 to 1.44 Å). About the same bond-length values were computed for the isolated molecule on the B3LYP/6-31+G(d,p) level of theory thus ruling out packing forces as the origin of the bond-length differences. The five-membered chelate ring stemming from the aliphatic diol adopts an envelope conformation on the spiro center C1 (puckering parameters: $Q_2 = 0.1274(16)$ Å, $\varphi_2 = 42.6(7)^\circ$ for the O1—C1—O2—C4—C3 ring).

Accordingly, the crystal packing is characterized by weak C—H \cdots X interactions whose H \cdots X distances are close to the sum of the van-der-Waals radii (vdWr). In terms of the vdWr criterion, the shortest tabulated hydrogen-bond, the C14—H14 \cdots O2 interaction, is 0.11 Å shorter than the radii sum. The weak interactions in (I) are thus less significant than those in the related 1-(ylomethyl)cyclopentyl 1',2'-phenylene orthocarbonate, where C—H \cdots O bonds are observed at the radii sum minus 0.35 Å (Betz & Klüfers, 2007c). Fig. 2 shows this interaction as well as the shortest C—H \cdots π bond which has one of the norbornane-bridgehead C—H functions as the donor. The other bridgehead methylidyne function acts as a donor in a still weaker bond. Moreover, another weak C—H \cdots O bond may be recognized with a diol-CH function as the donor (see the hydrogen bond table).

Experimental

The title compound was prepared based on a published procedure (Komatsu *et al.*, 1992) upon reaction of norbornane-*exo-cis*-2,3-diol with 2,2-dichlorobenzo[1.3]dioxol in dichloromethane in the presence of pyridine. Crystals suitable for X-ray analysis were obtained after recrystallization from boiling ethyl acetate.

Refinement

All H atoms were positioned geometrically and refined using a riding model, with C—H distances of 0.95, 0.99 and 1.00 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for all H atoms.

Figures




Fig. 1. The molecular structure of the title compound, showing displacement ellipsoids drawn at the 50% probability level.




Fig. 2. The crystal packing viewed along [0 1 0]. Green arrows: the strongest C—H...O interaction in terms of the H...O distance (C14—H14...O2). Green dotted lines: the strongest C—H... π interaction (C5—H5—C_g, C_g is the centroid of the phenylene residue).

4,7-methano-2,2'-spirobi[1,3-benzodioxole]

Crystal data

C ₁₄ H ₁₄ O ₄	$F_{000} = 520$
$M_r = 246.25$	$D_x = 1.445 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
Hall symbol: -P 2yn	$\lambda = 0.71073 \text{ \AA}$
$a = 7.9125 (3) \text{ \AA}$	Cell parameters from 14429 reflections
$b = 9.5545 (5) \text{ \AA}$	$\theta = 2.7\text{--}27.5^\circ$
$c = 15.2813 (6) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$\beta = 101.490 (3)^\circ$	$T = 200 (2) \text{ K}$
$V = 1132.11 (9) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.30 \times 0.25 \times 0.16 \text{ mm}$

Data collection

Nonius KappaCCD diffractometer	1716 reflections with $I > 2\sigma(I)$
Radiation source: rotating anode	$R_{\text{int}} = 0.054$
Monochromator: MONTEL, graded multilayered X-ray optics	$\theta_{\text{max}} = 27.5^\circ$
$T = 200(2) \text{ K}$	$\theta_{\text{min}} = 3.2^\circ$
CCD; rotation images; thick slices scans	$h = -9 \rightarrow 10$
Absorption correction: none	$k = -12 \rightarrow 11$
8399 measured reflections	$l = -19 \rightarrow 19$
2582 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	Only H-atom displacement parameters refined
$wR(F^2) = 0.119$	$w = 1/[\sigma^2(F_o^2) + (0.0569P)^2 + 0.1349P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
	$(\Delta/\sigma)_{\text{max}} = 0.001$

2582 reflections $\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$
 164 parameters $\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.39573 (14)	0.24072 (14)	0.47478 (7)	0.0460 (3)
O2	0.51722 (14)	0.20489 (14)	0.35623 (7)	0.0463 (3)
O3	0.27667 (13)	0.34038 (12)	0.34119 (7)	0.0408 (3)
O4	0.25466 (14)	0.10405 (11)	0.35921 (8)	0.0441 (3)
C1	0.3646 (2)	0.22289 (18)	0.38424 (11)	0.0384 (4)
C2	0.67414 (19)	0.32183 (17)	0.56079 (10)	0.0333 (4)
H2	0.6161	0.3766	0.6022	0.0490 (13)*
C3	0.56971 (19)	0.20081 (17)	0.51341 (10)	0.0342 (4)
H3	0.5742	0.1165	0.5526	0.0490 (13)*
C4	0.65286 (19)	0.17354 (18)	0.43193 (10)	0.0368 (4)
H4	0.6964	0.0755	0.4305	0.0490 (13)*
C5	0.7973 (2)	0.28116 (18)	0.44257 (11)	0.0378 (4)
H5	0.8407	0.3025	0.3869	0.0490 (13)*
C6	0.9350 (2)	0.2320 (2)	0.52188 (12)	0.0458 (5)
H6A	1.0433	0.2857	0.5259	0.0490 (13)*
H6B	0.9600	0.1310	0.5171	0.0490 (13)*
C7	0.8509 (2)	0.26146 (19)	0.60315 (11)	0.0439 (4)
H7A	0.8387	0.1743	0.6364	0.0490 (13)*
H7B	0.9196	0.3298	0.6443	0.0490 (13)*
C8	0.7165 (2)	0.40464 (17)	0.48246 (10)	0.0384 (4)
H8A	0.6123	0.4412	0.4420	0.0490 (13)*
H8B	0.7995	0.4814	0.5018	0.0490 (13)*
C9	0.12113 (18)	0.29423 (16)	0.29208 (9)	0.0307 (4)
C10	0.10837 (18)	0.15188 (16)	0.30274 (9)	0.0307 (4)
C11	-0.03269 (19)	0.07732 (17)	0.26098 (10)	0.0365 (4)
H11	-0.0406	-0.0211	0.2681	0.0490 (13)*
C12	-0.16353 (19)	0.15365 (18)	0.20764 (10)	0.0367 (4)
H12	-0.2643	0.1064	0.1778	0.0490 (13)*

supplementary materials

C13	-0.1506 (2)	0.29664 (18)	0.19700 (10)	0.0375 (4)
H13	-0.2426	0.3454	0.1600	0.0490 (13)*
C14	-0.0054 (2)	0.37112 (17)	0.23937 (10)	0.0350 (4)
H14	0.0049	0.4693	0.2320	0.0490 (13)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0254 (6)	0.0743 (9)	0.0367 (7)	0.0025 (5)	0.0027 (5)	-0.0082 (6)
O2	0.0296 (6)	0.0719 (9)	0.0354 (6)	0.0024 (6)	0.0016 (5)	-0.0142 (6)
O3	0.0331 (6)	0.0367 (7)	0.0486 (7)	-0.0050 (5)	-0.0018 (5)	-0.0036 (5)
O4	0.0333 (6)	0.0351 (7)	0.0556 (7)	0.0031 (5)	-0.0107 (5)	0.0028 (5)
C1	0.0293 (9)	0.0449 (10)	0.0388 (9)	0.0000 (7)	0.0014 (7)	-0.0057 (7)
C2	0.0337 (8)	0.0357 (9)	0.0301 (8)	0.0013 (7)	0.0053 (6)	-0.0024 (6)
C3	0.0283 (8)	0.0362 (9)	0.0369 (8)	-0.0013 (6)	0.0034 (6)	0.0023 (7)
C4	0.0305 (8)	0.0389 (9)	0.0389 (9)	0.0032 (7)	0.0016 (6)	-0.0092 (7)
C5	0.0297 (8)	0.0489 (10)	0.0358 (9)	-0.0022 (7)	0.0089 (6)	-0.0042 (7)
C6	0.0283 (9)	0.0509 (11)	0.0552 (11)	0.0031 (7)	0.0015 (7)	-0.0084 (8)
C7	0.0365 (9)	0.0506 (11)	0.0401 (10)	-0.0041 (8)	-0.0031 (7)	0.0004 (8)
C8	0.0419 (9)	0.0320 (9)	0.0419 (9)	-0.0018 (7)	0.0096 (7)	0.0023 (7)
C9	0.0267 (8)	0.0362 (9)	0.0293 (8)	-0.0017 (6)	0.0059 (6)	-0.0036 (6)
C10	0.0261 (8)	0.0336 (9)	0.0315 (8)	0.0037 (6)	0.0033 (6)	0.0010 (6)
C11	0.0320 (8)	0.0335 (9)	0.0432 (9)	-0.0027 (7)	0.0057 (6)	-0.0007 (7)
C12	0.0261 (8)	0.0478 (11)	0.0351 (8)	-0.0012 (7)	0.0030 (6)	-0.0050 (7)
C13	0.0318 (8)	0.0468 (11)	0.0326 (8)	0.0083 (7)	0.0035 (6)	0.0031 (7)
C14	0.0388 (9)	0.0328 (9)	0.0346 (8)	0.0043 (7)	0.0102 (7)	0.0046 (7)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.367 (2)	C5—H5	1.0000
O1—C3	1.438 (2)	C6—C7	1.547 (3)
O2—C1	1.370 (2)	C6—H6A	0.9900
O2—C4	1.444 (2)	C6—H6B	0.9900
O3—C9	1.380 (2)	C7—H7A	0.9900
O3—C1	1.412 (2)	C7—H7B	0.9900
O4—C10	1.377 (2)	C8—H8A	0.9900
O4—C1	1.435 (2)	C8—H8B	0.9900
C2—C3	1.518 (2)	C9—C14	1.367 (2)
C2—C8	1.527 (2)	C9—C10	1.376 (2)
C2—C7	1.532 (2)	C10—C11	1.370 (2)
C2—H2	1.0000	C11—C12	1.390 (2)
C3—C4	1.542 (2)	C11—H11	0.9500
C3—H3	1.0000	C12—C13	1.382 (2)
C4—C5	1.522 (2)	C12—H12	0.9500
C4—H4	1.0000	C13—C14	1.396 (2)
C5—C8	1.526 (2)	C13—H13	0.9500
C5—C6	1.533 (2)	C14—H14	0.9500
C1—O1—C3	110.32 (12)	C5—C6—H6A	111.1

C1—O2—C4	109.66 (12)	C7—C6—H6A	111.1
C9—O3—C1	107.62 (12)	C5—C6—H6B	111.1
C10—O4—C1	107.09 (12)	C7—C6—H6B	111.1
O1—C1—O2	109.88 (12)	H6A—C6—H6B	109.1
O1—C1—O3	110.18 (13)	C2—C7—C6	103.41 (13)
O2—C1—O3	109.73 (14)	C2—C7—H7A	111.1
O1—C1—O4	110.22 (14)	C6—C7—H7A	111.1
O2—C1—O4	110.04 (13)	C2—C7—H7B	111.1
O3—C1—O4	106.75 (11)	C6—C7—H7B	111.1
C3—C2—C8	101.79 (12)	H7A—C7—H7B	109.0
C3—C2—C7	106.20 (13)	C5—C8—C2	95.02 (13)
C8—C2—C7	101.65 (13)	C5—C8—H8A	112.7
C3—C2—H2	115.2	C2—C8—H8A	112.7
C8—C2—H2	115.2	C5—C8—H8B	112.7
C7—C2—H2	115.2	C2—C8—H8B	112.7
O1—C3—C2	112.31 (13)	H8A—C8—H8B	110.2
O1—C3—C4	103.94 (12)	C14—C9—C10	122.58 (14)
C2—C3—C4	103.56 (12)	C14—C9—O3	128.15 (15)
O1—C3—H3	112.1	C10—C9—O3	109.27 (13)
C2—C3—H3	112.1	C11—C10—C9	122.08 (14)
C4—C3—H3	112.1	C11—C10—O4	128.66 (14)
O2—C4—C5	111.59 (13)	C9—C10—O4	109.27 (12)
O2—C4—C3	104.08 (12)	C10—C11—C12	116.29 (15)
C5—C4—C3	103.40 (12)	C10—C11—H11	121.9
O2—C4—H4	112.4	C12—C11—H11	121.9
C5—C4—H4	112.4	C13—C12—C11	121.58 (14)
C3—C4—H4	112.4	C13—C12—H12	119.2
C4—C5—C8	101.52 (13)	C11—C12—H12	119.2
C4—C5—C6	106.13 (14)	C12—C13—C14	121.51 (14)
C8—C5—C6	102.10 (13)	C12—C13—H13	119.2
C4—C5—H5	115.1	C14—C13—H13	119.2
C8—C5—H5	115.1	C9—C14—C13	115.96 (15)
C6—C5—H5	115.1	C9—C14—H14	122.0
C5—C6—C7	103.11 (13)	C13—C14—H14	122.0
C3—O1—C1—O2	-14.26 (19)	O2—C4—C5—C6	-177.46 (13)
C3—O1—C1—O3	-135.27 (13)	C3—C4—C5—C6	71.25 (15)
C3—O1—C1—O4	107.18 (14)	C4—C5—C6—C7	-72.05 (16)
C4—O2—C1—O1	15.22 (19)	C8—C5—C6—C7	33.89 (17)
C4—O2—C1—O3	136.51 (13)	C3—C2—C7—C6	70.90 (16)
C4—O2—C1—O4	-106.32 (14)	C8—C2—C7—C6	-35.19 (16)
C9—O3—C1—O1	-119.98 (13)	C5—C6—C7—C2	0.83 (17)
C9—O3—C1—O2	118.91 (13)	C4—C5—C8—C2	54.90 (14)
C9—O3—C1—O4	-0.31 (16)	C6—C5—C8—C2	-54.59 (14)
C10—O4—C1—O1	120.17 (13)	C3—C2—C8—C5	-54.54 (14)
C10—O4—C1—O2	-118.49 (14)	C7—C2—C8—C5	54.98 (14)
C10—O4—C1—O3	0.52 (17)	C1—O3—C9—C14	-179.57 (15)
C1—O1—C3—C2	118.72 (15)	C1—O3—C9—C10	-0.02 (16)
C1—O1—C3—C4	7.43 (17)	C14—C9—C10—C11	0.0 (2)
C8—C2—C3—O1	-77.54 (15)	O3—C9—C10—C11	-179.53 (14)

supplementary materials

C7—C2—C3—O1	176.46 (13)	C14—C9—C10—O4	179.94 (14)
C8—C2—C3—C4	33.98 (15)	O3—C9—C10—O4	0.36 (17)
C7—C2—C3—C4	-72.02 (14)	C1—O4—C10—C11	179.34 (16)
C1—O2—C4—C5	-120.76 (14)	C1—O4—C10—C9	-0.54 (17)
C1—O2—C4—C3	-9.90 (17)	C9—C10—C11—C12	-0.6 (2)
O1—C3—C4—O2	1.48 (16)	O4—C10—C11—C12	179.57 (14)
C2—C3—C4—O2	-116.04 (13)	C10—C11—C12—C13	0.5 (2)
O1—C3—C4—C5	118.21 (13)	C11—C12—C13—C14	0.0 (2)
C2—C3—C4—C5	0.68 (15)	C10—C9—C14—C13	0.5 (2)
O2—C4—C5—C8	76.19 (15)	O3—C9—C14—C13	179.99 (14)
C3—C4—C5—C8	-35.11 (15)	C12—C13—C14—C9	-0.5 (2)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C3—H3...O4 ⁱ	1.00	2.71	3.623 (2)	151
C7—H7A...O4 ⁱ	0.99	2.76	3.663 (2)	151
C14—H14...O2 ⁱⁱ	0.95	2.61	3.501 (2)	156
C14—H14...O4 ⁱⁱ	0.95	2.88	3.514 (2)	125
C2—H2...Cg ⁱⁱⁱ	1.00	2.86	3.563 (2)	128
C5—H5...Cg ^{iv}	1.00	2.66	3.568 (2)	152

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x+1/2, y+1/2, -z+1/2$; (iii) $x-1/2, -y-1/2, z-1/2$; (iv) $x+1, y, z$.

Fig. 1




Fig. 2

