

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-Benzyl-6-phenylimino-5-(pyrrol-2-ylidene)hexahydropyrimidine-2,4-dione

 Rafael Tamazyan,<sup>a\*</sup> Armen Ayyazyan,<sup>a</sup> Vahan Martirosyan,<sup>b</sup> Kristine Avagyan<sup>b</sup> and Ashot Martirosyan<sup>b</sup>
<sup>a</sup>Molecule Structure Research Center, National Academy of Sciences RA, Azatutyan Avenue 26, 375014 Yerevan, Republic of Armenia, and <sup>b</sup>Institute of Fine Organic Chemistry, National Academy of Sciences RA, Azatutyan Avenue 26, 375014 Yerevan, Republic of Armenia

Correspondence e-mail: rafael@msrc.am


Received 12 December 2007; accepted 14 January 2008

 Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.065;  $wR$  factor = 0.129; data-to-parameter ratio = 15.8.

In the title compound,  $\text{C}_{21}\text{H}_{20}\text{N}_4\text{O}_2$ , a potential anti-human immunodeficiency virus type 1 (HIV-1) non-nucleoside reverse transcriptase inhibitor, the pyrrolidine ring adopts an envelope conformation, while the hydrogenated pyrimidine ring adopts a weakly expressed twist conformation. The molecules are connected into infinite chains *via*  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For related structures, see: Karapetyan *et al.* (2002); Tamazyan *et al.* (2002). For details of the pharmacological properties of similar compounds, see: De Clercq (1996).



## Experimental

### Crystal data

$\text{C}_{21}\text{H}_{20}\text{N}_4\text{O}_2$   
 $M_r = 360.41$   
 Triclinic,  $P\bar{1}$   
 $a = 5.7844$  (12) Å  
 $b = 10.378$  (2) Å  
 $c = 15.595$  (3) Å  
 $\alpha = 102.64$  (3)°  
 $\beta = 93.32$  (3)°  
 $\gamma = 102.45$  (3)°  
 $V = 886.6$  (3) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.09$  mm<sup>-1</sup>  
 $T = 293$  (2) K  
 $0.3 \times 0.27 \times 0.25$  mm

### Data collection

Enraf–Nonius CAD-4 diffractometer  
 Absorption correction: none  
 5619 measured reflections  
 5144 independent reflections  
 2944 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$   
 3 standard reflections  
 frequency: 180 min  
 intensity decay: none

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$   
 $wR(F^2) = 0.128$   
 $S = 1.04$   
 5144 reflections  
 325 parameters  
 All H-atom parameters refined  
 $\Delta\rho_{\text{max}} = 0.27$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.26$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N10}-\text{H10}\cdots\text{O7}$	0.90 (2)	2.03 (2)	2.664 (3)	127 (2)
$\text{N10}-\text{H10}\cdots\text{O7}^i$	0.90 (2)	2.22 (2)	2.891 (3)	131 (2)
$\text{N3}-\text{H3}\cdots\text{O8}^ii$	0.86 (2)	2.10 (2)	2.937 (3)	165 (2)

 Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $-x, -y, -z + 1$ .

Data collection: *CAD-4 Software* (Enraf–Nonius 1988); cell refinement: *CAD-4 Software*; data reduction: *HELENA* (Spek, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXTL*.

This research was carried out within the framework of the Armenian Science and Education Foundation (ANSEF, grant No. PS-chemorg-907). The authors express their thanks to ANSEF.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2060).

## References

- De Clercq, E. (1996). *Rev. Med. Virol.* **6**, 97–117.  
 Enraf–Nonius. (1988). *CAD-4 Software*. Version 5.0. Enraf–Nonius, Delft, The Netherlands.  
 Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.  
 Karapetyan, H., Tamazyan, R., Martirosyan, A., Hovhannesian, V. & Gasparyan, S. (2002). *Acta Cryst.* **C58**, o399–o401.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (1997). *HELENA*. University of Utrecht, The Netherlands.  
 Tamazyan, R., Karapetyan, H., Martirosyan, A., Hovhannesian, V. & Gasparyan, S. (2002). *Acta Cryst.* **C58**, o386–o388.

## supporting information

*Acta Cryst.* (2008). E64, o483 [doi:10.1107/S1600536808001384]

**1-Benzyl-6-phenylimino-5-(pyrrol-2-ylidene)hexahydropyrimidine-2,4-dione**

Rafael Tamazyan, Armen Ayvazyan, Vahan Martirosyan, Kristine Avagyan and Ashot Martirosyan

**S1. Comment**

The interest to X-ray structural investigation of the title compound, (I), was stimulated by its potentially HIV-1 RT inhibition properties. It belongs to the family of non-nucleoside reverse transcriptase inhibitors (NNRTIs).


A view of (I) with our numbering scheme is depicted in Fig. 1. All intramolecular interatomic distances in (I) are in good agreement with their mean statistical values. The crystal structure consists of infinite chains along  $[\bar{1}12]$  direction of crystal lattice. These chains are formed by molecules of (I) via  $O7\cdots H10-N10$  and  $O8\cdots H3-N3$  hydrogen bonds (Fig. 2).

**S2. Experimental**

The title compound was synthesized by the condensation of 6-anilino-1-benzyl-1,2,3,4-tetrahydro-2,4-pyrimidinedione with pyrrolidone-2. The crystals were grown from an ethanol solution. A suitable crystal of the size  $\sim 0.3$  mm was selected for X-ray diffraction experiment.


**S3. Refinement**

The positional parameters of all atoms, anisotropic displacement parameters of nonhydrogen atoms and isotropic thermal parameters of hydrogen atoms were refined without restraints.



**Figure 1**

A view of (I) with the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Unit cell packing of (I) showing infinite chains of molecules *via* hydrogen bonding; for clarity only H atoms participating in bonding have been included. Symmetry cods: (i)  $1 - x, 1 - y, 1 - z$ ; (ii)  $1 + x, 1 + y, z$ ; (iii)  $-x, -y, 1 - z$ .

### 1-Benzyl-6-phenylimino-5-(pyrrol-2-ylidene)hexahydropyrimidine-2,4-dione

#### Crystal data

$C_{21}H_{20}N_4O_2$

$M_r = 360.41$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 5.7844$  (12) Å

$b = 10.378$  (2) Å

$c = 15.595$  (3) Å

$\alpha = 102.64$  (3)°

$\beta = 93.32$  (3)°

$\gamma = 102.45$  (3)°

$V = 886.6$  (3) Å<sup>3</sup>

$Z = 2$

$F(000) = 380$

$D_x = 1.350$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 22 reflections

$\theta = 13$ – $16^\circ$

$\mu = 0.09$  mm<sup>-1</sup>

$T = 293$  K

Prism, colourless

$0.3 \times 0.27 \times 0.25$  mm

#### Data collection

Enraf–Nonius CAD-4

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\theta/2\theta$  scans

5619 measured reflections

5144 independent reflections  
 2944 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$   
 $\theta_{\text{max}} = 30.0^\circ$ ,  $\theta_{\text{min}} = 1.4^\circ$   
 $h = 0 \rightarrow 8$

$k = -14 \rightarrow 14$   
 $l = -21 \rightarrow 21$   
 3 standard reflections every 180 min  
 intensity decay: none

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.065$   
 $wR(F^2) = 0.128$   
 $S = 1.04$   
 5144 reflections  
 325 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: difference Fourier map  
 All H-atom parameters refined  
 $w = 1/[\sigma^2(F_o^2) + (0.0198P)^2 + 0.561P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.27 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.26 \text{ e } \text{\AA}^{-3}$   
 Extinction correction: *SHELXL97* (Sheldrick,  
 2008),  $F_c^* = kFc^*[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.0124 (13)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.4204 (3)	-0.03850 (16)	0.35608 (11)	0.0320 (4)
C2	0.2737 (4)	-0.0209 (2)	0.42132 (14)	0.0329 (5)
H3	0.163 (4)	0.120 (2)	0.4942 (16)	0.043 (7)*
N3	0.2672 (4)	0.11039 (18)	0.45813 (13)	0.0374 (5)
C4	0.4114 (4)	0.2259 (2)	0.44199 (13)	0.0312 (5)
C5	0.5484 (4)	0.20422 (19)	0.36835 (13)	0.0291 (4)
C6	0.5188 (4)	0.0671 (2)	0.31427 (14)	0.0295 (4)
O7	0.4116 (3)	0.33653 (15)	0.49159 (10)	0.0430 (4)
O8	0.1540 (3)	-0.11624 (15)	0.44595 (11)	0.0469 (4)
C9	0.7255 (4)	0.3145 (2)	0.35785 (13)	0.0300 (4)
H10	0.638 (4)	0.464 (2)	0.4363 (16)	0.044 (7)*
N10	0.7427 (4)	0.44206 (18)	0.39838 (13)	0.0393 (5)
C11	0.9511 (5)	0.5384 (2)	0.3816 (2)	0.0465 (6)
H11A	1.017 (5)	0.610 (3)	0.438 (2)	0.075 (9)*
H11B	0.906 (5)	0.585 (3)	0.3357 (18)	0.060 (8)*
C12	1.1099 (5)	0.4461 (3)	0.34836 (19)	0.0462 (6)
H12A	1.208 (5)	0.435 (3)	0.4041 (19)	0.066 (9)*
H12B	1.215 (5)	0.476 (3)	0.3058 (17)	0.058 (8)*

C13	0.9361 (4)	0.3101 (3)	0.30583 (17)	0.0401 (6)
H13A	0.993 (5)	0.228 (3)	0.3025 (17)	0.054 (8)*
H13B	0.890 (5)	0.310 (3)	0.245 (2)	0.071 (9)*
N14	0.5687 (3)	0.02327 (18)	0.23576 (12)	0.0367 (4)
C15	0.5951 (4)	0.0974 (2)	0.16998 (14)	0.0342 (5)
C16	0.4413 (5)	0.1784 (3)	0.15504 (17)	0.0457 (6)
H16	0.325 (4)	0.192 (2)	0.1957 (16)	0.048 (7)*
C17	0.4605 (6)	0.2394 (3)	0.08445 (19)	0.0578 (8)
H17	0.351 (5)	0.292 (3)	0.0734 (19)	0.069 (9)*
C18	0.6315 (6)	0.2211 (3)	0.02769 (18)	0.0593 (8)
H18	0.646 (5)	0.263 (3)	-0.024 (2)	0.072 (9)*
C19	0.7829 (5)	0.1407 (3)	0.04185 (17)	0.0533 (7)
H19	0.904 (5)	0.127 (3)	0.002 (2)	0.072 (9)*
C20	0.7642 (5)	0.0783 (3)	0.11150 (16)	0.0439 (6)
H20	0.866 (5)	0.023 (3)	0.1218 (17)	0.056 (8)*
C21	0.4193 (4)	-0.1793 (2)	0.31229 (15)	0.0350 (5)
H21A	0.575 (4)	-0.179 (2)	0.2917 (14)	0.033 (6)*
H21B	0.403 (4)	-0.229 (2)	0.3603 (15)	0.037 (6)*
C22	0.2243 (4)	-0.2461 (2)	0.23746 (14)	0.0334 (5)
C23	0.2296 (5)	-0.3733 (2)	0.18596 (16)	0.0417 (6)
H23	0.367 (5)	-0.412 (3)	0.1979 (17)	0.058 (8)*
C24	0.0439 (5)	-0.4443 (3)	0.12153 (18)	0.0528 (7)
H24	0.049 (5)	-0.533 (3)	0.0869 (17)	0.059 (8)*
C25	-0.1440 (5)	-0.3884 (3)	0.10493 (18)	0.0546 (7)
H25	-0.276 (5)	-0.439 (3)	0.0577 (18)	0.064 (8)*
C26	-0.1472 (5)	-0.2606 (3)	0.15307 (17)	0.0482 (6)
H26	-0.285 (5)	-0.219 (3)	0.1429 (17)	0.061 (8)*
C27	0.0355 (5)	-0.1909 (2)	0.21906 (16)	0.0415 (6)
H27	0.030 (4)	-0.103 (2)	0.2539 (15)	0.038 (6)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0374 (10)	0.0241 (8)	0.0323 (9)	0.0033 (7)	0.0113 (8)	0.0042 (7)
C2	0.0368 (12)	0.0286 (10)	0.0328 (11)	0.0050 (9)	0.0096 (9)	0.0071 (9)
N3	0.0432 (11)	0.0282 (9)	0.0404 (11)	0.0049 (8)	0.0219 (9)	0.0063 (8)
C4	0.0339 (12)	0.0288 (10)	0.0307 (11)	0.0062 (9)	0.0062 (9)	0.0066 (9)
C5	0.0318 (11)	0.0260 (10)	0.0284 (10)	0.0030 (8)	0.0086 (9)	0.0062 (8)
C6	0.0288 (11)	0.0259 (10)	0.0330 (11)	0.0030 (8)	0.0090 (9)	0.0072 (8)
O7	0.0566 (11)	0.0281 (8)	0.0424 (9)	0.0071 (7)	0.0204 (8)	0.0032 (7)
O8	0.0581 (11)	0.0319 (8)	0.0532 (10)	0.0052 (8)	0.0276 (9)	0.0153 (7)
C9	0.0301 (11)	0.0318 (11)	0.0268 (10)	0.0046 (9)	0.0026 (8)	0.0069 (8)
N10	0.0418 (12)	0.0279 (9)	0.0462 (12)	0.0024 (8)	0.0149 (10)	0.0079 (8)
C11	0.0464 (15)	0.0323 (12)	0.0540 (16)	-0.0060 (11)	0.0077 (13)	0.0106 (12)
C12	0.0374 (14)	0.0424 (14)	0.0541 (16)	-0.0033 (11)	0.0071 (12)	0.0136 (12)
C13	0.0340 (13)	0.0371 (13)	0.0453 (14)	0.0011 (10)	0.0120 (11)	0.0067 (11)
N14	0.0441 (11)	0.0309 (9)	0.0336 (10)	0.0047 (8)	0.0141 (8)	0.0063 (8)
C15	0.0361 (12)	0.0310 (11)	0.0289 (11)	-0.0022 (9)	0.0075 (9)	0.0022 (9)

C16	0.0410 (14)	0.0533 (15)	0.0411 (14)	0.0068 (12)	0.0079 (12)	0.0111 (11)
C17	0.0546 (18)	0.0641 (18)	0.0554 (17)	0.0070 (15)	-0.0062 (14)	0.0257 (15)
C18	0.066 (2)	0.0638 (18)	0.0395 (15)	-0.0107 (15)	-0.0006 (14)	0.0215 (13)
C19	0.0581 (18)	0.0554 (16)	0.0340 (13)	-0.0086 (14)	0.0138 (13)	0.0035 (12)
C20	0.0500 (15)	0.0402 (13)	0.0382 (13)	0.0056 (12)	0.0165 (12)	0.0045 (11)
C21	0.0407 (13)	0.0280 (11)	0.0371 (12)	0.0089 (10)	0.0116 (10)	0.0065 (9)
C22	0.0390 (12)	0.0267 (10)	0.0333 (11)	0.0026 (9)	0.0114 (10)	0.0081 (8)
C23	0.0543 (16)	0.0312 (12)	0.0383 (13)	0.0081 (11)	0.0136 (12)	0.0050 (10)
C24	0.0671 (19)	0.0366 (14)	0.0437 (14)	0.0029 (13)	0.0119 (14)	-0.0061 (11)
C25	0.0545 (18)	0.0550 (17)	0.0405 (14)	-0.0035 (14)	0.0009 (13)	-0.0001 (12)
C26	0.0418 (15)	0.0514 (15)	0.0472 (15)	0.0050 (12)	0.0026 (12)	0.0094 (12)
C27	0.0456 (14)	0.0316 (12)	0.0443 (13)	0.0068 (10)	0.0089 (11)	0.0040 (10)

*Geometric parameters (Å, °)*

N1—C2	1.369 (3)	C15—C16	1.391 (3)
N1—C6	1.427 (2)	C15—C20	1.392 (3)
N1—C21	1.470 (3)	C16—C17	1.383 (4)
C2—O8	1.228 (2)	C16—H16	0.96 (3)
C2—N3	1.368 (3)	C17—C18	1.381 (4)
N3—C4	1.386 (3)	C17—H17	0.95 (3)
N3—H3	0.86 (2)	C18—C19	1.373 (4)
C4—O7	1.236 (2)	C18—H18	0.99 (3)
C4—C5	1.437 (3)	C19—C20	1.379 (4)
C5—C9	1.408 (3)	C19—H19	0.97 (3)
C5—C6	1.455 (3)	C20—H20	0.93 (3)
C6—N14	1.282 (3)	C21—C22	1.507 (3)
C9—N10	1.316 (3)	C21—H21A	0.97 (2)
C9—C13	1.506 (3)	C21—H21B	0.99 (2)
N10—C11	1.467 (3)	C22—C27	1.381 (3)
N10—H10	0.90 (2)	C22—C23	1.395 (3)
C11—C12	1.499 (4)	C23—C24	1.384 (4)
C11—H11A	1.01 (3)	C23—H23	0.99 (3)
C11—H11B	1.00 (3)	C24—C25	1.375 (4)
C12—C13	1.531 (3)	C24—H24	0.96 (3)
C12—H12A	1.05 (3)	C25—C26	1.379 (4)
C12—H12B	0.98 (3)	C25—H25	1.00 (3)
C13—H13A	0.97 (3)	C26—C27	1.384 (4)
C13—H13B	0.97 (3)	C26—H26	1.00 (3)
N14—C15	1.407 (3)	C27—H27	0.96 (2)
C2—N1—C6	122.57 (17)	C16—C15—C20	118.3 (2)
C2—N1—C21	116.70 (17)	C16—C15—N14	122.3 (2)
C6—N1—C21	118.51 (17)	C20—C15—N14	119.0 (2)
O8—C2—N3	121.31 (19)	C17—C16—C15	120.3 (3)
O8—C2—N1	122.72 (19)	C17—C16—H16	121.3 (15)
N3—C2—N1	115.97 (18)	C15—C16—H16	118.4 (15)
C2—N3—C4	126.16 (19)	C18—C17—C16	120.8 (3)

C2—N3—H3	115.4 (16)	C18—C17—H17	119.4 (18)
C4—N3—H3	118.5 (16)	C16—C17—H17	119.7 (18)
O7—C4—N3	117.81 (19)	C19—C18—C17	119.1 (3)
O7—C4—C5	126.25 (19)	C19—C18—H18	119.3 (17)
N3—C4—C5	115.94 (18)	C17—C18—H18	121.6 (17)
C9—C5—C4	118.04 (18)	C18—C19—C20	120.6 (3)
C9—C5—C6	122.50 (18)	C18—C19—H19	120.2 (18)
C4—C5—C6	119.06 (17)	C20—C19—H19	119.2 (18)
N14—C6—N1	113.57 (18)	C19—C20—C15	120.8 (3)
N14—C6—C5	131.61 (19)	C19—C20—H20	121.4 (16)
N1—C6—C5	114.81 (17)	C15—C20—H20	117.7 (16)
N10—C9—C5	124.5 (2)	N1—C21—C22	115.08 (19)
N10—C9—C13	107.55 (19)	N1—C21—H21A	107.5 (13)
C5—C9—C13	127.79 (19)	C22—C21—H21A	110.5 (13)
C9—N10—C11	114.7 (2)	N1—C21—H21B	104.6 (13)
C9—N10—H10	120.3 (16)	C22—C21—H21B	110.5 (13)
C11—N10—H10	124.8 (16)	H21A—C21—H21B	108.3 (18)
N10—C11—C12	101.68 (19)	C27—C22—C23	118.0 (2)
N10—C11—H11A	109.7 (17)	C27—C22—C21	123.7 (2)
C12—C11—H11A	115.6 (17)	C23—C22—C21	118.2 (2)
N10—C11—H11B	110.5 (16)	C24—C23—C22	120.5 (3)
C12—C11—H11B	111.4 (16)	C24—C23—H23	121.3 (16)
H11A—C11—H11B	108 (2)	C22—C23—H23	118.2 (16)
C11—C12—C13	103.9 (2)	C25—C24—C23	120.6 (3)
C11—C12—H12A	106.9 (15)	C25—C24—H24	119.8 (16)
C13—C12—H12A	108.7 (15)	C23—C24—H24	119.5 (16)
C11—C12—H12B	115.1 (15)	C24—C25—C26	119.6 (3)
C13—C12—H12B	110.6 (15)	C24—C25—H25	120.6 (16)
H12A—C12—H12B	111 (2)	C26—C25—H25	119.9 (16)
C9—C13—C12	103.0 (2)	C25—C26—C27	119.9 (3)
C9—C13—H13A	113.3 (15)	C25—C26—H26	120.6 (16)
C12—C13—H13A	117.7 (16)	C27—C26—H26	119.4 (16)
C9—C13—H13B	109.5 (18)	C22—C27—C26	121.4 (2)
C12—C13—H13B	107.9 (17)	C22—C27—H27	118.9 (14)
H13A—C13—H13B	105 (2)	C26—C27—H27	119.7 (14)
C6—N14—C15	125.54 (19)		
C6—N1—C2—O8	166.1 (2)	N10—C9—C13—C12	18.0 (3)
C21—N1—C2—O8	3.3 (3)	C5—C9—C13—C12	-157.6 (2)
C6—N1—C2—N3	-13.8 (3)	C11—C12—C13—C9	-28.8 (3)
C21—N1—C2—N3	-176.6 (2)	N1—C6—N14—C15	158.8 (2)
O8—C2—N3—C4	173.1 (2)	C5—C6—N14—C15	-22.4 (4)
N1—C2—N3—C4	-7.0 (3)	C6—N14—C15—C16	-42.4 (3)
C2—N3—C4—O7	-167.7 (2)	C6—N14—C15—C20	144.7 (2)
C2—N3—C4—C5	12.0 (3)	C20—C15—C16—C17	-1.0 (4)
O7—C4—C5—C9	9.8 (3)	N14—C15—C16—C17	-174.0 (2)
N3—C4—C5—C9	-169.9 (2)	C15—C16—C17—C18	0.1 (4)
O7—C4—C5—C6	-177.4 (2)	C16—C17—C18—C19	0.3 (4)



N3—C4—C5—C6	3.0 (3)	C17—C18—C19—C20	0.3 (4)
C2—N1—C6—N14	-154.0 (2)	C18—C19—C20—C15	-1.3 (4)
C21—N1—C6—N14	8.5 (3)	C16—C15—C20—C19	1.6 (3)
C2—N1—C6—C5	27.1 (3)	N14—C15—C20—C19	174.8 (2)
C21—N1—C6—C5	-170.4 (2)	C2—N1—C21—C22	84.4 (2)
C9—C5—C6—N14	-26.9 (4)	C6—N1—C21—C22	-79.1 (2)
C4—C5—C6—N14	160.5 (2)	N1—C21—C22—C27	-10.3 (3)
C9—C5—C6—N1	151.8 (2)	N1—C21—C22—C23	172.27 (19)
C4—C5—C6—N1	-20.7 (3)	C27—C22—C23—C24	-3.3 (3)
C4—C5—C9—N10	-15.1 (3)	C21—C22—C23—C24	174.3 (2)
C6—C5—C9—N10	172.4 (2)	C22—C23—C24—C25	2.5 (4)
C4—C5—C9—C13	159.9 (2)	C23—C24—C25—C26	-0.1 (4)
C6—C5—C9—C13	-12.7 (4)	C24—C25—C26—C27	-1.6 (4)
C5—C9—N10—C11	176.1 (2)	C23—C22—C27—C26	1.6 (3)
C13—C9—N10—C11	0.3 (3)	C21—C22—C27—C26	-175.8 (2)
C9—N10—C11—C12	-18.9 (3)	C25—C26—C27—C22	0.8 (4)
N10—C11—C12—C13	28.4 (3)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N10—H10 $\cdots$ O7	0.90 (2)	2.03 (2)	2.664 (3)	127 (2)
N10—H10 $\cdots$ O7 <sup>i</sup>	0.90 (2)	2.22 (2)	2.891 (3)	131 (2)
N3—H3 $\cdots$ O8 <sup>ii</sup>	0.86 (2)	2.10 (2)	2.937 (3)	165 (2)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x, -y, -z+1$ .