

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 2-[(phenylsulfonyl)methyl]-1-(phenylsulfonyl)-1*H*-indole-3-carboxylateG. Chakkaravarthi,^{a*} V. Dhayalan,^b
A. K. Mohanakrishnan^b and V. Manivannan^c^aDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India,^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India, and ^cDepartment of Physics, Presidency College, Chennai 600 005, India

Correspondence e-mail: chakkaravarthi_2005@yahoo.com


Received 22 December 2007; accepted 29 December 2007

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.048; wR factor = 0.123; data-to-parameter ratio = 14.6.

In the title compound, $\text{C}_{24}\text{H}_{21}\text{NO}_4\text{S}_2$, the phenyl rings form dihedral angles of 85.77 (9) and 85.22 (9)° and the ester group forms an angle of 12.61 (10)° with the indane ring. The molecular structure is stabilized by weak intramolecular C—H...O interactions.

Related literature

For related literature, see: Allen *et al.* (1987); Nieto *et al.* (2005); Satis Kumar *et al.* (2006). A similar compound has been reported by Chakkaravarthi *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{24}\text{H}_{21}\text{NO}_4\text{S}_2$
 $M_r = 451.56$ Monoclinic, $P2_1/c$
 $a = 11.745$ (1) Å $b = 7.7140$ (2) Å
 $c = 26.770$ (1) Å
 $\beta = 116.020$ (2)°
 $V = 2179.6$ (2) Å³
 $Z = 4$ Mo $K\alpha$ radiation
 $\mu = 0.28$ mm⁻¹
 $T = 295$ (2) K
 $0.24 \times 0.20 \times 0.20$ mm

Data collection

Bruker Kappa APEX2
diffractometer
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.927$, $T_{\max} = 0.947$ 9731 measured reflections
4116 independent reflections
2128 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.122$
 $S = 0.82$
4116 reflections
281 parameters9 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.23$ e Å⁻³
 $\Delta\rho_{\min} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C7-H7A\cdots O1$	0.97	2.38	2.813 (4)	107
$C7-H7B\cdots O3$	0.97	2.26	2.977 (4)	130
$C11-H11\cdots O4$	0.93	2.42	2.932 (3)	115
$C14-H14\cdots O2$	0.93	2.41	2.966 (3)	119
$C24-H24\cdots O2$	0.93	2.52	2.898 (3)	105

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2; data reduction: APEX2; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2072).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2004). APEX2. Version 1.0-27. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chakkaravarthi, G., Ramesh, N., Mohanakrishnan, A. K. & Manivannan, V. (2007). *Acta Cryst. E* **63**, o3564.
- Nieto, M. J., Alovero, F. L., Manzo, R. H. & Mazzieri, M. R. (2005). *Eur. J. Med. Chem.* **40**, 361–369.
- Satis Kumar, B. K., Gayathri, D., Velmurugan, D., Ravikumar, K. & Poornachandran, M. (2006). *Acta Cryst. E* **62**, o5388–o5389.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o392 [doi:10.1107/S1600536807068778]

Ethyl 2-[(phenylsulfanyl)methyl]-1-(phenylsulfonyl)-1*H*-indole-3-carboxylate

G. Chakkaravarthi, V. Dhayalan, A. K. Mohanakrishnan and V. Manivannan

Comment

The benzenesulfonamide derivatives exhibit significant biological activities, such as antibacterial (Nieto *et al.*, 2005). In the molecule of title compound (I) (Fig. 1), the bond lengths and angles are agree with the reported similar structures (Allen *et al.* 1987; Chakkaravarthi *et al.*, 2007; Satis Kumar *et al.*, 2006).

The phenyl rings C1–C6 and C19–C24 form the dihedral angles with the indane ring system of 85.77 (9)° and 85.22 (9)°, respectively. The dihedral angle between these phenyl rings is 4.26 (9)°. The ester group C16/O3/O4/C17/C18 forms a dihedral angle with the indane ring system of 12.61 (10)°.

The molecular structure of (I) (Fig. 1) is stabilized by intramolecular C—H···O interactions. A similar compound has been reported by Chakkaravarthi *et al.* (2007).

Experimental

To a well stirred suspension of sodium hydride (0.17 g, 3.55 mmol) in dry tetrahydrofuran (THF) (10 ml) at 273 K, a solution of thiophenol (0.36 ml, 3.55 mmol) in dry THF (10 ml) was slowly added over a period of 10 min. After the evaluation of hydrogen gas ceased a solution of ethyl 2-(bromomethyl)-1-(phenylsulfonyl)-1*H*-indole-3-carboxylate (1 g, 2.36 mmol) in dry THF (10 ml) was added in dropwise with vigorous stirring. Then the reaction mixture was stirred for 2 h and then poured over crushed ice. The precipitated solid was filtered, washed with water and dried over calcium chloride. The crude sulfide was recrystallized from methanol. Single crystals suitable for X-ray analysis were grown by slow evaporation of a methanol solution at room temperature.

Refinement

H atoms were positioned geometrically and refined using riding model with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic CH, C—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH₂ and C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for CH₃. The bond distances C19—C20, C19—24, C20—C21, C21—C22, C22—C23, C23—C24, C3—C4, C4—C5 and C5—C6 were restrained to be 1.39 (3) Å and the bond distance C17—C18 was restrained to be 1.55 (3) Å.

Figures




Fig. 1. The molecular structure of (I), with atom labeling scheme. Displacement ellipsoids are drawn at 50% probability level. H atoms are presented as a small spheres of arbitrary radius. Intramolecular H-bonds are shown as dotted lines.

Ethyl 2-[(phenylsulfanyl)methyl]-1-(phenylsulfonyl)-1*H*-indole- 3-carboxylate

Crystal data

$C_{24}H_{21}NO_4S_2$

$M_r = 451.56$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 11.745$ (1) Å

$b = 7.7140$ (2) Å

$c = 26.770$ (1) Å

$\beta = 116.020$ (2)°

$V = 2179.6$ (2) Å³

$Z = 4$

$F_{000} = 944$

$D_x = 1.376$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 10291 reflections

$\theta = 2.8$ – 26.0 °

$\mu = 0.28$ mm⁻¹

$T = 295$ (2) K

Block, colourless

$0.24 \times 0.20 \times 0.20$ mm

Data collection

Bruker Kappa APEX2
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295$ (2) K

ω - and φ -scan

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.927$, $T_{\max} = 0.947$

9731 measured reflections

4116 independent reflections

2128 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

$\theta_{\text{max}} = 28.7$ °

$\theta_{\text{min}} = 1.9$ °

$h = -15 \rightarrow 15$

$k = -10 \rightarrow 8$

$l = -36 \rightarrow 35$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.122$	$w = 1/[\sigma^2(F_o^2) + (0.0481P)^2]$
$S = 0.82$	where $P = (F_o^2 + 2F_c^2)/3$
4116 reflections	$(\Delta/\sigma)_{\max} < 0.001$
281 parameters	$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
9 restraints	$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All s.u.'s (except the s.u.'s in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.75890 (7)	0.46511 (8)	0.02189 (3)	0.03776 (17)
S2	1.06434 (7)	0.72740 (9)	0.09115 (4)	0.0563 (2)
O1	0.84440 (19)	0.5349 (2)	0.00269 (8)	0.0471 (5)
O2	0.64341 (19)	0.5547 (2)	0.01066 (8)	0.0494 (5)
O3	1.2184 (2)	0.3917 (3)	0.22110 (10)	0.0725 (6)
O4	1.11279 (18)	0.3805 (3)	0.27290 (8)	0.0545 (5)
N1	0.8387 (2)	0.4502 (2)	0.09151 (9)	0.0362 (5)
C1	1.2195 (3)	0.7980 (3)	0.13616 (12)	0.0438 (6)
C2	1.2297 (3)	0.9673 (4)	0.15435 (13)	0.0546 (7)
H2	1.1570	1.0329	0.1457	0.066*
C3	1.3477 (3)	1.0396 (4)	0.18538 (15)	0.0661 (9)
H3	1.3537	1.1543	0.1970	0.079*
C4	1.4568 (3)	0.9438 (4)	0.19933 (15)	0.0700 (10)
H4	1.5360	0.9927	0.2203	0.084*
C5	1.4459 (3)	0.7738 (4)	0.18144 (14)	0.0628 (9)
H5	1.5184	0.7070	0.1911	0.075*
C6	1.3281 (2)	0.7019 (4)	0.14923 (13)	0.0552 (7)
H6	1.3221	0.5886	0.1364	0.066*
C7	1.0649 (2)	0.4937 (3)	0.10300 (12)	0.0423 (6)
H7A	1.0453	0.4325	0.0685	0.051*
H7B	1.1486	0.4581	0.1300	0.051*
C8	0.9695 (2)	0.4468 (3)	0.12382 (11)	0.0362 (6)
C9	0.9929 (2)	0.4064 (3)	0.17792 (10)	0.0367 (6)

supplementary materials

C10	0.8719 (2)	0.3803 (3)	0.17934 (11)	0.0368 (6)
C11	0.8367 (3)	0.3384 (4)	0.22147 (12)	0.0470 (6)
H11	0.8978	0.3236	0.2579	0.056*
C12	0.7090 (3)	0.3194 (4)	0.20795 (13)	0.0539 (7)
H12	0.6848	0.2903	0.2356	0.065*
C13	0.6167 (3)	0.3430 (4)	0.15376 (13)	0.0513 (7)
H13	0.5318	0.3299	0.1459	0.062*
C14	0.6486 (3)	0.3858 (3)	0.11120 (12)	0.0431 (6)
H14	0.5871	0.4008	0.0749	0.052*
C15	0.7775 (2)	0.4055 (3)	0.12549 (10)	0.0358 (5)
C16	1.1197 (3)	0.3927 (3)	0.22444 (12)	0.0442 (6)
C17	1.2305 (3)	0.3585 (4)	0.32177 (12)	0.0626 (8)
H17A	1.2932	0.4401	0.3216	0.075*
H17B	1.2627	0.2419	0.3233	0.075*
C18	1.2026 (4)	0.3919 (5)	0.37179 (13)	0.0733 (10)
H18A	1.1788	0.5110	0.3717	0.110*
H18B	1.2770	0.3676	0.4055	0.110*
H18C	1.1346	0.3182	0.3694	0.110*
C19	0.7233 (2)	0.2500 (3)	-0.00112 (10)	0.0379 (5)
C20	0.8218 (3)	0.1388 (3)	0.00635 (12)	0.0507 (7)
H20	0.9054	0.1755	0.0254	0.061*
C21	0.7946 (3)	-0.0279 (3)	-0.01486 (13)	0.0594 (9)
H21	0.8597	-0.1040	-0.0105	0.071*
C22	0.6695 (3)	-0.0800 (4)	-0.04255 (13)	0.0607 (9)
H22	0.6507	-0.1915	-0.0571	0.073*
C23	0.5720 (3)	0.0317 (3)	-0.04881 (15)	0.0663 (9)
H23	0.4885	-0.0057	-0.0674	0.080*
C24	0.5977 (2)	0.1988 (3)	-0.02768 (14)	0.0576 (8)
H24	0.5326	0.2739	-0.0312	0.069*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0378 (4)	0.0385 (3)	0.0341 (4)	-0.0018 (3)	0.0132 (3)	0.0011 (3)
S2	0.0334 (4)	0.0581 (4)	0.0664 (6)	-0.0052 (3)	0.0118 (4)	0.0160 (4)
O1	0.0506 (12)	0.0494 (10)	0.0405 (11)	-0.0102 (9)	0.0194 (11)	0.0004 (8)
O2	0.0432 (12)	0.0476 (10)	0.0507 (13)	0.0090 (8)	0.0146 (11)	0.0063 (9)
O3	0.0350 (12)	0.1220 (18)	0.0579 (15)	0.0079 (12)	0.0179 (12)	0.0098 (13)
O4	0.0348 (11)	0.0834 (14)	0.0379 (11)	-0.0033 (10)	0.0092 (10)	0.0114 (10)
N1	0.0306 (11)	0.0451 (10)	0.0327 (12)	-0.0052 (9)	0.0136 (11)	-0.0034 (9)
C1	0.0366 (15)	0.0576 (16)	0.0390 (15)	-0.0023 (12)	0.0183 (14)	0.0082 (12)
C2	0.0505 (19)	0.0595 (17)	0.0534 (19)	0.0039 (14)	0.0223 (18)	0.0035 (14)
C3	0.072 (3)	0.0648 (19)	0.065 (2)	-0.0187 (18)	0.032 (2)	-0.0098 (17)
C4	0.051 (2)	0.097 (3)	0.056 (2)	-0.0244 (18)	0.019 (2)	-0.0037 (19)
C5	0.0363 (16)	0.084 (2)	0.067 (2)	-0.0003 (15)	0.0225 (18)	0.0078 (17)
C6	0.0403 (17)	0.0656 (18)	0.064 (2)	-0.0032 (14)	0.0267 (17)	-0.0010 (15)
C7	0.0326 (15)	0.0515 (13)	0.0454 (16)	-0.0013 (11)	0.0195 (15)	0.0035 (12)
C8	0.0307 (14)	0.0386 (12)	0.0393 (15)	-0.0014 (10)	0.0154 (13)	-0.0002 (10)

C9	0.0312 (14)	0.0415 (12)	0.0368 (15)	-0.0044 (10)	0.0142 (13)	-0.0031 (10)
C10	0.0331 (14)	0.0396 (12)	0.0380 (15)	-0.0005 (10)	0.0160 (13)	-0.0026 (11)
C11	0.0418 (16)	0.0620 (16)	0.0353 (15)	-0.0069 (13)	0.0151 (14)	-0.0027 (12)
C12	0.0500 (18)	0.0751 (18)	0.0461 (18)	-0.0146 (15)	0.0298 (17)	-0.0089 (14)
C13	0.0359 (16)	0.0649 (17)	0.059 (2)	-0.0088 (13)	0.0263 (16)	-0.0090 (15)
C14	0.0309 (14)	0.0548 (14)	0.0408 (16)	-0.0034 (11)	0.0132 (14)	-0.0052 (12)
C15	0.0363 (14)	0.0374 (12)	0.0349 (14)	-0.0072 (10)	0.0166 (13)	-0.0060 (10)
C16	0.0366 (15)	0.0502 (14)	0.0431 (16)	-0.0009 (11)	0.0150 (15)	0.0036 (12)
C17	0.0382 (17)	0.086 (2)	0.0483 (19)	-0.0086 (15)	0.0052 (17)	0.0149 (17)
C18	0.060 (2)	0.103 (3)	0.045 (2)	-0.0090 (19)	0.0121 (19)	0.0156 (17)
C19	0.0389 (15)	0.0421 (12)	0.0274 (13)	0.0013 (11)	0.0095 (13)	0.0045 (10)
C20	0.0489 (18)	0.0488 (15)	0.0510 (18)	-0.0011 (12)	0.0186 (17)	-0.0036 (13)
C21	0.068 (2)	0.0468 (15)	0.057 (2)	0.0111 (15)	0.021 (2)	0.0027 (14)
C22	0.079 (3)	0.0418 (15)	0.0470 (19)	-0.0043 (15)	0.014 (2)	0.0021 (13)
C23	0.051 (2)	0.0597 (18)	0.067 (2)	-0.0175 (15)	0.007 (2)	-0.0056 (16)
C24	0.0490 (19)	0.0538 (16)	0.062 (2)	-0.0009 (13)	0.0170 (18)	-0.0037 (15)

Geometric parameters (Å, °)

S1—O1	1.4195 (18)	C10—C15	1.393 (4)
S1—O2	1.4318 (19)	C10—C11	1.398 (4)
S1—N1	1.683 (2)	C11—C12	1.388 (4)
S1—C19	1.755 (2)	C11—H11	0.9300
S2—C1	1.773 (3)	C12—C13	1.390 (4)
S2—C7	1.830 (3)	C12—H12	0.9300
O3—C16	1.201 (3)	C13—C14	1.386 (4)
O4—C16	1.338 (3)	C13—H13	0.9300
O4—C17	1.437 (3)	C14—C15	1.398 (4)
N1—C8	1.394 (3)	C14—H14	0.9300
N1—C15	1.427 (3)	C17—C18	1.533 (3)
C1—C6	1.381 (4)	C17—H17A	0.9700
C1—C2	1.381 (4)	C17—H17B	0.9700
C2—C3	1.383 (4)	C18—H18A	0.9600
C2—H2	0.9300	C18—H18B	0.9600
C3—C4	1.381 (3)	C18—H18C	0.9600
C3—H3	0.9300	C19—C20	1.383 (2)
C4—C5	1.383 (3)	C19—C24	1.385 (2)
C4—H4	0.9300	C20—C21	1.385 (2)
C5—C6	1.385 (3)	C20—H20	0.9300
C5—H5	0.9300	C21—C22	1.383 (4)
C6—H6	0.9300	C21—H21	0.9300
C7—C8	1.498 (3)	C22—C23	1.382 (3)
C7—H7A	0.9700	C22—H22	0.9300
C7—H7B	0.9700	C23—C24	1.387 (3)
C8—C9	1.386 (3)	C23—H23	0.9300
C9—C10	1.453 (3)	C24—H24	0.9300
C9—C16	1.467 (4)		
O1—S1—O2	119.78 (11)	C10—C11—H11	120.6
O1—S1—N1	106.73 (11)	C11—C12—C13	121.1 (3)

supplementary materials

O2—S1—N1	106.35 (11)	C11—C12—H12	119.4
O1—S1—C19	109.19 (11)	C13—C12—H12	119.4
O2—S1—C19	108.93 (11)	C14—C13—C12	121.4 (3)
N1—S1—C19	104.79 (10)	C14—C13—H13	119.3
C1—S2—C7	105.10 (12)	C12—C13—H13	119.3
C16—O4—C17	116.6 (2)	C13—C14—C15	116.8 (3)
C8—N1—C15	109.02 (19)	C13—C14—H14	121.6
C8—N1—S1	127.98 (17)	C15—C14—H14	121.6
C15—N1—S1	122.05 (17)	C10—C15—C14	122.9 (2)
C6—C1—C2	119.4 (3)	C10—C15—N1	107.4 (2)
C6—C1—S2	124.5 (2)	C14—C15—N1	129.8 (2)
C2—C1—S2	115.7 (2)	O3—C16—O4	122.9 (3)
C1—C2—C3	120.1 (3)	O3—C16—C9	126.2 (3)
C1—C2—H2	119.9	O4—C16—C9	110.9 (2)
C3—C2—H2	119.9	O4—C17—C18	106.6 (2)
C4—C3—C2	120.9 (3)	O4—C17—H17A	110.4
C4—C3—H3	119.6	C18—C17—H17A	110.4
C2—C3—H3	119.6	O4—C17—H17B	110.4
C3—C4—C5	118.7 (3)	C18—C17—H17B	110.4
C3—C4—H4	120.7	H17A—C17—H17B	108.6
C5—C4—H4	120.7	C17—C18—H18A	109.5
C4—C5—C6	120.8 (3)	C17—C18—H18B	109.5
C4—C5—H5	119.6	H18A—C18—H18B	109.5
C6—C5—H5	119.6	C17—C18—H18C	109.5
C1—C6—C5	120.1 (3)	H18A—C18—H18C	109.5
C1—C6—H6	120.0	H18B—C18—H18C	109.5
C5—C6—H6	120.0	C20—C19—C24	122.2 (2)
C8—C7—S2	110.86 (17)	C20—C19—S1	118.70 (18)
C8—C7—H7A	109.5	C24—C19—S1	119.09 (17)
S2—C7—H7A	109.5	C19—C20—C21	119.2 (3)
C8—C7—H7B	109.5	C19—C20—H20	120.4
S2—C7—H7B	109.5	C21—C20—H20	120.4
H7A—C7—H7B	108.1	C22—C21—C20	119.3 (3)
C9—C8—N1	108.2 (2)	C22—C21—H21	120.3
C9—C8—C7	127.3 (2)	C20—C21—H21	120.3
N1—C8—C7	124.4 (2)	C23—C22—C21	120.8 (3)
C8—C9—C10	108.1 (2)	C23—C22—H22	119.6
C8—C9—C16	124.4 (2)	C21—C22—H22	119.6
C10—C9—C16	127.5 (2)	C22—C23—C24	120.6 (3)
C15—C10—C11	119.0 (2)	C22—C23—H23	119.7
C15—C10—C9	107.3 (2)	C24—C23—H23	119.7
C11—C10—C9	133.8 (3)	C19—C24—C23	117.8 (2)
C12—C11—C10	118.8 (3)	C19—C24—H24	121.1
C12—C11—H11	120.6	C23—C24—H24	121.1
O1—S1—N1—C8	-20.4 (2)	C11—C12—C13—C14	0.2 (4)
O2—S1—N1—C8	-149.37 (19)	C12—C13—C14—C15	-0.5 (4)
C19—S1—N1—C8	95.3 (2)	C11—C10—C15—C14	-1.9 (3)
O1—S1—N1—C15	171.99 (17)	C9—C10—C15—C14	178.2 (2)
O2—S1—N1—C15	43.0 (2)	C11—C10—C15—N1	178.6 (2)

C19—S1—N1—C15	-72.3 (2)	C9—C10—C15—N1	-1.3 (2)
C7—S2—C1—C6	34.8 (3)	C13—C14—C15—C10	1.4 (4)
C7—S2—C1—C2	-151.5 (2)	C13—C14—C15—N1	-179.3 (2)
C6—C1—C2—C3	0.0 (4)	C8—N1—C15—C10	2.0 (2)
S2—C1—C2—C3	-174.0 (2)	S1—N1—C15—C10	171.65 (15)
C1—C2—C3—C4	-0.9 (5)	C8—N1—C15—C14	-177.5 (2)
C2—C3—C4—C5	0.3 (5)	S1—N1—C15—C14	-7.8 (3)
C3—C4—C5—C6	1.3 (5)	C17—O4—C16—O3	-2.5 (4)
C2—C1—C6—C5	1.5 (4)	C17—O4—C16—C9	177.3 (2)
S2—C1—C6—C5	175.0 (2)	C8—C9—C16—O3	-10.4 (4)
C4—C5—C6—C1	-2.2 (5)	C10—C9—C16—O3	169.3 (3)
C1—S2—C7—C8	119.3 (2)	C8—C9—C16—O4	169.8 (2)
C15—N1—C8—C9	-1.9 (2)	C10—C9—C16—O4	-10.5 (3)
S1—N1—C8—C9	-170.77 (17)	C16—O4—C17—C18	166.6 (3)
C15—N1—C8—C7	-177.4 (2)	O1—S1—C19—C20	48.4 (2)
S1—N1—C8—C7	13.7 (3)	O2—S1—C19—C20	-179.2 (2)
S2—C7—C8—C9	-103.3 (3)	N1—S1—C19—C20	-65.7 (2)
S2—C7—C8—N1	71.4 (3)	O1—S1—C19—C24	-129.6 (2)
N1—C8—C9—C10	1.1 (3)	O2—S1—C19—C24	2.9 (3)
C7—C8—C9—C10	176.5 (2)	N1—S1—C19—C24	116.4 (2)
N1—C8—C9—C16	-179.2 (2)	C24—C19—C20—C21	2.1 (4)
C7—C8—C9—C16	-3.8 (4)	S1—C19—C20—C21	-175.8 (2)
C8—C9—C10—C15	0.2 (3)	C19—C20—C21—C22	-0.6 (4)
C16—C9—C10—C15	-179.6 (2)	C20—C21—C22—C23	-0.6 (5)
C8—C9—C10—C11	-179.7 (3)	C21—C22—C23—C24	0.3 (5)
C16—C9—C10—C11	0.6 (4)	C20—C19—C24—C23	-2.3 (5)
C15—C10—C11—C12	1.5 (4)	S1—C19—C24—C23	175.6 (2)
C9—C10—C11—C12	-178.6 (3)	C22—C23—C24—C19	1.0 (5)
C10—C11—C12—C13	-0.7 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C7—H7A...O1	0.97	2.38	2.813 (4)	107
C7—H7B...O3	0.97	2.26	2.977 (4)	130
C11—H11...O4	0.93	2.42	2.932 (3)	115
C14—H14...O2	0.93	2.41	2.966 (3)	119
C24—H24...O2	0.93	2.52	2.898 (3)	105

Fig. 1

