

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5,5'-Bis[(trimethylsilyl)methyl]-2,2'-bipyridine

Murray S. Davies,* Christopher R. K. Glasson and George V. Meehan

School of Pharmacy and Molecular Sciences, James Cook University, Townsville, QLD 4811, Australia

Correspondence e-mail: murray.davies@jcu.edu.au


Received 17 May 2007; accepted 21 October 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.064; wR factor = 0.192; data-to-parameter ratio = 16.7.

The molecule of the title compound, $\text{C}_{18}\text{H}_{28}\text{N}_2\text{Si}_2$, occupies a special position on an inversion centre. The $\text{Si}-\text{CH}_2-\text{C}(\text{ipso})$ plane is approximately orthogonal to the plane of the pyridine rings, the corresponding dihedral angle being 82.0 (2)°.

Related literature

For related chemistry, see: Fraser *et al.* (1997); Hochwimmer *et al.* (1998); Perkins *et al.* (2006); Schubert *et al.* (1998). For recently reported similar structures, see: Khan *et al.* (2004); Lindoy *et al.* (2004). For related literature, see: Lindoy *et al.* (2004).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{28}\text{N}_2\text{Si}_2$
 $M_r = 328.60$
 Triclinic, $P\bar{1}$
 $a = 6.279$ (3) Å
 $b = 6.575$ (3) Å

$c = 14.030$ (6) Å
 $\alpha = 76.599$ (7)°
 $\beta = 88.415$ (7)°
 $\gamma = 64.859$ (6)°
 $V = 508.4$ (4) Å³

$Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.17$ mm⁻¹

$T = 293$ (2) K
 $0.25 \times 0.15 \times 0.10$ mm

Data collection

Bruker SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.973$, $T_{\max} = 0.974$

4192 measured reflections
 2057 independent reflections
 1294 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$
 $wR(F^2) = 0.192$
 $S = 1.04$
 2057 reflections
 123 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.36$ e Å⁻³
 $\Delta\rho_{\min} = -0.16$ e Å⁻³

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT and XPREP (Bruker, 2001); program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and ORTEPIII (Burnett & Johnson, 1996); software used to prepare material for publication: WinGX (Farrugia, 1999).

The authors acknowledge financial support by the Australian Research Council.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: YA2054).

References

- Bruker (2001). SMART, SAINT and XPREP. Bruker Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Burnett, M. N. & Johnson, C. K. (1996). ORTEPIII. Report ORNL-6895. Oak Ridge National Laboratory, Oak Ridge, Tennessee, U. S.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Fraser, C. L., Anastasi, N. R. & Lamba, J. J. S. (1997). *J. Org. Chem.* **62**, 9314–9317.
 Hochwimmer, G., Nuyken, O. & Schubert, U. S. (1998). *Macromol. Rapid Commun.* **19**, 309–313.
 Khan, M. S., Ahrens, B., Male, L. & Raithby, P. R. (2004). *Acta Cryst.* **E60**, o915–o916.
 Lindoy, L. F., McMurtrie, J. C. & Price, J. R. (2004). *Acta Cryst.* **E60**, o886–o888.
 Perkins, D. F., Lindoy, L. F., McAuley, A., Meehan, G. V. & Turner, P. (2006). *Proc. Natl Acad. Sci. USA*, **103**, 532–537.
 Schubert, U. S., Eschbaumer, C. & Hochwimmer, G. (1998). *Tetrahedron Lett.* **39**, 8643–8644.
 Sheldrick, G. M. (1996). SADABS: University of Göttingen, Germany.
 Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.

supplementary materials

Acta Cryst. (2008). E64, o364 [doi:10.1107/S1600536807052154]

5,5'-Bis[(trimethylsilyl)methyl]-2,2'-bipyridine

M. S. Davies, C. R. K. Glasson and G. V. Meehan

Comment

As a part of our work on cryptates derived from 5,5'-disubstituted-2,2'-bipyridines (Perkins *et al.*, 2006), we have studied methyl functionalization reactions of 5,5'-dimethyl-2,2'-bipyridine as a model for similar chemistry proposed for its more complex analogue, 5,5''-dimethyl-2,2':5',5'':2'',2''''-quaterpyridine. In contrast to the previous report by Schubert *et al.* (1998), we have been able to promote bis-lithiation of 5,5'-dimethyl-2,2'-bipyridine with lithium diisopropylamide (LDA) in THF by use of a coordinating co-solvent, hexamethylphosphoramide (HMPA). Subsequent bis-silylation with trimethylsilyl chloride afforded (I) in good yield.

The molecule of the title compound (Fig.1) occupies a special position in the inversion centre. The SiMe₃ groups are *trans* disposed relative to the plane of the two pyridyl rings giving the molecule a zigzag shape (Fig. 2). The dihedral angle between the plane of the bipyridyl rings, and that of trimethylsilylmethyl substituent, as defined by Si1—C6—C4, is 82.0 (2)°.

The molecules in crystal show stacking arrangement with methylenetrimethylsilyl groups of the adjacent molecules oriented in the same direction (Fig. 2).

Experimental

A solution of LDA, prepared from *n*-BuLi (1.9 M, 1.42 ml, 2.7 mmol), and dry diisopropylamine (0.42 ml, 3.0 mmol) in dry THF (8 ml) was cooled to -78° C and a solution of 5,5'-dimethyl-2,2'-bipyridine (100 mg, 0.54 mmol) and dry HMPA (1.13 ml, 6.5 mmol) in dry THF (5 ml) was added dropwise, resulting in a deep red/brown opaque reaction mixture. This was stirred for 2 h, then trimethylsilyl chloride (217 mg, 2.0 mmol) was added, and the stirring was continued for 0.5 h more at -78° C. The resulting transparent red solution was quenched with 2 ml of absolute ethanol. Fortuitously, this solution precipitated crystals of (I) suitable for X-ray structure determination when its volume was reduced by rotary evaporation. To the remaining material, saturated NaHCO₃ (10 ml) was added and the product was extracted with ethyl acetate (3 × 40 ml). The combined organic fractions were dried over anhydrous Na₂SO₄ and the solvent removed under vacuum. The resulting solid was purified by chromatography on deactivated silica gel, affording (I) as a greasy white solid (142 mg, 80%). δH (300 MHz; CDCl₃) 0.02 (18H, s, SiMe₃) 2.12 (4H, s, CH₂), 7.45 (2H, dd, J = 7.8, 1.8 Hz, H-4,4'), 8.22 (2H, d, J = 7.8 Hz, H-3,3'), 8.34 (2H, d, 1.8 Hz, H-6,6'); δC (75 MHz; CDCl₃) 2.06 (SiMe₃), 23.92 (CH₂), 120.16, 136.11, 136.18, 148.22, 152.21.

Refinement

All aromatic and methylene H atoms were located in the difference map and refined with isotropic thermal parameters [C—H 0.94 (2) – 1.03 (3) Å]. Methyl H atoms were positioned geometrically and refined in a riding model approximation with C—H bond distances of 0.96 Å and $U_{iso}(H) = 1.5 U_{eq}$ of the parent C atom.

Figures




Fig. 1. ORTEP (Burnett and Johnson (1996), Farrugia (1997)) drawing of (I) with displacement ellipsoids shown at 50% probability level. Only the non-hydrogen atoms in the asymmetric unit are labelled; unlabelled atoms are derived from the corresponding labelled atoms by means of the $(1 - x, 3 - y, -z)$ transformation.




Fig. 2. Crystal packing diagram of (I) viewed down the a axis of the crystal.

5,5'-Bis[(trimethylsilyl)methyl]-2,2'-bipyridine

Crystal data

$C_{18}H_{28}N_2Si_2$

$M_r = 328.60$

Triclinic, PT

Hall symbol: $-P 1$

$a = 6.279$ (3) Å

$b = 6.575$ (3) Å

$c = 14.030$ (6) Å

$\alpha = 76.599$ (7)°

$\beta = 88.415$ (7)°

$\gamma = 64.859$ (6)°

$V = 508.4$ (4) Å³

$Z = 1$

$F_{000} = 178$

$D_x = 1.073$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71069$ Å

Cell parameters from 1147 reflections

$\theta = 0.9$ – 26.4 °

$\mu = 0.17$ mm⁻¹

$T = 293$ (2) K

Prism, colourless

$0.25 \times 0.15 \times 0.10$ mm

Data collection

Bruker SMART 1000 CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ (2) K

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.973$, $T_{\max} = 0.974$

4192 measured reflections

2057 independent reflections

1294 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.042$

$\theta_{\max} = 26.4$ °

$\theta_{\min} = 3.0$ °

$h = -7 \rightarrow 7$

$k = -8 \rightarrow 8$

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.064$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of

	independent and constrained refinement
$wR(F^2) = 0.192$	$w = 1/[\sigma^2(F_o^2) + (0.093P)^2]$
	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.04$	$(\Delta/\sigma)_{\max} = 0.007$
2057 reflections	$\Delta\rho_{\max} = 0.36 \text{ e } \text{\AA}^{-3}$
123 parameters	$\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. An empirical absorption correction determined with *SADABS* (Sheldrick, 1996) was applied to the data. The data integration and reduction were undertaken with *SAINTE* and *XPREP* (Bruker, 2001). The data reduction included the application of Lorentz and polarization corrections. The reflection data were merged including Friedel opposites. The structure was solved in the space group P-1 by direct methods with *SHELXS97* (Sheldrick, 1997) within the WinG-X (Farrugia, 1999) interface and extended and refined with *SHELXL97* (Sheldrick 1997). Anisotropic thermal parameters were refined for the non-hydrogen atoms. All aromatic and methylene H atoms were located and refined with isotropic thermal parameters. Methyl H atoms were constrained as riding atoms, fixed to the parent C atom with a distance of 0.96 Å. U_{iso} values were set to 1.5 U_{eq} of the parent C atom.

Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Si1	0.78941 (12)	0.61211 (12)	0.82038 (5)	0.0664 (3)
C1	0.5188 (4)	1.3836 (4)	0.52943 (15)	0.0544 (5)
N1	0.7386 (3)	1.2150 (4)	0.53910 (15)	0.0681 (6)
C4	0.6028 (4)	0.9417 (4)	0.63513 (16)	0.0594 (6)
C2	0.3355 (4)	1.3408 (5)	0.57193 (18)	0.0646 (7)
C5	0.7751 (5)	1.0028 (5)	0.59098 (19)	0.0686 (7)
C6	0.6608 (5)	0.6974 (5)	0.69012 (18)	0.0660 (7)
C3	0.3787 (5)	1.1207 (5)	0.62426 (18)	0.0673 (7)
C9	0.5840 (5)	0.8130 (5)	0.8897 (2)	0.0951 (9)
H9A	0.5359	0.9699	0.8525	0.143*
H9B	0.6620	0.7910	0.9517	0.143*
H9C	0.4478	0.7823	0.9009	0.143*
C7	0.8328 (6)	0.3086 (5)	0.8742 (2)	0.1086 (11)
H7B	0.8969	0.2606	0.9412	0.163*
H7C	0.9400	0.2082	0.8371	0.163*
H7A	0.6838	0.3008	0.8719	0.163*

supplementary materials

C8	1.0830 (5)	0.6243 (6)	0.8206 (2)	0.1070 (11)
H8A	1.0637	0.7764	0.7865	0.160*
H8B	1.1908	0.5123	0.7882	0.160*
H8C	1.1446	0.5904	0.8871	0.160*
H6A	0.773 (4)	0.589 (4)	0.6581 (18)	0.074 (8)*
H6B	0.513 (5)	0.676 (4)	0.6947 (18)	0.087 (8)*
H5	0.923 (4)	0.883 (4)	0.5873 (17)	0.075 (7)*
H3	0.243 (5)	1.082 (4)	0.6515 (19)	0.093 (8)*
H2	0.179 (4)	1.470 (4)	0.5684 (16)	0.067 (7)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Si1	0.0679 (5)	0.0709 (5)	0.0656 (5)	-0.0393 (4)	0.0016 (3)	-0.0070 (3)
C1	0.0476 (13)	0.0684 (14)	0.0500 (12)	-0.0274 (12)	0.0054 (10)	-0.0148 (10)
N1	0.0537 (12)	0.0677 (13)	0.0784 (14)	-0.0267 (10)	0.0164 (10)	-0.0096 (11)
C4	0.0635 (15)	0.0713 (15)	0.0509 (12)	-0.0374 (13)	0.0049 (11)	-0.0123 (11)
C2	0.0457 (13)	0.0728 (17)	0.0686 (15)	-0.0243 (13)	0.0056 (11)	-0.0077 (13)
C5	0.0546 (15)	0.0663 (16)	0.0765 (17)	-0.0221 (13)	0.0141 (12)	-0.0104 (14)
C6	0.0720 (18)	0.0705 (16)	0.0665 (16)	-0.0400 (15)	0.0085 (13)	-0.0187 (13)
C3	0.0567 (15)	0.0804 (18)	0.0696 (16)	-0.0381 (14)	0.0088 (12)	-0.0107 (13)
C9	0.108 (2)	0.116 (2)	0.0779 (19)	-0.060 (2)	0.0187 (17)	-0.0319 (18)
C7	0.127 (3)	0.084 (2)	0.110 (2)	-0.052 (2)	-0.006 (2)	-0.0014 (18)
C8	0.080 (2)	0.136 (3)	0.113 (3)	-0.061 (2)	-0.0108 (18)	-0.015 (2)

Geometric parameters (\AA , $^\circ$)

Si1—C9	1.853 (3)	C5—H5	0.94 (2)
Si1—C7	1.864 (3)	C6—H6A	0.95 (2)
Si1—C6	1.878 (3)	C6—H6B	1.00 (3)
Si1—C8	1.879 (3)	C3—H3	1.03 (3)
C1—N1	1.339 (3)	C9—H9A	0.9600
C1—C2	1.385 (3)	C9—H9B	0.9600
C1—C1 ⁱ	1.483 (4)	C9—H9C	0.9600
N1—C5	1.340 (3)	C7—H7B	0.9600
C4—C3	1.382 (4)	C7—H7C	0.9600
C4—C5	1.390 (3)	C7—H7A	0.9600
C4—C6	1.501 (4)	C8—H8A	0.9600
C2—C3	1.376 (3)	C8—H8B	0.9600
C2—H2	0.98 (2)	C8—H8C	0.9600
C9—Si1—C7	111.14 (16)	Si1—C6—H6B	105.3 (15)
C9—Si1—C6	109.21 (14)	H6A—C6—H6B	110 (2)
C7—Si1—C6	107.55 (14)	C2—C3—C4	120.9 (2)
C9—Si1—C8	110.33 (15)	C2—C3—H3	120.9 (15)
C7—Si1—C8	109.30 (16)	C4—C3—H3	118.1 (15)
C6—Si1—C8	109.25 (13)	Si1—C9—H9A	109.5
N1—C1—C2	121.3 (2)	Si1—C9—H9B	109.5
N1—C1—C1 ⁱ	116.9 (2)	H9A—C9—H9B	109.5

C2—C1—C1 ⁱ	121.8 (2)	Si1—C9—H9C	109.5
C1—N1—C5	117.7 (2)	H9A—C9—H9C	109.5
C3—C4—C5	115.2 (2)	H9B—C9—H9C	109.5
C3—C4—C6	123.4 (2)	Si1—C7—H7B	109.5
C5—C4—C6	121.4 (2)	Si1—C7—H7C	109.5
C3—C2—C1	119.6 (2)	H7B—C7—H7C	109.5
C3—C2—H2	120.8 (13)	Si1—C7—H7A	109.5
C1—C2—H2	119.5 (13)	H7B—C7—H7A	109.5
N1—C5—C4	125.4 (2)	H7C—C7—H7A	109.5
N1—C5—H5	115.7 (15)	Si1—C8—H8A	109.5
C4—C5—H5	118.0 (15)	Si1—C8—H8B	109.5
C4—C6—Si1	115.70 (17)	H8A—C8—H8B	109.5
C4—C6—H6A	111.0 (14)	Si1—C8—H8C	109.5
Si1—C6—H6A	105.7 (14)	H8A—C8—H8C	109.5
C4—C6—H6B	108.8 (15)	H8B—C8—H8C	109.5

Symmetry codes: (i) $-x+1, -y+3, -z+1$.

Fig. 1




Fig. 2

