

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Diaquabis(4-bromobenzoato- κ^2O,O')-zinc(II)Tuncer Hökelek,^a Nagihan Çaylak^b and Hacali Necefoğlu^{c*}^aHacettepe University, Department of Physics, 06800 Beytepe, Ankara, Turkey,^bSakarya University, Faculty of Arts and Science, Department of Physics, 54187 Esentepe, Adapazarı, Turkey, and ^cKafkas University, Department of Chemistry, 63100 Kars, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 28 January 2008; accepted 4 February 2008

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.011$ Å; R factor = 0.064; wR factor = 0.167; data-to-parameter ratio = 14.3.

The monomeric title Zn^{II} complex, $[\text{Zn}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{H}_2\text{O})_2]$, contains two 4-bromobenzoate (BB) ligands and two coordinated water molecules around a Zn^{II} atom on a twofold rotation axis. The BB ions act as bidentate ligands, with two very dissimilar coordination distances. The sixfold coordination around the Zn^{II} may be described as highly distorted octahedral, with the two aqua ligands arranged *cis*. Hydrogen bonding involving the carboxylate O atoms has an effect on the delocalization in the carboxylate groups. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into chains parallel to the c axis and stacked along the b axis.

Related literature

For general background, see: Antolini *et al.* (1982); Chen & Chen (2002); Amiraslanov *et al.* (1979); Hauptmann *et al.* (2000); Shnulin *et al.* (1981); Antsyshkina *et al.* (1980); Adiwidjaja *et al.* (1978); Catterick *et al.* (1974). For related literature, see: Guseinov *et al.* (1984); Clegg *et al.* (1986*a,b*, 1987); Capilla & Aranda (1979); van Niekerk *et al.* (1953); Usabaliev *et al.* (1992); Musaev *et al.* (1983); Nadzhafov *et al.* (1981); Day & Selbin (1969); Amiraslanov *et al.* (1980); Necefoğlu *et al.* (2002); Hökelek *et al.* (2008, 2007); Hökelek & Necefoğlu (1996, 2001, 2007); Greenaway *et al.* (1984).



Experimental

Crystal data

 $[\text{Zn}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{H}_2\text{O})_2]$ $M_r = 501.43$ Monoclinic, $C2/c$ $a = 26.9067$ (3) Å $b = 5.0704$ (4) Å $c = 12.0371$ (5) Å $\beta = 104.95$ (2)° $V = 1586.6$ (2) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 6.61$ mm⁻¹ $T = 294$ (2) K $0.25 \times 0.20 \times 0.15$ mm

Data collection

Enraf–Nonius TurboCAD-4

diffractometer

Absorption correction: ψ scan(North *et al.*, 1968) $T_{\text{min}} = 0.214$, $T_{\text{max}} = 0.370$

1648 measured reflections

1613 independent reflections

1133 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.031$

3 standard reflections

frequency: 120 min

intensity decay: 1%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.063$ $wR(F^2) = 0.166$ $S = 1.04$

1613 reflections

113 parameters

4 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 1.42$ e Å⁻³ $\Delta\rho_{\text{min}} = -1.83$ e Å⁻³

Table 1

Selected bond lengths (Å).

Zn—O1	2.010 (5)	Zn—O3	1.993 (5)
Zn—O2	2.468 (5)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O3}-\text{H31}\cdots\text{O2}^{\text{i}}$	0.97 (7)	1.82 (6)	2.746 (7)	157 (9)
$\text{O3}-\text{H32}\cdots\text{O1}^{\text{ii}}$	0.95 (8)	1.86 (8)	2.765 (7)	160 (9)

Symmetry codes: (i) $-x + 2, -y, -z + 1$; (ii) $-x + 2, y - 1, -z + \frac{1}{2}$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors acknowledge the purchase of the CAD-4 diffractometer under grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BG2161).

References

- Adiwidjaja, G., Rossmanith, E. & Küppers, H. (1978). *Acta Cryst.* **B34**, 3079–3083.
- Amiraslanov, I. R., Mamedov, Kh. S., Movsumov, E. M., Musaev, F. N. & Nadzhafov, G. N. (1979). *Zh. Strukt. Khim.* **20**, 1075–1080.
- Amiraslanov, I. R., Nadzhafov, G. N., Usabaliev, B. T., Musaev, A. A., Movsumov, E. M. & Mamedov, Kh. S. (1980). *Zh. Strukt. Khim.* **21**, 140–145.

- Antolini, L., Battaglia, L. P., Corradi, A. B., Marcotrigiano, G., Menabue, L., Pellacani, G. C. & Saladini, M. (1982). *Inorg. Chem.* **21**, 1391–1395.
- Antsyshkina, A. S., Chiragov, F. M. & Poray-Koshits, M. A. (1980). *Koord. Khim.* **15**, 1098–1103.
- Capilla, A. V. & Aranda, R. A. (1979). *Cryst. Struct. Commun.* **8**, 795–798.
- Catterick, J., Hursthouse, M. B., New, D. B. & Thornton, P. (1974). *J. Chem. Soc. Chem. Commun.* pp. 843–844.
- Chen, H. J. & Chen, X. M. (2002). *Inorg. Chim. Acta*, **329**, 13–21.
- Clegg, W., Little, I. R. & Straughan, B. P. (1986a). *Acta Cryst.* **C42**, 919–920.
- Clegg, W., Little, I. R. & Straughan, B. P. (1986b). *Acta Cryst.* **C42**, 1701–1703.
- Clegg, W., Little, I. R. & Straughan, B. P. (1987). *Acta Cryst.* **C43**, 456–457.
- Day, M. C. & Selbin, J. (1969). *Theoretical Inorganic Chemistry*, p. 109. New York: Van Nostrand Reinhold.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Greenaway, F. T., Pezeshk, A., Cordes, A. W., Noble, M. C. & Sorenson, J. R. J. (1984). *Inorg. Chim. Acta*, **93**, 67–71.
- Guseinov, G. A., Musaev, F. N., Usabaliev, B. T., Amiraslanov, I. R. & Mamedov, Kh. S. (1984). *Koord. Khim.* **10**, 117–122.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Hauptmann, R., Kondo, M. & Kitagawa, S. (2000). *Z. Kristallogr. New Cryst. Struct.* **215**, 169–172.
- Hökelek, T., Çaylak, N. & Necefoğlu, H. (2007). *Acta Cryst.* **E63**, m2561–m2562.
- Hökelek, T., Çaylak, N. & Necefoğlu, H. (2008). *Acta Cryst.* **E64**, m460–m461.
- Hökelek, T. & Necefoğlu, H. (1996). *Acta Cryst.* **C52**, 1128–1131.
- Hökelek, T. & Necefoğlu, H. (2001). *Anal. Sci.* **17**, 1241–1242.
- Hökelek, T. & Necefoğlu, H. (2007). *Acta Cryst.* **E63**, m821–m823.
- Musaev, F. N., Nadzhafov, G. N. & Mamedov, Kh. S. (1983). *Koord. Khim.* **12**, 37–46.
- Nadzhafov, G. N., Usabaliev, B. T., Amiraslanov, I. R., Movsumov, E. M. & Mamedov, Kh. S. (1981). *Koord. Khim.* **7**, 770–775.
- Necefoğlu, H., Hökelek, T., Ersanlı, C. C. & Erdönmez, A. (2002). *Acta Cryst.* **E58**, m758–m761.
- Niekerk, J. N. van, Schoening, F. R. L. & Talbot, J. H. (1953). *Acta Cryst.* **6**, 720–723.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Shnulin, A. N., Nadzhafov, G. N., Amiraslanov, I. R., Usabaliev, B. T. & Mamedov, Kh. S. (1981). *Koord. Khim.* **7**, 1409–1416.
- Usabaliev, B. T., Guliev, F. I., Musaev, F. N., Ganbarov, D. M., Ashurova, S. A. & Movsumov, E. M. (1992). *Zh. Strukt. Khim.* **33**, 203–207.

supplementary materials

Acta Cryst. (2008). E64, m458-m459 [doi:10.1107/S1600536808003759]

Diaquabis(4-bromobenzoato- κ^2O,O')zinc(II)

T. Hökelek, N. Çaylak and H. Necefoglu

Comment

Transition metal complexes with biochemical molecules show interesting physical and/or chemical properties, through which they may find applications in biological systems (Antolini *et al.*, 1982). Some benzoic acid derivatives, such as 4-aminobenzoic acid, have been extensively reported in coordination chemistry, as bifunctional organic ligands, due to the varieties of their coordination modes (Chen & Chen, 2002, Amiraslanov *et al.*, 1979; Hauptmann *et al.*, 2000).

The structure-function-coordination relationships of the arylcarboxylate ion in Zn^{II} complexes of benzoic acid derivatives may also change depending on the nature and position of the substituted groups on the benzene ring, the nature of the additional ligand molecule or solvent, and the pH and temperature of synthesis, as in Co^{II} complexes (Shnulin *et al.*, 1981; Antsyshkina *et al.*, 1980; Adiwidjaja *et al.*, 1978). When pyridine and its derivatives are used instead of water molecules, the structure is completely different (Catterick *et al.*, 1974).

The solid-state structures of anhydrous zinc(II) carboxylates include one-dimensional (Guseinov *et al.*, 1984; Clegg *et al.*, 1986a), two-dimensional (Clegg *et al.*, 1986b, 1987) and three-dimensional (Capilla & Aranda, 1979) polymeric motifs of different types, while discrete monomeric complexes with octahedral or tetrahedral coordination geometry are found if water or other donor molecules are coordinated to Zn (van Niekerk *et al.*, 1953; Usabaliev *et al.*, 1992). In hexaaquazinc(II) bis(4-hydroxybenzoate) dihydrate, $[Zn(H_2O)_6](4-HOC_6H_4COO)_2 \cdot 2H_2O$, [(II); Musaev *et al.*, 1983], which is isostructural with the corresponding Mg^{II} , Co^{II} , Ni^{II} and Mn^{II} compounds, the carboxylate ion lies outside the coordination sphere of the Zn atom, while $[Zn(4-HOC_6H_4COO)_2] \cdot 4C_5H_5N$ [(III); Nadzhafov *et al.*, 1981], forms a clathrate, consisting of $[Zn(4-HOC_6H_4COO)_2(C_5H_5N)_2]$ units with tetrahedral coordination geometry and free pyridine molecules.

The structure determination of the title compound, (I), a zinc complex with two bromobenzoate (BB) ligands and two water molecules, was undertaken in order to determine the ligand properties of (BB) and also to compare the results obtained with those reported previously.

In the monomeric title complex, $[Zn(C_7H_4O_2Br)_2(H_2O)_2]$, (I), the Zn atom lies on a twofold rotation axis and is surrounded by two 4-bromobenzoate (BB), acting as bidentate ligands, and two coordinated water molecules (Fig. 1).

The Zn coordination polyhedron is formed by four clear basal bonds and two close contacts of the symmetry related $O2$ and $O2^i$ atoms, [(i) $2 - x, y, 1/2 - z, Zn \cdots O2 = 2.468(5) \text{ \AA}$, in double dashed lines in Fig. 1] occupying apical positions and completing the six-coordination; this distance is greater than the sum of the corresponding ionic radii (2.14 \AA ; Day & Selbin, 1969), but similar $Zn \cdots O$ contacts have already been reported, viz.: 2.50 (1) \AA in (III), 2.494 (8) \AA in $[Zn(p-H_2NC_6H_4COO)_2] \cdot 1.5nH_2O$ [(IV); Amiraslanov *et al.*, 1980], 2.404 (2) \AA in $[Zn(C_6H_6N_2O_2)_2(C_7H_5O_3)_2]$ [(V), (Necefoglu *et al.*, 2002)] and 2.458 (3) \AA in $[Zn(C_7H_4O_2F)_2(C_6H_6N_2O_2)] \cdot H_2O$ [(VI); Hökelek *et al.*, 2008]. The sixfold coordination around Zn^{II} may thus be described as highly distorted octahedral (Table 1), with the two aqua ligands arranged *cis*.

supplementary materials

In the binuclear complex $[\text{Zn}_2(\text{C}_7\text{H}_5\text{O}_3)_4(\text{C}_{10}\text{H}_{14}\text{N}_2\text{O})_2(\text{H}_2\text{O})_2]$ [(VII); Hökelek & Necefoğlu, 1996], the average Zn—O bond length [1.953 (2) Å] is shorter than the corresponding value in (I) [2.157 (5) Å], but Zn is four coordinate. In complexes (V), $[\text{Zn}(\text{C}_7\text{H}_4\text{FO}_2)_2(\text{DNA})_2(\text{H}_2\text{O})_2]$ [(VIII); Hökelek *et al.*, 2007] and $[\text{Zn}(\text{C}_7\text{H}_5\text{O}_3)(\text{OH}_2)_3(\text{C}_6\text{H}_6\text{N}_2\text{O})].\text{C}_7\text{O}_3\text{H}_5$ [(IX); Hökelek & Necefoğlu, 2001], (where Zn atoms are five, six and five coordinates) the average Zn—O bond lengths are 2.107 (2) Å, 2.117 (2) Å and 2.047 (5) Å, respectively. In (I), the O1—Zn···O2 angle is 57.27 (18)°. The corresponding O—M···O (where *M* is a metal) angles are 58.79 (6)° in (V), 57.04 (10)° in (VI), 58.3 (3)° in (VII) and 55.2 (1)° in $[\text{Cu}(\text{Asp})_2(\text{py})_2]$ (where Asp is acetylsalicylate and py is pyridine) [(X); Greenaway *et al.*, 1984].

The near equality of C1—O1 [1.289 (8) Å] and C1—O2 [1.230 (9) Å] bonds in the carboxylate group indicates a delocalized bonding arrangement, rather than localized single and double bonds, as in (V) and $[\text{Mn}(\text{C}_9\text{H}_{10}\text{NO}_2)_2(\text{H}_2\text{O})_4].2\text{H}_2\text{O}$ [(XI); Hökelek & Necefoğlu, 2007]. This may be due to the intermolecular hydrogen bonds of the carboxyl O atoms (Table 2). The Zn atom is out of the least-squares plane of the carboxyl group (O1/C1/O2) by 0.055 (1) Å. The dihedral angle between the planar carboxyl group and the benzene ring (C2—C7) is 18.62 (44)°. The corresponding value is reported as 5.54 (43)° in (XI).

The molecules of (I) are linked by intermolecular O—H···O hydrogen bonds (Table 2), forming infinite chains along the [001] direction, which are in turn stacked along the *b* axis.

Experimental

The title compound, (I), was prepared by the reaction of ZnSO_4 (1.61 g, 10 mmol) in H_2O (100 ml) and *p*-bromobenzoate (4.00 g, 20 mmol) in H_2O (100 ml). The mixture was filtered and set aside to crystallize at ambient temperature for several days, giving colorless single crystals.

Refinement

H atoms of water molecules were located in difference syntheses and refined isotropically with restraints [O—H = 0.97 (7) and 0.95 (8) Å; $U_{\text{iso}}(\text{H}) = 0.09$ (4) and 0.09 (4) Å²]. The remaining H atoms were positioned geometrically with C—H = 0.93 Å, for aromatic H atoms and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. A drawing of the title molecule with the atom-numbering scheme. The displacement ellipsoids are drawn at the 50% probability level [symmetry code: (i) 2 - *x*, *y*, 1/2 - *z*].

Diaquabis(4-bromobenzoato- $\kappa^2\text{O},\text{O}'$)zinc(II)

Crystal data

$[\text{Zn}(\text{C}_7\text{H}_4\text{BrO}_2)_2(\text{H}_2\text{O})_2]$

$M_r = 501.43$

$F_{000} = 976$

$D_x = 2.099 \text{ Mg m}^{-3}$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 26.9067\ (3)\ \text{\AA}$

$b = 5.0704\ (4)\ \text{\AA}$

$c = 12.0371\ (5)\ \text{\AA}$

$\beta = 104.95\ (2)^\circ$

$V = 1586.6\ (2)\ \text{\AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 6.7\text{--}10.8^\circ$

$\mu = 6.61\ \text{mm}^{-1}$

$T = 294\ (2)\ \text{K}$

Block, colourless

$0.25 \times 0.20 \times 0.15\ \text{mm}$

Data collection

Enraf–Nonius TurboCAD-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 294\ (2)\ \text{K}$

non-profiled ω scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.214$, $T_{\max} = 0.370$

1648 measured reflections

1613 independent reflections

1133 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 26.3^\circ$

$\theta_{\min} = 3.1^\circ$

$h = -33 \rightarrow 0$

$k = -6 \rightarrow 0$

$l = -14 \rightarrow 15$

3 standard reflections

every 120 min

intensity decay: 1%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.063$

$wR(F^2) = 0.166$

$S = 1.04$

1613 reflections

113 parameters

4 restraints

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H atoms treated by a mixture of
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.1112P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 1.42\ \text{e}\ \text{\AA}^{-3}$

$\Delta\rho_{\min} = -1.83\ \text{e}\ \text{\AA}^{-3}$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

supplementary materials

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br	1.22102 (3)	1.19239 (18)	0.64660 (7)	0.0502 (4)
Zn	1.0000	0.0782 (2)	0.2500	0.0335 (4)
O1	1.0587 (2)	0.3361 (10)	0.2873 (4)	0.0342 (12)
O2	1.0366 (2)	0.2503 (10)	0.4458 (4)	0.0379 (12)
O3	0.9709 (2)	-0.1998 (10)	0.3320 (4)	0.0404 (13)
H31	0.960 (4)	-0.19 (2)	0.403 (5)	0.09 (4)*
H32	0.953 (4)	-0.343 (15)	0.289 (7)	0.09 (4)*
C1	1.0641 (3)	0.3712 (14)	0.3959 (6)	0.0314 (16)
C2	1.1029 (3)	0.5711 (14)	0.4540 (5)	0.0292 (16)
C3	1.0994 (3)	0.6756 (16)	0.5586 (6)	0.0361 (17)
H3	1.0732	0.6214	0.5909	0.043*
C4	1.1352 (3)	0.8615 (16)	0.6150 (6)	0.0404 (19)
H4	1.1329	0.9337	0.6845	0.048*
C5	1.1735 (3)	0.9357 (14)	0.5673 (6)	0.0316 (16)
C6	1.1776 (3)	0.8399 (17)	0.4632 (7)	0.0415 (19)
H6	1.2036	0.8983	0.4311	0.050*
C7	1.1418 (3)	0.6525 (17)	0.4067 (6)	0.0392 (18)
H7	1.1442	0.5826	0.3369	0.047*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br	0.0509 (6)	0.0389 (5)	0.0485 (5)	-0.0172 (4)	-0.0093 (4)	0.0011 (4)
Zn	0.0403 (7)	0.0166 (6)	0.0429 (7)	0.000	0.0097 (5)	0.000
O1	0.050 (3)	0.028 (3)	0.021 (2)	0.001 (2)	0.002 (2)	-0.006 (2)
O2	0.049 (3)	0.030 (3)	0.031 (3)	-0.014 (2)	0.005 (2)	0.000 (2)
O3	0.068 (4)	0.023 (3)	0.030 (3)	-0.009 (3)	0.012 (3)	-0.004 (2)
C1	0.038 (4)	0.026 (4)	0.027 (3)	0.005 (3)	0.000 (3)	-0.001 (3)
C2	0.039 (4)	0.019 (3)	0.023 (3)	0.001 (3)	-0.003 (3)	-0.003 (3)
C3	0.045 (4)	0.041 (4)	0.026 (3)	-0.012 (4)	0.014 (3)	-0.008 (3)
C4	0.054 (5)	0.035 (4)	0.030 (4)	-0.011 (4)	0.008 (3)	-0.009 (3)
C5	0.037 (4)	0.024 (3)	0.025 (3)	-0.006 (3)	-0.007 (3)	0.003 (3)
C6	0.043 (4)	0.043 (5)	0.040 (4)	-0.007 (4)	0.014 (4)	0.002 (4)
C7	0.044 (4)	0.050 (5)	0.023 (3)	-0.006 (4)	0.007 (3)	-0.009 (3)

Geometric parameters (\AA , $^\circ$)

Br—C5	1.901 (7)	C1—C2	1.494 (10)
Zn—O1	2.010 (5)	C2—C3	1.392 (10)

Zn—O1 ⁱ	2.010 (5)	C2—C7	1.376 (11)
Zn—O2	2.468 (5)	C3—H3	0.9300
Zn—O2 ⁱ	2.468 (5)	C4—C3	1.393 (10)
Zn—O3	1.993 (5)	C4—H4	0.9300
Zn—O3 ⁱ	1.993 (5)	C5—C4	1.354 (11)
O1—C1	1.289 (8)	C5—C6	1.375 (11)
O2—C1	1.230 (9)	C6—H6	0.9300
O3—H31	0.97 (7)	C7—C6	1.399 (11)
O3—H32	0.95 (8)	C7—H7	0.9300
O1—Zn—O1 ⁱ	98.8 (3)	O2—C1—Zn	70.5 (4)
O1—Zn—O2	57.27 (18)	O2—C1—O1	120.1 (7)
O1 ⁱ —Zn—O2	94.62 (19)	O2—C1—C2	123.0 (6)
O1—Zn—O2 ⁱ	94.62 (19)	C2—C1—Zn	165.9 (5)
O1 ⁱ —Zn—O2 ⁱ	57.27 (18)	C3—C2—C1	118.7 (7)
O2—Zn—O2 ⁱ	138.6 (3)	C7—C2—C1	121.6 (6)
O3 ⁱ —Zn—O1	100.6 (2)	C7—C2—C3	119.7 (7)
O3—Zn—O1	137.4 (2)	C2—C3—C4	120.0 (7)
O3 ⁱ —Zn—O1 ⁱ	137.4 (2)	C2—C3—H3	120.0
O3—Zn—O1 ⁱ	100.6 (2)	C4—C3—H3	120.0
O3 ⁱ —Zn—O2	127.7 (2)	C3—C4—H4	120.4
O3—Zn—O2	83.58 (18)	C5—C4—C3	119.1 (7)
O3 ⁱ —Zn—O2 ⁱ	83.58 (18)	C5—C4—H4	120.4
O3—Zn—O2 ⁱ	127.7 (2)	C4—C5—Br	117.6 (5)
O3 ⁱ —Zn—O3	90.0 (3)	C4—C5—C6	122.5 (7)
C1—O1—Zn	101.1 (5)	C6—C5—Br	119.8 (6)
C1—O2—Zn	81.5 (4)	C5—C6—C7	118.3 (7)
Zn—O3—H32	119 (6)	C5—C6—H6	120.8
Zn—O3—H31	131 (6)	C7—C6—H6	120.8
H32—O3—H31	106 (4)	C2—C7—C6	120.3 (7)
O1—C1—Zn	49.6 (4)	C2—C7—H7	119.8
O1—C1—C2	116.8 (7)	C6—C7—H7	119.8
O1 ⁱ —Zn—O1—C1	89.0 (4)	O1—C1—C2—C7	-19.9 (10)
O2—Zn—O1—C1	-0.8 (4)	O2—C1—C2—C7	162.8 (7)
O2 ⁱ —Zn—O1—C1	146.6 (4)	Zn—C1—C2—C3	146.9 (18)
O3 ⁱ —Zn—O1—C1	-129.1 (4)	Zn—C1—C2—C7	-33 (2)
O3—Zn—O1—C1	-27.4 (6)	O1—C1—C2—C3	160.3 (7)
C1 ⁱ —Zn—O1—C1	118.5 (4)	O2—C1—C2—C3	-17.0 (11)
O1—Zn—O2—C1	0.9 (4)	C1—C2—C3—C4	179.7 (7)
O1 ⁱ —Zn—O2—C1	-96.7 (4)	C7—C2—C3—C4	-0.1 (12)
O2 ⁱ —Zn—O2—C1	-53.4 (4)	C1—C2—C7—C6	-180.0 (7)
O3 ⁱ —Zn—O2—C1	78.2 (5)	C3—C2—C7—C6	-0.2 (12)
O3—Zn—O2—C1	163.1 (5)	C5—C4—C3—C2	-0.8 (12)
C1 ⁱ —Zn—O2—C1	-78.8 (6)	Br—C5—C4—C3	179.4 (6)
Zn—O1—C1—O2	1.6 (8)	C6—C5—C4—C3	1.9 (12)

supplementary materials

Zn—O1—C1—C2	-175.8 (5)	Br—C5—C6—C7	-179.6 (6)
Zn—O2—C1—O1	-1.3 (6)	C4—C5—C6—C7	-2.1 (12)
Zn—O2—C1—C2	175.9 (7)	C2—C7—C6—C5	1.2 (12)

Symmetry codes: (i) $-x+2, y, -z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H31 \cdots O2 ⁱⁱ	0.97 (7)	1.82 (6)	2.746 (7)	157 (9)
O3—H32 \cdots O1 ⁱⁱⁱ	0.95 (8)	1.86 (8)	2.765 (7)	160 (9)

Symmetry codes: (ii) $-x+2, -y, -z+1$; (iii) $-x+2, y-1, -z+1/2$.

Fig. 1

