

(E)-2-(2-Fluorobenzylidene)butanoic acid

Muhammad Niaz,^a M. Nawaz Tahir,^{b*} Zia-ur-Rehman,^a
Saqib Ali^a and Islam Ullah Khan^c

^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan,^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan, and^cDepartment of Chemistry, Government College University, Lahore, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 16 January 2008; accepted 15 March 2008

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.050; wR factor = 0.165; data-to-parameter ratio = 22.8.

In the crystal structure of the title compound, $\text{C}_{11}\text{H}_{11}\text{FO}_2$, the methine CH forms an intramolecular hydrogen bond with the carboxyl O atom. The molecules form dimers through hydrogen bonding between carboxyl groups. These dimers are linked to each other by C–H···O contacts between the benzene and carbonyl groups of adjoining molecules. In addition, there are weak intermolecular C–H···F contacts.

Related literature

For related literature, see: Burns & Hagaman (1993); Burt (2004); Forgó *et al.* (2005); Hertog *et al.* (1995); Muhammad *et al.* (2007). For details of the Cambridge Structural Database, see: Allen (2002).

**Experimental***Crystal data*

$\text{C}_{11}\text{H}_{11}\text{FO}_2$
 $M_r = 194.20$
Monoclinic, $P2_{1}/c$
 $a = 4.1895 (4)\text{ \AA}$
 $b = 17.4362 (19)\text{ \AA}$

$c = 13.8134 (15)\text{ \AA}$
 $\beta = 96.719 (3)^\circ$
 $V = 1002.12 (18)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation radiation

$\mu = 0.10\text{ mm}^{-1}$
 $T = 296 (2)\text{ K}$

$0.25 \times 0.18 \times 0.12\text{ mm}$

Data collection

Bruker Kappa APEXII CCD
diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)
 $T_{\min} = 0.935$, $T_{\max} = 0.958$

8632 measured reflections
2981 independent reflections
1704 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.164$
 $S = 1.04$
2981 reflections
131 parameters

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.41\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1···O2 ⁱ	0.97 (2)	1.66 (2)	2.6325 (18)	177.7 (12)
C3—H3···O1	0.93	2.32	2.713 (2)	105
C6—H6···O2 ⁱⁱ	0.93	2.53	3.421 (2)	160
C8—H8···F1 ⁱⁱⁱ	0.93	2.55	3.266 (2)	134 (1)

Symmetry codes: (i) $-x + 2, -y, -z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the Higher Education Commission, Islamabad, Pakistan, for funding the purchase of the diffractometer and Bana International, Karachi, Pakistan, for technical support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BV2091).

References

- Allen, F. H. (2002). *Acta Cryst. B* **58**, 380–388.
- Bruker (2005). *SADABS*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc. Madison, Wisconsin, USA.
- Burns, J. H. & Hagaman, E. W. (1993). *Acta Cryst. C* **49**, 1393–1396.
- Burt, S. (2004). *Int. J. Food Microbiol.* **94**, 223–253.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Forgó, P., Felföldi, K. & Pálinkó, I. (2005). *J. Mol. Struct.* **744–747**, 273–276.
- Hertog, M. G., Kromhout, D., Aravanis, C., Blackburn, H., Buzina, R., Fidanza, F., Giampaoli, S., Jansen, A., Menotti, A. & Nedeljkovic, S. (1995). *Arch. Intern. Med.* **155**, 381–386.
- Muhammad, N., Zia-ur-Rehman, Ali, S. & Meetsma, A. (2007). *Acta Cryst. E* **63**, o2174–o2175.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supporting information

Acta Cryst. (2008). E64, o733 [doi:10.1107/S1600536808007149]

(E)-2-(2-Fluorobenzylidene)butanoic acid

Muhammad Niaz, M. Nawaz Tahir, Zia-ur-Rehman, Saqib Ali and Islam Ullah Khan

S1. Comment


Cinnamic acid derivatives are well known for their antibacterial, antifungal antiparasitic (Burt, 2004), and antitumor activity (Hertog *et al.*, 1995). They are also used in the shikmic acid metabolic pathways of higher plants (Forgó *et al.*, 2005).

In the structure of the title compound (I), there are two C-atoms between the carboxylate and 2-fluorophenyl C-atoms. A search of CCDC (Allen, 2002) shows that a structure of 2-amino-2-(2-fluorophenyl)acetic acid (Burns *et al.*, 1993) has been published in which there is only a single C-atom between carboxylate and phenyl ring. Moreover, there is no structure of this kind with a different position for the F-atom.


The C1=O2 bond distance [1.2301 (18) Å], is significantly shorter than the C1—O1 distance [1.3006 (18) Å]. The C1—O1 bond lengthened due to the formation of intramolecular and intermolecular H-bonds. The value of C2=C3 is 1.334 (2) Å. The phenyl ring bond distances are in the normal range but the C4—C5—C6 bond angle is 124.27 (18)°, due to the influence of the F substituent attached to C5. The dihedral angle between the planes formed by (O1, C1, and O2) and (C2, C10, and C11) is 80.97 (18)°, and the dihedral angles between these planes and the phenyl ring are 52.88 (10)° and 67.17 (15)° respectively. The molecules are stabilized by intramolecular and intermolecular H-bonds. The title compound forms dimers through H-bonding, O1—H1···O2ⁱ [symmetry code i = -x + 2, -y, -z] as shown in Fig 2. These dimers are linked to each other through a C6—H6···O2ⁱⁱ interaction [symmetry code ii = -x + 1, y + 1/2, -z + 1/2]. Details of the H-bonding are given in Table 1. In addition there is a weak C8—H8···F1ⁱⁱⁱ intermolecular interaction [symmetry code iii = x, 1/2 - y, 1/2 + z] with a distance 3.2658 (25) Å between C8 and F1ⁱⁱⁱ.

S2. Experimental

Compound (I) was synthesized as reported earlier (Niaz, *et al.*, 2007). A mixture of 2-fluorobenzaldehyde (1.05 ml, 10 mmol), ethylmalonic acid (2.64 g, 20 mmol) and piperidine (1.98 ml, 20 mmol) in a pyridine (12.5 ml) solution was heated on a steam-bath for 24 h. The reaction mixture was cooled and added to a mixture of 25 ml of concentrated HCl and 50 g of ice. The precipitate formed in the acidified mixture was filtered off and washed with ice-cold water. The product was recrystallized from ethanol. The yield was 65%, m.p. 94 °C.

**Figure 1**

ORTEP drawing of the title compound, $C_{11}H_{11}F_1O_2$ with the atom numbering scheme. The thermal ellipsoids are drawn at the 50% probability level. H-atoms are shown by small circles of arbitrary radii.

**Figure 2**

The packing figure (PLATON: Spek, 2003) which shows the dimeric nature of the compound owing to inter molecular hydrogen bonding and also showing a link between dimers.

(E)-2-(2-Fluorobenzylidene)butanoic acid*Crystal data*

$C_{11}H_{11}FO_2$
 $M_r = 194.20$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 4.1895 (4)$ Å
 $b = 17.4362 (19)$ Å
 $c = 13.8134 (15)$ Å
 $\beta = 96.719 (3)^\circ$
 $V = 1002.12 (18)$ Å³
 $Z = 4$

$F(000) = 408$
 $D_x = 1.287$ Mg m⁻³
Mo $K\alpha$ radiation radiation, $\lambda = 0.71073$ Å
Cell parameters from 2981 reflections
 $\theta = 2.3\text{--}30.6^\circ$
 $\mu = 0.10$ mm⁻¹
 $T = 296$ K
Prismatic, colourless
 $0.25 \times 0.18 \times 0.12$ mm

Data collection

Bruker KappaAPEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 7.2 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2005)
 $T_{\min} = 0.935$, $T_{\max} = 0.958$

8632 measured reflections
2981 independent reflections
1704 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$
 $\theta_{\max} = 30.6^\circ$, $\theta_{\min} = 2.3^\circ$
 $h = -5 \rightarrow 6$
 $k = -24 \rightarrow 23$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.164$
 $S = 1.04$
2981 reflections
131 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0731P)^2 + 0.1527P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.41$ e Å⁻³
 $\Delta\rho_{\min} = -0.22$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.5705 (5)	0.28288 (7)	0.22076 (10)	0.1113 (6)
O1	1.0669 (3)	0.08331 (7)	0.07503 (9)	0.0575 (4)
H1	1.104 (5)	0.0617 (11)	0.0126 (15)	0.069*

O2	0.8191 (3)	-0.02692 (7)	0.09434 (9)	0.0609 (4)
C1	0.8986 (4)	0.03770 (8)	0.12367 (11)	0.0406 (4)
C2	0.8045 (3)	0.06721 (8)	0.21699 (11)	0.0394 (4)
C3	0.8530 (4)	0.14131 (9)	0.23791 (12)	0.0443 (4)
H3	0.9555	0.1699	0.1938	0.053*
C4	0.7616 (4)	0.18258 (9)	0.32326 (11)	0.0449 (4)
C5	0.6226 (5)	0.25413 (10)	0.31195 (13)	0.0593 (5)
C6	0.5260 (6)	0.29671 (11)	0.38672 (16)	0.0705 (6)
H6	0.4286	0.3443	0.3749	0.085*
C7	0.5763 (6)	0.26755 (11)	0.47922 (15)	0.0659 (6)
H7	0.5120	0.2953	0.5311	0.079*
C8	0.7214 (5)	0.19750 (12)	0.49543 (14)	0.0691 (6)
H8	0.7586	0.1781	0.5585	0.083*
C9	0.8126 (5)	0.15554 (11)	0.41831 (13)	0.0601 (5)
H9	0.9105	0.1081	0.4304	0.072*
C10	0.6348 (4)	0.01081 (9)	0.27580 (12)	0.0458 (4)
H10A	0.4779	-0.0170	0.2319	0.055*
H10B	0.5185	0.0392	0.3209	0.055*
C11	0.8524 (5)	-0.04706 (11)	0.33354 (14)	0.0614 (5)
H11A	0.7248	-0.0808	0.3682	0.092*
H11B	1.0041	-0.0205	0.3791	0.092*
H11C	0.9657	-0.0764	0.2897	0.092*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.2170 (19)	0.0646 (8)	0.0569 (8)	0.0505 (10)	0.0350 (9)	0.0161 (6)
O1	0.0854 (9)	0.0450 (7)	0.0468 (7)	-0.0124 (6)	0.0275 (6)	-0.0072 (5)
O2	0.0912 (10)	0.0434 (7)	0.0528 (7)	-0.0149 (6)	0.0288 (7)	-0.0118 (5)
C1	0.0485 (8)	0.0360 (8)	0.0382 (8)	0.0015 (6)	0.0091 (7)	0.0007 (6)
C2	0.0426 (8)	0.0401 (8)	0.0357 (7)	0.0046 (6)	0.0061 (6)	0.0003 (6)
C3	0.0549 (9)	0.0404 (8)	0.0382 (8)	0.0029 (6)	0.0085 (7)	-0.0005 (6)
C4	0.0575 (10)	0.0386 (8)	0.0393 (8)	0.0012 (7)	0.0088 (7)	-0.0041 (6)
C5	0.0947 (14)	0.0413 (9)	0.0438 (10)	0.0086 (9)	0.0156 (9)	0.0029 (7)
C6	0.1047 (17)	0.0438 (10)	0.0653 (13)	0.0165 (10)	0.0194 (12)	-0.0084 (9)
C7	0.0898 (15)	0.0588 (12)	0.0521 (11)	0.0015 (10)	0.0203 (10)	-0.0183 (9)
C8	0.0999 (16)	0.0685 (13)	0.0391 (9)	0.0122 (11)	0.0095 (10)	-0.0042 (9)
C9	0.0866 (14)	0.0525 (10)	0.0406 (9)	0.0170 (9)	0.0047 (9)	-0.0027 (8)
C10	0.0478 (9)	0.0460 (9)	0.0456 (9)	-0.0026 (7)	0.0136 (7)	-0.0018 (7)
C11	0.0694 (12)	0.0513 (10)	0.0654 (12)	-0.0014 (8)	0.0163 (10)	0.0179 (9)

Geometric parameters (\AA , $^\circ$)

F1—C5	1.350 (2)	C6—H6	0.9300
O1—C1	1.3006 (18)	C7—C8	1.371 (3)
O1—H1	0.97 (2)	C7—H7	0.9300
O2—C1	1.2301 (18)	C8—C9	1.383 (2)
C1—C2	1.483 (2)	C8—H8	0.9300

C2—C3	1.334 (2)	C9—H9	0.9300
C2—C10	1.506 (2)	C10—C11	1.521 (2)
C3—C4	1.470 (2)	C10—H10A	0.9700
C3—H3	0.9300	C10—H10B	0.9700
C4—C5	1.378 (2)	C11—H11A	0.9600
C4—C9	1.388 (2)	C11—H11B	0.9600
C5—C6	1.371 (3)	C11—H11C	0.9600
C6—C7	1.368 (3)		
C1—O1—H1	112.0 (12)	C6—C7—H7	120.0
O2—C1—O1	122.15 (14)	C8—C7—H7	120.0
O2—C1—C2	120.97 (14)	C7—C8—C9	120.19 (18)
O1—C1—C2	116.88 (13)	C7—C8—H8	119.9
C3—C2—C1	118.35 (14)	C9—C8—H8	119.9
C3—C2—C10	125.80 (14)	C8—C9—C4	121.56 (17)
C1—C2—C10	115.63 (13)	C8—C9—H9	119.2
C2—C3—C4	126.91 (15)	C4—C9—H9	119.2
C2—C3—H3	116.5	C2—C10—C11	115.09 (14)
C4—C3—H3	116.5	C2—C10—H10A	108.5
C5—C4—C9	115.53 (15)	C11—C10—H10A	108.5
C5—C4—C3	119.92 (15)	C2—C10—H10B	108.5
C9—C4—C3	124.53 (14)	C11—C10—H10B	108.5
F1—C5—C6	118.18 (17)	H10A—C10—H10B	107.5
F1—C5—C4	117.52 (16)	C10—C11—H11A	109.5
C6—C5—C4	124.27 (18)	C10—C11—H11B	109.5
C7—C6—C5	118.38 (18)	H11A—C11—H11B	109.5
C7—C6—H6	120.8	C10—C11—H11C	109.5
C5—C6—H6	120.8	H11A—C11—H11C	109.5
C6—C7—C8	120.03 (18)	H11B—C11—H11C	109.5
O2—C1—C2—C3	-169.70 (16)	C3—C4—C5—C6	179.3 (2)
O1—C1—C2—C3	10.1 (2)	F1—C5—C6—C7	179.6 (2)
O2—C1—C2—C10	5.2 (2)	C4—C5—C6—C7	1.5 (4)
O1—C1—C2—C10	-174.93 (14)	C5—C6—C7—C8	0.3 (4)
C1—C2—C3—C4	176.46 (15)	C6—C7—C8—C9	-1.0 (4)
C10—C2—C3—C4	2.1 (3)	C7—C8—C9—C4	0.1 (3)
C2—C3—C4—C5	-136.30 (19)	C5—C4—C9—C8	1.5 (3)
C2—C3—C4—C9	45.5 (3)	C3—C4—C9—C8	179.80 (19)
C9—C4—C5—F1	179.57 (19)	C3—C2—C10—C11	-106.93 (19)
C3—C4—C5—F1	1.2 (3)	C1—C2—C10—C11	78.57 (19)
C9—C4—C5—C6	-2.4 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1—H1 \cdots O2 ²	0.97 (2)	1.66 (2)	2.6325 (18)	177.7 (12)
C3—H3 \cdots O1	0.93	2.32	2.713 (2)	105

C6—H6···O2 ⁱⁱ	0.93	2.53	3.421 (2)	160
C8—H8···F1 ⁱⁱⁱ	0.93	2.55	3.266 (2)	134 (1)

Symmetry codes: (i) $-x+2, -y, -z$; (ii) $-x+1, y+1/2, -z+1/2$; (iii) $x, -y+1/2, z+1/2$.