

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## Pyromellitic acid–sarcosine (1/2)

Sérgio R. Domingos, Manuela Ramos Silva,\* Nuno D. Martins, Ana Matos Beja and J. A. Paixão

CEMDRX, Physics Department, University of Coimbra, P-3004-516 Coimbra, Portugal

Correspondence e-mail: manuela@pollux.fis.uc.pt


Received 18 March 2008; accepted 3 April 2008

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.115; data-to-parameter ratio = 14.2.

The title compound,  $\text{C}_{10}\text{H}_6\text{O}_8 \cdot 2\text{C}_3\text{H}_7\text{NO}_2$ , crystallizes as an adduct with the acid and amino acid molecules in their neutral forms. The asymmetric unit contains one half of a centrosymmetric pyromellitic acid molecule and one sarcosine molecule. The sarcosine has the amine group protonated and the carboxyl group deprotonated, as is usual for amino acids (zwitterionic form). The pyromellitic acid molecules retain the four carboxyl H atoms with the carboxyl groups rotated out of the ring plane [O—C—C—C torsion angles =  $24.1$  (3) and  $61.6$  (2)°]. There is a three-dimensional hydrogen-bond network linking the molecules.

## Related literature

For related compounds, see: Yaghi *et al.* (1997); Arora & Pedireddi (2003); Rochon & Massarweh (2001); Kumagai *et al.* (2003).



## Experimental

## Crystal data

$\text{C}_{10}\text{H}_6\text{O}_8 \cdot 2\text{C}_3\text{H}_7\text{NO}_2$   
 $M_r = 432.34$   
 Monoclinic,  $P2_1/c$   
 $a = 8.8894$  (3) Å  
 $b = 5.4118$  (2) Å  
 $c = 20.2205$  (7) Å  
 $\beta = 104.388$  (2)°

$V = 942.25$  (6) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.13$  mm<sup>-1</sup>  
 $T = 293$  (2) K  
 $0.47 \times 0.10 \times 0.07$  mm

## Data collection

Bruker APEX CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 2000)  
 $T_{\min} = 0.915$ ,  $T_{\max} = 0.998$   
 16656 measured reflections  
 2351 independent reflections  
 1643 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.047$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.115$   
 $S = 1.01$   
 2351 reflections  
 166 parameters  
 Only H-atom coordinates refined  
 $\Delta\rho_{\text{max}} = 0.24$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.21$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O2}-\text{H2} \cdots \text{O5}^i$	0.93 (2)	1.61 (2)	2.5216 (18)	166 (2)
$\text{O3}-\text{H3} \cdots \text{O5}$	0.98 (2)	1.62 (2)	2.6026 (18)	179 (2)
$\text{N1}-\text{H1A} \cdots \text{O6}^{ii}$	0.87 (2)	2.11 (2)	2.854 (2)	143.4 (17)
$\text{N1}-\text{H1A} \cdots \text{O1}^{iii}$	0.87 (2)	2.28 (2)	2.7262 (19)	111.8 (15)
$\text{N1}-\text{H1B} \cdots \text{O4}^{iv}$	0.88 (2)	2.15 (2)	2.917 (2)	145.5 (16)

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $-x, y-\frac{1}{2}, -z+\frac{3}{2}$ ; (iii)  $-x, y+\frac{1}{2}, -z+\frac{3}{2}$ ; (iv)  $-x+1, y+\frac{1}{2}, -z+\frac{3}{2}$ .

Data collection: SMART (Bruker, 2003); cell refinement: SAINT-Plus (Bruker, 2003); data reduction: SAINT-Plus (Bruker, 2003); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEPIII (Burnett & Johnson, 1996), ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2003); software used to prepare material for publication: SHELXL97.

This work was supported by Fundação para a Ciência e a Tecnologia (FCT) under project POCI/FIS/57876/2004.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2326).

## References

- Arora, K. K. & Pedireddi, V. R. (2003). *J. Org. Chem.* **68**, 9177–9185.  
 Bruker (2003). SMART and SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Burnett, M. N. & Johnson, C. K. (1996). ORTEPIII. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Kumagai, H., Chapman, K. W., Kepert, C. J. & Kurmoo, M. (2003). *Polyhedron*, **22**, 1921–1927.  
 Rochon, F. D. & Massarweh, G. (2001). *Inorg. Chim. Acta*, **314**, 163–171.  
 Sheldrick, G. M. (2000). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.  
 Yaghi, O. M., Davis, C. E., Li, G. M. & Li, H. L. (1997). *J. Am. Chem. Soc.* **119**, 2861–2868.

**supplementary materials**

*Acta Cryst.* (2008). E64, o826 [ doi:10.1107/S1600536808009045 ]

## Pyromellitic acid-sarcosine (1/2)

S. R. Domingos, M. Ramos Silva, N. D. Martins, A. Matos Beja and J. A. Paixão

### Comment

1,2,4,5-benzenetetracarboxylic acid (pyromellitic) is frequently chosen as a building block for crystal engineering due to its predictable properties and interesting supramolecular properties: It has provided three-dimensional porous networks (Yaghi *et al.*, 1997), host-guest systems (Arora & Pedireddi, 2003), mixed metallic systems (Rochon & Massarweh, 2001) and complex magnetic behaviours (Kumagai *et al.*, 2003). In an attempt to synthesize a low dimensional compound with copper, 1,2,4,5-benzenetetracarboxylic acid and sarcosine (as an auxiliary ligand), we have obtained the title compound, (I).

The midpoint of the acidic molecule lies on an inversion centre thus these molecules exhibit a  $C_i$  symmetry (Fig. 1). All four carboxylic groups retain the hydrogen atom and rotate around the C—C bond. Torsion angles O1—C1—C2—C4 24.1 (3)° and C4—C3—C5—O4 61.6 (2)° show different degrees of rotation. Sarcosine (*N*-methyl-glycine) crystallizes in the zwitterionic form with the amine group protonated and the carboxylic group deprotonated. The molecule when viewed along the C6—C7 bond shows the oxygen atoms anti to each other and the nitrogen atom synperiplanar to O6 [O6—C6—C7—N1 4.9 (2)°]. There is an extensive three-dimensional network of hydrogen bonds linking the molecules. Sarcosine molecules are assembled in chains *via* the N1—H1A···O6 bond (Table 1), running along the *b* axis. The chains are all interconnected through the remaining H bonds, since each sarcosine molecule is H-bonded to four benzenetetracarboxylic neighbours, (Fig. 2).

### Experimental

0.5 mmol of copper hydroxyfluoride were added to a 20 ml warmed ethanolic solution containing 1.5 mmol of 1,2,4,5-benzenetetracarboxylic acid and 1.5 mmol of sarcosine. After a few weeks, transparent, colourless crystals could be isolated from the solution.

### Refinement

H atoms coordinates were located from a difference Fourier map and refined freely. The  $U_{iso}(H)$  were restrained to be  $1.2U_{eq}(\text{parent atom})$ .

### Figures




Fig. 1. View of the title compound (I). Displacement ellipsoids are drawn at the 50% level. H atoms are represented as small spheres of arbitrary radii. H bond is represented as dashed line. [Symmetry code: (i) 1-x, 1-y, 2-z].




Fig. 2. Packing view of the title compound down the *b* axis. Hydrogen bonds are depicted as dashed lines. H atoms not involved in hydrogen bondings have been omitted for clarity. [Symmetry codes: (ii)  $-x, y-1/2, -z+3/2$ ; (iii)  $-x, y+1/2, -z+3/2$ ; (iv)  $-x+1, y+1/2, -z+3/2$ ]

## Pyromellitic acid–sarcosine (1/2)

### Crystal data

$C_{10}H_6O_8 \cdot 2C_3H_7NO_2$

$M_r = 432.34$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2ybc$

$a = 8.8894\ (3)\ \text{\AA}$

$b = 5.4118\ (2)\ \text{\AA}$

$c = 20.2205\ (7)\ \text{\AA}$

$\beta = 104.388\ (2)^\circ$

$V = 942.25\ (6)\ \text{\AA}^3$

$Z = 2$

$F_{000} = 452$

$D_x = 1.524\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2870 reflections

$\theta = 2.4\text{--}23.7^\circ$

$\mu = 0.13\ \text{mm}^{-1}$

$T = 293\ (2)\ \text{K}$

Prism, colourless

$0.47 \times 0.10 \times 0.07\ \text{mm}$

### Data collection

Bruker APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293\ (2)\ \text{K}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 2000)

$T_{\min} = 0.915, T_{\max} = 0.998$

16656 measured reflections

2351 independent reflections

1643 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$

$\theta_{\max} = 28.4^\circ$

$\theta_{\min} = 2.1^\circ$

$h = -11 \rightarrow 11$

$k = -7 \rightarrow 7$

$l = -27 \rightarrow 27$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.115$

$S = 1.01$

2351 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

Only H-atom coordinates refined

$w = 1/[\sigma^2(F_o^2) + (0.056P)^2 + 0.2509P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.25\ \text{e \AA}^{-3}$

166 parameters

$$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.19992 (15)	0.1100 (3)	0.89313 (8)	0.0531 (4)
O2	0.43230 (15)	-0.0346 (2)	0.88997 (7)	0.0415 (4)
H2	0.389 (3)	-0.163 (4)	0.8612 (11)	0.050*
C1	0.33933 (19)	0.1205 (3)	0.90904 (8)	0.0291 (4)
C2	0.42466 (17)	0.3192 (3)	0.95451 (8)	0.0236 (3)
C3	0.57683 (17)	0.3865 (3)	0.95565 (8)	0.0239 (3)
C4	0.65079 (18)	0.5655 (3)	1.00139 (8)	0.0256 (3)
H1	0.758 (2)	0.621 (3)	1.0018 (9)	0.031*
C5	0.66506 (19)	0.2833 (3)	0.90749 (8)	0.0284 (4)
O4	0.78636 (14)	0.1742 (3)	0.92796 (7)	0.0418 (4)
O3	0.61224 (16)	0.3342 (3)	0.84210 (6)	0.0382 (3)
H3	0.515 (2)	0.430 (4)	0.8318 (11)	0.046*
O5	0.35723 (14)	0.5924 (2)	0.81422 (6)	0.0377 (3)
O6	0.12981 (14)	0.7404 (2)	0.75521 (6)	0.0365 (3)
C6	0.23381 (18)	0.5845 (3)	0.76612 (8)	0.0261 (4)
C7	0.21727 (19)	0.3602 (3)	0.71983 (9)	0.0284 (4)
H7A	0.205 (2)	0.202 (4)	0.7448 (9)	0.034*
H7B	0.310 (2)	0.343 (3)	0.7009 (9)	0.034*
N1	0.07624 (17)	0.3848 (3)	0.66375 (8)	0.0283 (3)
H1A	-0.004 (2)	0.404 (4)	0.6805 (10)	0.034*
H1B	0.087 (2)	0.516 (4)	0.6393 (10)	0.034*
C8	0.0406 (3)	0.1733 (4)	0.61676 (12)	0.0437 (5)
H8A	-0.047 (3)	0.212 (4)	0.5816 (12)	0.052*
H8B	0.123 (3)	0.159 (4)	0.5931 (11)	0.052*
H8C	0.033 (3)	0.037 (5)	0.6395 (12)	0.052*

### Atomic displacement parameters ( $\text{\AA}^2$ )

$U^{11}$        $U^{22}$        $U^{33}$        $U^{12}$        $U^{13}$        $U^{23}$

## supplementary materials

---

O1	0.0296 (7)	0.0604 (10)	0.0672 (10)	-0.0132 (6)	0.0080 (6)	-0.0350 (8)
O2	0.0364 (7)	0.0329 (7)	0.0517 (8)	-0.0029 (6)	0.0042 (6)	-0.0215 (6)
C1	0.0313 (9)	0.0282 (9)	0.0271 (8)	-0.0057 (7)	0.0061 (7)	-0.0040 (7)
C2	0.0250 (8)	0.0231 (8)	0.0213 (8)	-0.0009 (6)	0.0030 (6)	-0.0020 (6)
C3	0.0240 (7)	0.0247 (8)	0.0227 (8)	0.0016 (6)	0.0054 (6)	-0.0013 (6)
C4	0.0227 (7)	0.0282 (9)	0.0255 (8)	-0.0026 (6)	0.0052 (6)	-0.0021 (7)
C5	0.0275 (8)	0.0283 (9)	0.0298 (9)	-0.0015 (7)	0.0079 (7)	-0.0059 (7)
O4	0.0333 (7)	0.0487 (8)	0.0437 (8)	0.0126 (6)	0.0105 (6)	-0.0056 (6)
O3	0.0410 (7)	0.0487 (8)	0.0271 (7)	0.0070 (6)	0.0128 (5)	-0.0032 (6)
O5	0.0416 (7)	0.0326 (7)	0.0321 (7)	0.0032 (6)	-0.0034 (5)	-0.0080 (5)
O6	0.0374 (7)	0.0330 (7)	0.0393 (7)	0.0073 (6)	0.0098 (6)	-0.0068 (6)
C6	0.0295 (8)	0.0253 (8)	0.0242 (8)	-0.0017 (7)	0.0079 (6)	-0.0009 (7)
C7	0.0292 (9)	0.0258 (9)	0.0274 (9)	0.0031 (7)	0.0019 (7)	-0.0028 (7)
N1	0.0255 (7)	0.0285 (8)	0.0292 (8)	0.0006 (6)	0.0036 (6)	-0.0038 (6)
C8	0.0486 (12)	0.0370 (11)	0.0390 (11)	0.0027 (10)	-0.0015 (10)	-0.0142 (9)

### *Geometric parameters (Å, °)*

O1—C1	1.202 (2)	O5—C6	1.273 (2)
O2—C1	1.302 (2)	O6—C6	1.230 (2)
O2—H2	0.93 (2)	C6—C7	1.518 (2)
C1—C2	1.493 (2)	C7—N1	1.473 (2)
C2—C4 <sup>i</sup>	1.390 (2)	C7—H7A	1.02 (2)
C2—C3	1.396 (2)	C7—H7B	0.995 (19)
C3—C4	1.387 (2)	N1—C8	1.471 (2)
C3—C5	1.502 (2)	N1—H1A	0.87 (2)
C4—C2 <sup>i</sup>	1.390 (2)	N1—H1B	0.88 (2)
C4—H1	0.997 (18)	C8—H8A	0.94 (2)
C5—O4	1.209 (2)	C8—H8B	0.97 (2)
C5—O3	1.318 (2)	C8—H8C	0.88 (2)
O3—H3	0.98 (2)		
C1—O2—H2	118.4 (14)	O5—C6—C7	115.53 (14)
O1—C1—O2	125.37 (16)	N1—C7—C6	109.64 (13)
O1—C1—C2	122.06 (16)	N1—C7—H7A	106.5 (11)
O2—C1—C2	112.57 (14)	C6—C7—H7A	112.0 (10)
C4 <sup>i</sup> —C2—C3	119.49 (14)	N1—C7—H7B	109.9 (11)
C4 <sup>i</sup> —C2—C1	117.78 (14)	C6—C7—H7B	110.5 (11)
C3—C2—C1	122.70 (14)	H7A—C7—H7B	108.2 (15)
C4—C3—C2	119.36 (14)	C8—N1—C7	115.63 (15)
C4—C3—C5	117.05 (14)	C8—N1—H1A	106.3 (13)
C2—C3—C5	123.53 (14)	C7—N1—H1A	109.5 (13)
C3—C4—C2 <sup>i</sup>	121.14 (14)	C8—N1—H1B	107.4 (13)
C3—C4—H1	120.5 (10)	C7—N1—H1B	108.1 (13)
C2 <sup>i</sup> —C4—H1	118.3 (11)	H1A—N1—H1B	109.8 (18)
O4—C5—O3	120.72 (15)	N1—C8—H8A	108.6 (14)
O4—C5—C3	121.66 (15)	N1—C8—H8B	108.0 (14)
O3—C5—C3	117.36 (14)	H8A—C8—H8B	103.5 (18)
C5—O3—H3	113.4 (12)	N1—C8—H8C	110.2 (16)

O6—C6—O5	125.48 (16)	H8A—C8—H8C	115 (2)
O6—C6—C7	118.98 (15)	H8B—C8—H8C	111 (2)
O1—C1—C2—C4 <sup>i</sup>	24.1 (3)	C5—C3—C4—C2 <sup>i</sup>	176.64 (15)
O2—C1—C2—C4 <sup>i</sup>	-155.56 (15)	C4—C3—C5—O4	61.6 (2)
O1—C1—C2—C3	-157.90 (18)	C2—C3—C5—O4	-121.1 (2)
O2—C1—C2—C3	22.4 (2)	C4—C3—C5—O3	-112.73 (18)
C4 <sup>i</sup> —C2—C3—C4	0.7 (3)	C2—C3—C5—O3	64.5 (2)
C1—C2—C3—C4	-177.23 (15)	O6—C6—C7—N1	4.9 (2)
C4 <sup>i</sup> —C2—C3—C5	-176.47 (15)	O5—C6—C7—N1	-175.62 (15)
C1—C2—C3—C5	5.5 (3)	C6—C7—N1—C8	-176.75 (17)
C2—C3—C4—C2 <sup>i</sup>	-0.8 (3)		

Symmetry codes: (i)  $-x+1, -y+1, -z+2$ .

*Hydrogen-bond geometry* ( $\text{\AA}, ^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O2—H2 $\cdots$ O5 <sup>ii</sup>	0.93 (2)	1.61 (2)	2.5216 (18)	166 (2)
O3—H3 $\cdots$ O5	0.98 (2)	1.62 (2)	2.6026 (18)	179 (2)
N1—H1A $\cdots$ O6 <sup>iii</sup>	0.87 (2)	2.11 (2)	2.854 (2)	143.4 (17)
N1—H1A $\cdots$ O1 <sup>iv</sup>	0.87 (2)	2.28 (2)	2.7262 (19)	111.8 (15)
N1—H1B $\cdots$ O4 <sup>v</sup>	0.88 (2)	2.15 (2)	2.917 (2)	145.5 (16)

Symmetry codes: (ii)  $x, y-1, z$ ; (iii)  $-x, y-1/2, -z+3/2$ ; (iv)  $-x, y+1/2, -z+3/2$ ; (v)  $-x+1, y+1/2, -z+3/2$ .

Fig. 1




Fig. 2

