

Three-dimensional hydrogen-bonded supramolecular assembly in tetrakis-(1,3,5-triaza-7-phosphaadamantane)-copper(I) chloride hexahydrate

Alexander M. Kirillov,^a Piotr Smoleński,^a M. Fátima C. Guedes da Silva,^{a,b*} Maximilian N. Kopylovich^a and Armando J. L. Pombeiro^a

^aCentro de Química Estrutural, Complexo Interdisciplinar, Instituto Superior Técnico, TU Lisbon, Av. Rovisco Pais, 1049-001 Lisbon, Portugal, and ^bUniversidade Lusófona de Humanidades e Tecnologias, ULHT Lisbon, Av. do Campo Grande 376, 1749-024 Lisbon, Portugal

Correspondence e-mail: fatima.guedes@ist.utl.pt


Received 22 March 2008; accepted 26 March 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{N}-\text{C}) = 0.003$ Å; R factor = 0.034; wR factor = 0.093; data-to-parameter ratio = 16.0.

The structure of the title compound, $[\text{Cu}(\text{PTA})_4]\text{Cl}\cdot 6\text{H}_2\text{O}$ (PTA is 1,3,5-triaza-7-phosphaadamantane, $\text{C}_6\text{H}_{12}\text{N}_3\text{P}$), is composed of discrete monomeric $[\text{Cu}(\text{PTA})_4]^+$ cations, chloride anions and uncoordinated water molecules. The Cu^{I} atom exhibits tetrahedral coordination geometry, involving four symmetry-equivalent P-bound PTA ligands. The structure is extended to a regular three-dimensional supramolecular framework *via* numerous equivalent $\text{O}-\text{H}\cdots\text{N}$ hydrogen bonds between all solvent water molecules (six per cation) and all PTA N atoms, thus simultaneously bridging each $[\text{Cu}(\text{PTA})_4]^+$ cation with 12 neighbouring units in multiple directions. The study also shows that PTA can be a convenient ligand in crystal engineering for the construction of supramolecular architectures.

Related literature

For general background, see: Kirillov *et al.* (2007, 2008); Karabach *et al.* (2006); Di Nicola *et al.* (2007). For a comprehensive review of PTA chemistry, see: Phillips *et al.* (2004). For PTA-derived polymeric networks, see: Lidrissi *et al.* (2005); Frost *et al.* (2006); Mohr *et al.* (2006). For related compounds, see: Forward *et al.* (1996); Darensbourg *et al.* (1997, 1999).



Experimental

Crystal data

$[\text{Cu}(\text{C}_6\text{H}_{12}\text{N}_3\text{P})_4]\text{Cl}\cdot 6\text{H}_2\text{O}$
 $M_r = 835.71$
 Cubic, $Fd\bar{3}m$
 $a = 19.795$ (4) Å
 $V = 7757$ (3) Å³

$Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.85$ mm⁻¹
 $T = 150$ (2) K
 $0.20 \times 0.17 \times 0.12$ mm

Data collection

Bruker APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\text{min}} = 0.848$, $T_{\text{max}} = 0.905$
 3022 measured reflections
 447 independent reflections
 361 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.092$
 $S = 1.08$
 447 reflections

28 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.75$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}10-\text{H}10\cdots\text{N}1$	0.81	2.04	2.843 (3)	174

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SIR97 (Altomare *et al.*, 1999); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP III (Burnett & Johnson, 1996), PLATON (Spek, 2003) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXL97.

This work has been supported by the FCT, Portugal, and its POCI 2010 programme (FEDER funded).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2329).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bruker (2004). *APEX2* and *S SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEP III*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Darensbourg, D. J., Decuir, T. J., Stafford, N. W., Robertson, J. B., Draper, J. D., Reibenspies, J. H., Katho, A. & Joo, F. (1997). *Inorg. Chem.* **36**, 4218–4226.
- Darensbourg, D. J., Robertson, J. B., Larkins, D. L. & Reibenspies, J. H. (1999). *Inorg. Chem.* **38**, 2473–2481.
- Di Nicola, C., Karabach, Y. Y., Kirillov, A. M., Monari, M., Pandolfo, L., Pettinari, C. & Pombeiro, A. J. L. (2007). *Inorg. Chem.* **46**, 221–230.
- Forward, J. M., Assefa, Z., Staples, R. J. & Fackler, J. P. Jr (1996). *Inorg. Chem.* **35**, 16–22.
- Frost, B. J., Bautista, C. M., Huang, R. C. & Shearer, J. (2006). *Inorg. Chem.* **45**, 3481–3483.
- Karabach, Y. Y., Kirillov, A. M., da Silva, M. F. C. G., Kopylovich, M. N. & Pombeiro, A. J. L. (2006). *Cryst. Growth Des.* **6**, 2200–2203.
- Kirillov, A. M., Karabach, Y. Y., Haukka, M., Guedes da Silva, M. F. C., Sanchiz, J., Kopylovich, M. N. & Pombeiro, A. J. L. (2008). *Inorg. Chem.* **47**, 162–175.
- Kirillov, A. M., Smoleński, P., Guedes da Silva, M. F. C. & Pombeiro, A. J. L. (2007). *Eur. J. Inorg. Chem.* pp. 2686–2692.
- Lidrissi, C., Romerosa, A., Saoud, M., Serrano-Ruiz, M., Gonsalvi, L. & Peruzzini, M. (2005). *Angew. Chem. Int. Ed.* **44**, 2568–2572.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Mohr, F., Falvello, L. R. & Laguna, M. (2006). *Eur. J. Inorg. Chem.* pp. 3152–3154.
- Phillips, A. D., Gonsalvi, L., Romerosa, A., Vizza, F. & Peruzzini, M. (2004). *Coord. Chem. Rev.* **248**, 955–993.
- Sheldrick, G. M. (2003). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, m603-m604 [doi:10.1107/S1600536808008179]

Three-dimensional hydrogen-bonded supramolecular assembly in tetrakis(1,3,5-triaza-7-phosphaadamantane)copper(I) chloride hexahydrate

A. M. Kirillov, P. Smolenski, M. F. C. Guedes da Silva, M. N. Kopylovich and A. J. L. Pombeiro

Comment

1,3,5-triaza-7-phosphaadamantane (PTA) is a water soluble aminophosphine that has sparked recent interest in coordination chemistry in view of the significance of transition metal PTA complexes in aqueous phase catalysis, photochemistry and medicinal chemistry (Phillips *et al.*, 2004). Besides, PTA and its derivatives can also be convenient building blocks for the construction of polymeric networks (Lidriissi *et al.*, 2005; Frost *et al.*, 2006; Mohr *et al.*, 2006) due to several potentially available coordination sites, protonation ability of N atoms, and strong affinity towards hydrogen bonds. Nevertheless, the use of PTA ligands in crystal design and engineering has remained little explored. Hence, in pursuit of our recent studies directed towards the synthesis of new copper compounds including PTA complexes (Kirillov *et al.*, 2007) and various coordination polymers, supramolecular frameworks and host–guest systems with other ligands (Karabach *et al.*, 2006; Di Nicola *et al.*, 2007; Kirillov *et al.*, 2008), we have prepared compound (I) whose crystal structure and supramolecular features are reported herein.

The moiety formula of (I) consists of the $[\text{Cu}(\text{PTA})_4]^+$ cation (Fig. 1), one chloride anion and six symmetry equivalent crystallization water molecules. The $[\text{Cu}(\text{PTA})_4]^+$ unit possesses a very high symmetry, being generated from only five symmetry nonequivalent atoms (Cu1, P1, N1, C1 and C2). The Cu^I atom lies on $-43m$ site symmetry and its coordination environment is filled by four equivalent *P*-bound PTA ligands, arranged in a perfect tetrahedral coordination geometry with the corresponding P—Cu—P angles of 109.47 (2)°. The Cu—P bond distances of 2.2598 (6) Å as well as other bonding parameters within the cage-like PTA cores are comparable to those reported for tetrahedral PTA complexes of Cu (Kirillov *et al.*, 2007), Au (Forward *et al.*, 1996), Pt (Darensbourg *et al.*, 1999) and Ni (Darensbourg *et al.*, 1997).

An interesting feature of (I) consists in the extensive intermolecular hydrogen bonding that arises from only one type of O—H⋯N H-bond (Table 1). Hence, each crystallization water molecule (O10) repeatedly acts as a double H-bond donor bridging to two N1 atoms of two different $[\text{Cu}(\text{PTA})_4]^+$ units. This results in the extensive interlinkage in multiple directions of every monomeric copper unit with twelve neighbouring ones (Fig. 2), thus leading to the formation of a regular three-dimensional supramolecular framework (Fig. 3). That framework has the shortest Cu⋯Cu separation of 13.977 (1) Å and possesses the repeating channels (*ca* 4.8 Å diameter) filled by water molecules.

Experimental

To the ethanolic solution (5 ml) of CuCl₂ (27 mg, 0.20 mmol) was added solid PTA (126 mg, 0.80 mmol). The obtained mixture was refluxed for 3 h resulting in a white suspension. This was filtered off and the colourless filtrate was left to evaporate in a beaker in air and at ambient temperature. A small crop of the colourless X-ray quality crystals of (I) was formed in several days. ¹H NMR data are similar to those reported for $[\text{Cu}(\text{PTA})_4]\text{NO}_3$ (Kirillov *et al.*, 2007). FT–IR (KBr pellet), cm⁻¹: 3430 m, br and 3195 w [$\nu(\text{H}_2\text{O})$], 2940 m and 2901 m [$\nu_{\text{as}}(\text{C—H})$], 2863 m and 2808 w [$\nu_{\text{s}}(\text{C—H})$], 1645 w

supplementary materials

br [$\delta(\text{H}_2\text{O})$], 1437 m, 1413 m, 1365 m, 1296 s, 1242 s, 1180 m, 1105 m, 1037 w, 1015 s, 971 s, 906 w, 890 m, 808 s, 797 s, 744 m, 694 m, 670 w, 582 s, 551 w, 451 s, 406 m [PTA bands]. FAB-MS⁺ (*m*-nitrobenzylalcohol), *m/z*: 691 [Cu(PTA)₄]⁺.

Refinement

All H atoms attached to C atoms were fixed geometrically and treated as riding with C—H = 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atom of the water molecule were located in difference Fourier maps and included in the subsequent refinement using restraint (O—H = 0.82 (1)Å) with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. In the last stage of refinement, it was treated as riding on the O atom.

Figures




Fig. 1. Molecular view of the cation with the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen atoms are omitted for clarity. [Symmetry codes: (i) z, x, y ; (ii) y, z, x ; (iii) $-x+1/4, y, -z+1/4$; (iv) $-x+1/4, -y+1/4, z$; (v) $x, -y+1/4, -z+1/4$]




Fig. 2. Fragment of the crystal packing diagram of (I) showing the simultaneous multidimensional interlinkage of the central monomeric [Cu(PTA)₄]⁺ unit (black coloured) with twelve neighbouring ones (each represented by different colour) via repeating O10—H10...N1 hydrogen bonding interactions (black dashed lines) between crystallization water molecules O10 (coloured balls) and PTA N1 atoms. H and Cl atoms are omitted for clarity.




Fig. 3. Fragment of the crystal packing diagram of (I) (view along the *a* axis) showing the extensive hydrogen bonding interactions (pale blue dashed lines) resulting in the formation of a regular three-dimensional H-bonded supramolecular assembly. H atoms are omitted for clarity. Cu, green; P, orange; N, blue; C, grey; O, red (balls); Cl, yellow (balls).

tetrakis(1,3,5-triaza-7-phosphaadamantane)copper(I) chloride hexahydrate

Crystal data

[Cu(C₆H₁₂N₃P)₄]Cl·6H₂O

$M_r = 835.71$

Cubic, $Fd\bar{3}m$

Hall symbol: -F 4vw 2vw 3

$a = 19.795$ (4) Å

$b = 19.795$ (4) Å

$Z = 8$

$F_{000} = 3536$

$D_x = 1.431$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71069$ Å

Cell parameters from 743 reflections

$\theta = 2.9\text{--}27.0^\circ$

$c = 19.795(4) \text{ \AA}$
 $\alpha = 90^\circ$
 $\beta = 90^\circ$
 $\gamma = 90^\circ$
 $V = 7757(3) \text{ \AA}^3$
 $\mu = 0.85 \text{ mm}^{-1}$
 $T = 150(2) \text{ K}$
 Prism, colourless
 $0.20 \times 0.17 \times 0.12 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer
 Radiation source: fine-focus sealed tube
 Monochromator: graphite
 $T = 150(2) \text{ K}$
 φ and ω scans
 Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\min} = 0.848$, $T_{\max} = 0.905$
 3022 measured reflections
 447 independent reflections
 361 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$
 $\theta_{\max} = 27.0^\circ$
 $\theta_{\min} = 2.9^\circ$
 $h = -24 \rightarrow 23$
 $k = -16 \rightarrow 11$
 $l = -6 \rightarrow 25$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.092$
 $S = 1.08$
 447 reflections
 28 parameters
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0463P)^2 + 19.2954P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.75 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$
 Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
-----	-----	-----	----------------------------------	-----------

supplementary materials

C1	0.25075 (10)	0.15137 (15)	0.25075 (10)	0.0199 (6)	
H1A	0.2258	0.1228	0.2818	0.024*	0.50
H1B	0.2818	0.1228	0.2258	0.024*	0.50
C2	0.33080 (15)	0.24509 (11)	0.24509 (11)	0.0239 (7)	
H2A	0.3607	0.2726	0.2726	0.029*	
H2B	0.3587	0.2166	0.2166	0.029*	
N1	0.29002 (8)	0.20160 (12)	0.29002 (8)	0.0212 (6)	
Cu1	0.1250	0.1250	0.1250	0.0134 (3)	
P1	0.19090 (4)	0.19090 (4)	0.19090 (4)	0.0156 (3)	
Cl1	0.3750	0.3750	0.3750	0.0165 (5)	
O10	0.3750	0.12300 (14)	0.3750	0.0240 (7)	
H10	0.3521	0.1480	0.3521	0.036*	

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0193 (8)	0.0212 (14)	0.0193 (8)	-0.0005 (7)	-0.0053 (11)	-0.0005 (7)
C2	0.0193 (15)	0.0262 (10)	0.0262 (10)	-0.0036 (8)	-0.0036 (8)	-0.0027 (12)
N1	0.0218 (8)	0.0202 (13)	0.0218 (8)	-0.0019 (7)	-0.0056 (10)	-0.0019 (7)
Cu1	0.0134 (3)	0.0134 (3)	0.0134 (3)	0.000	0.000	0.000
P1	0.0156 (3)	0.0156 (3)	0.0156 (3)	-0.0009 (3)	-0.0009 (3)	-0.0009 (3)
Cl1	0.0165 (5)	0.0165 (5)	0.0165 (5)	0.000	0.000	0.000
O10	0.0255 (10)	0.0210 (16)	0.0255 (10)	0.000	-0.0083 (12)	0.000

Geometric parameters (\AA , $^\circ$)

C1—N1	1.482 (3)	C2—H2A	0.9700
C1—P1	1.849 (3)	C2—H2B	0.9700
C1—H1A	0.9700	Cu1—P1	2.2596 (13)
C1—H1B	0.9700	P1—C1 ⁱ	1.849 (3)
C2—N1 ⁱ	1.478 (2)	O10—H10	0.8104
C2—N1	1.478 (2)		
N1—C1—P1	112.8 (2)	P1—Cu1—P1 ^{iv}	109.5
N1—C1—H1A	109.0	P1 ⁱⁱⁱ —Cu1—P1 ^{iv}	109.5
P1—C1—H1B	109.0	P1—Cu1—P1 ^v	109.5
H1A—C1—H1B	107.8	P1 ⁱⁱⁱ —Cu1—P1 ^v	109.5
N1 ⁱ —C2—N1	113.7 (3)	P1 ^{iv} —Cu1—P1 ^v	109.5
N1—C2—H2A	108.8	C1 ⁱⁱ —P1—C1 ⁱ	97.57 (12)
N1—C2—H2B	108.8	C1 ⁱⁱ —P1—C1	97.57 (12)
H2A—C2—H2B	107.7	C1 ⁱ —P1—C1	97.57 (12)
C2 ⁱⁱ —N1—C2	108.5 (3)	C1 ⁱⁱ —P1—Cu1	119.70 (9)
C2 ⁱⁱ —N1—C1	111.21 (16)	C1 ⁱ —P1—Cu1	119.70 (9)
C2—N1—C1	111.21 (16)	C1—P1—Cu1	119.70 (9)
P1—Cu1—P1 ⁱⁱⁱ	109.5		

Symmetry codes: (i) z, x, y ; (ii) y, z, x ; (iii) $-x+1/4, y, -z+1/4$; (iv) $-x+1/4, -y+1/4, z$; (v) $x, -y+1/4, -z+1/4$.

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
O10—H10···N1	0.81	2.04	2.843 (3)	174

Fig. 1




Fig. 2




Fig. 3

