Acta Crystallographica Section E Structure Reports Online

ISSN 1600-5368

Bis[(2-pyridylmethyl)(triisopropylsilyl)amido]zinc(II)—toluene—tetrahydrofuran (4/2/1)

Christian Koch, Helmar Görls and Matthias Westerhausen*

Institute of Inorganic and Analytical Chemistry, Friedrich-Schiller-Universität Jena, August-Bebel-Strasse 2, D-07743 Jena, Germany Correspondence e-mail: m.we@uni-jena.de

Received 5 March 2008; accepted 2 April 2008

Key indicators: single-crystal X-ray study; T = 183 K; mean σ (C–C) = 0.010 Å; disorder in solvent or counterion; R factor = 0.062; wR factor = 0.174; data-to-parameter ratio = 21.5.

The transamination reaction of (2-pyridylmethyl)(triisopropylsilyl)amine with bis{bis(trimethylsilyl)amido}zinc(II) yields the colorless title solvate, $[Zn(C_{15}H_{27}N_2Si)_2] \cdot 0.5C_7H_8 \cdot 0.25C_4H_8O$. The title compound was crystallized from toluene and tetrahydrofuran. There are two independent molecules in the asymmetric unit. In each molecule, the Zn atom is tetrahedrally coordinated by four N atoms. The two molecules differ in the orientation of the isopropyl groups. The molecules show large N-Zn-N angles [143.0 (2) and 145.7 (2)° between the amide groups].

Related literature

For related literature, see: Koch et al. (2007), Westerhausen et al. (2001, 2002).





Experimental

Crystal data

$$\begin{split} & [Zn(C_{15}H_{27}N_2Si)_2] \cdot 0.5C_7H_8 \cdots \\ & 0.25C_4H_8O \\ & M_r = 656.44 \\ & \text{Monoclinic, } P2_1 \\ & a = 13.2410 \ (3) \text{ \AA} \\ & b = 17.6223 \ (4) \text{ \AA} \\ & c = 16.7978 \ (5) \text{ \AA} \end{split}$$

Data collection

Nonius KappaCCD diffractometer Absorption correction: none 26798 measured reflections

Refinement

$$\begin{split} R[F^2 > 2\sigma(F^2)] &= 0.061 & \text{H-atom parameters constrained} \\ wR(F^2) &= 0.173 & \Delta\rho_{\text{max}} &= 1.04 \text{ e } \text{ Å}^{-3} \\ S &= 1.01 & \Delta\rho_{\text{min}} &= -0.38 \text{ e } \text{ Å}^{-3} \\ 16235 \text{ reflections} & \text{Absolute structure: Flack (1983),} \\ 754 \text{ parameters} & 7003 \text{ Friedel Pairs} \\ 1 \text{ restraint} & \text{Flack parameter: } -0.026 (12) \end{split}$$

 $\beta = 91.018 \ (2)^{\circ}$

Z = 4

V = 3918.93 (17) Å³

 $0.04 \times 0.04 \times 0.03 \text{ mm}$

16235 independent reflections

11581 reflections with $I > 2\sigma(I)$

Mo $K\alpha$ radiation

 $\mu = 0.71 \text{ mm}^{-1}$

T = 183 (2) K

 $R_{\rm int} = 0.051$

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997); data reduction: *DENZO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

We thank the Deutsche Forschungsgemeinschaft (DFG, Bonn–Band Godesberg, Germany) for generous financial support. We also acknowledge funding by the Fonds der Chemischen Industrie (Frankfurt/Main, Germany).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2104).

References

- Flack, H. D. (1983). Acta Cryst. A39, 876-881.
- Koch, C., Malassa, A., Agthe, C., Görls, H., Biedermann, R. & Krautscheid, H. (2007). Z. Anorg. Allg. Chem. 633, 375–382.
- Nonius (1998). COLLECT, Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). Acta Cryst. A64, 112-122.
- Westerhausen, M., Bollwein, T., Makropoulos, N., Rotter, T. M., Habereder, T., Suter, M. & Nöth, H. (2001). Eur. J. Inorg. Chem. pp. 851–857.
- Westerhausen, M., Bollwein, T., Makropoulos, N., Schneiderbauer, S., Suter, M., Nöth, H., Mayer, P., Piotrowski, H., Polborn, K. & Pfitzner, A. (2002). *Eur. J. Inorg. Chem.* pp. 389–404.

supporting information

Acta Cryst. (2008). E64, m664 [doi:10.1107/S160053680800888X]

Bis[(2-pyridylmethyl)(triisopropylsilyl)amido]zinc(II)-toluene-tetrahydrofuran (4/2/1)

Christian Koch, Helmar Görls and Matthias Westerhausen

S1. Comment

In the past, metallated (2-pyridylmethyl)(trialkysilyl)amines were used for oxidative C–C coupling reactions in order to prepare tetradendate bases. Zincation of (2-pyridylmethyl)(trialkylsilyl)amine (**A**; trialkylsilyl=*tert*BuMe₂Si, *i*Pr₃Si) yields dimeric methylzinc-(2-pyridylmethyl)(trialkylsilyl)amide (**B**). Further addition of dimethylzinc to a toluene solution of **B** at raised temperatures yields the C–C coupling product bis(methylzinc)-[1,2-dipyridyl-1,2-bis(trialkylsilylamido)ethane] (Westerhausen *et al.*, 2002). Thermal decomposition of **B** at 150°C yields [bis {(2-pyridylmethyl)(*tert*-butyldimethyl-silyl)amide} zinc(II)] **C** (Westerhausen *et al.*, 2001, Westerhausen *et al.*, 2002). The transamination reaction of (2-pyridylmethyl)(*tert*-butylsilyl)amine with *M*[N(SiMe₃)₂]₂ (*M*=Mg, Mn, Fe, Co and Zn) leads to the formation of the corresponding homoleptic [bis {(2-pyridylmethyl)(*tert*-butylsilyl)amido} metal (II)] (Koch *et al.*, 2007). The synthesis of **1** is similar (Koch *et al.*, 2007). The metal ions are in distorted tetrahedral environments. These compounds crystallize isotypically and show large N(2)–M–N(2) angles between 142° (Mg) and 156° (Mn). Due to the lower steric demand of the silyl moiety in **1** the corresponding N(2 A)–ZnA–N(4 A) angle with the value of 145.65 (19)° is smaller than 149.95 (7)° in the analogous zinc bis[(2-pyridylmethyl)(*tert*-butylsilyl)amide] **C** (Koch *et al.* 2007).

S2. Experimental

All manipulations were carried out in an atmosphere of argon using standard Schlenk techniques. Toluene, THF and pentane were dried (Na/benzophenone) and distilled prior to use. 2-Pyridylmethylamine and butyllithium were purchased form Aldrich. Triisopropylchlorosilane was purchased from Merck.¹H-NMR and ¹³C-NMR spectra were recorded at C_6D_6 solution at ambient temperature on a Bruker AC 400 MHz s pectrometer and were referenced to deuterated benzene as an internal standard.

Zinc bis[(2-pyridylmethyl)(triisopropylsilyl)amide] was prepared according to a literature procedure (Koch *et al*.2007) and recrystallized from pentane, toluene and thf.

Spectroscopic data:

¹H NMR (200 MHz, [D₆]benzene) δ = 8.29 (d, ³*J*(H¹,H²) = 6.0, 1H, Pyr1); 6.79–7.12 (m, 1H, Pyr3); 6.78 (d, ³*J*(H⁴,H³) = 7.8, 1H, Pyr4); 6.44–6.49 (m, 1H, Pyr2); 4.52 (s, 2H, CH₂); 1.30 (s, 18H, SiCH*Me*₂); 1.27 (s, 3H, SiC*H*Me₂).

¹³C NMR (50 MHz, [D₆]benzene) δ = 165.28 (Pyr5); 144.94 (Pyr1); 137.70 (Pyr3); 122.47 (Pyr2); 122.10 (Pyr4); 53.17 (²*J*, CH2); 19.59 (Si*C*HMe₂); 14.00 (SiCHMe₂).

IR (cm⁻¹): 3376, 2925, 2854, 1592, 1571, 1463, 1377, 1251.

S3. Refinement

All hydrogen atoms were set to idealized positions and were refined with isotropic thermal parameters 1.2 times that of the attached atom (1.5 for methyl groups). The methyl groups of both molecules were allowed to rotate but not to tip. The

thf molecule was modeled with a 50% occupancy.



Figure 1

The molecular structure of **1**, showing 40% probability displacement ellipsoids and the atom-numbering scheme for molecule A H atoms have been omitted for clarity.



Figure 2

The molecular structure of **1**, showing 40% probability displacement ellipsoids and the atom-numbering scheme for molecule B H atoms have been omitted for clarity.

Bis[(2-pyridylmethyl)(triisopropylsilyl)amido]zinc(II)- toluene-tetrahydrofuran (4/2/1)

Crystal data	
$[Zn(C_{15}H_{27}N_2Si)_2] \cdot 0.5(C_7H_8) \cdot 0.25(C_4H_8O)$	F(000) = 1420
$M_r = 656.44$	$D_{\rm x} = 1.113 {\rm Mg m^{-3}}$
Monoclinic, $P2_1$	Mo <i>K</i> α radiation, $\lambda = 0.71073$ Å
Hall symbol: P 2yb	Cell parameters from 26798 reflections
a = 13.2410(3) Å	$\theta = 2.3 - 27.5^{\circ}$
b = 17.6223 (4) Å	$\mu = 0.72 \ \mathrm{mm^{-1}}$
c = 16.7978 (5) Å	T = 183 K
$\beta = 91.018 \ (2)^{\circ}$	Prism, colourless
$V = 3918.93 (17) Å^3$	$0.04 \times 0.04 \times 0.03 \text{ mm}$
Z = 4	
Data collection	
Nonius KappaCCD	11581 reflections with $I > 2\sigma(I)$
diffractometer	$R_{\rm int} = 0.051$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 27.5^{\circ}, \ \theta_{\text{min}} = 2.3^{\circ}$
Graphite monochromator	$h = -16 \rightarrow 17$
φ and ω scans	$k = -22 \rightarrow 21$
26798 measured reflections	$l = -21 \rightarrow 21$
16235 independent reflections	

Refinement

Refinement on F^2	Hydrogen site location: inferred from
Least-squares matrix: full	neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.061$	H-atom parameters constrained
$wR(F^2) = 0.173$	$w = 1/[\sigma^2(F_o^2) + (0.0995P)^2]$
S = 1.01	where $P = (F_o^2 + 2F_c^2)/3$
16235 reflections	$(\Delta/\sigma)_{\rm max} = 0.001$
754 parameters	$\Delta \rho_{\rm max} = 1.04 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta \rho_{\rm min} = -0.38 \text{ e} \text{ Å}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 7003 Friedel Pairs
Secondary atom site location: difference Fourier	Absolute structure parameter: -0.026 (12)
map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on F^2 , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\hat{A}^2)

				TT ¥/TT	$O_{\rm eff}(z_1)$
	x	У	Z	$U_{\rm iso} - U_{\rm eq}$	Occ. (<1)
Zn1A	0.33924 (4)	1.03552 (3)	0.86223 (3)	0.02695 (15)	
SilA	0.36403 (13)	1.01974 (8)	1.05506 (9)	0.0336 (3)	
Si2A	0.39898 (11)	1.19976 (8)	0.78840 (9)	0.0291 (3)	
N1A	0.4218 (3)	0.9452 (2)	0.8080 (3)	0.0288 (10)	
N2A	0.3895 (3)	0.9918 (2)	0.9601 (3)	0.0303 (10)	
N3A	0.1902 (3)	1.0067 (2)	0.8238 (3)	0.0283 (9)	
N4A	0.3154 (3)	1.1277 (2)	0.8040 (3)	0.0295 (10)	
C1A	0.4342 (4)	0.9266 (3)	0.7307 (3)	0.0361 (13)	
H1AA	0.4037	0.9576	0.6908	0.043*	
C2A	0.4888 (4)	0.8654 (3)	0.7081 (4)	0.0386 (14)	
H2AA	0.4946	0.8532	0.6533	0.046*	
C3A	0.5357 (4)	0.8209 (3)	0.7653 (4)	0.0391 (14)	
НЗАА	0.5751	0.7784	0.7505	0.047*	
C4A	0.5242 (4)	0.8395 (3)	0.8452 (4)	0.0375 (13)	
H4AA	0.5549	0.8094	0.8859	0.045*	
C5A	0.4672 (4)	0.9027 (3)	0.8644 (3)	0.0294 (11)	
C6A	0.4554 (4)	0.9278 (3)	0.9496 (3)	0.0364 (13)	
H6AA	0.4291	0.8846	0.9807	0.044*	
H6AB	0.5229	0.9408	0.9719	0.044*	
C7A	0.3222 (5)	0.9345 (3)	1.1157 (3)	0.0392 (14)	
H7AA	0.3820	0.8999	1.1185	0.047*	
C8A	0.2971 (8)	0.9531 (4)	1.2026 (4)	0.074 (3)	
H8AA	0.2888	0.9059	1.2325	0.111*	

H8AB	0.2342	0.9825	1.2040	0.111*
H8AC	0.3521	0.9829	1.2267	0.111*
C9A	0.2378 (5)	0.8886 (4)	1.0755 (4)	0.0481 (16)
H9AA	0.2219	0.8445	1.1085	0.072*
Н9АВ	0.2598	0.8714	1.0231	0.072*
H9AC	0.1775	0.9204	1.0691	0.072*
C10A	0.4798 (6)	1 0600 (4)	1 1099 (4)	0.072
H10A	0.4551	1.0833	1 1602	0.062*
C11A	0.1331 0.5289(7)	1.1234 (5)	1.0628 (6)	0.002
H11A	0.5257	1.1254 (5)	1.0026 (0)	0.115*
H11R	0.3757	1 1579	1.0270	0.115*
HIIC	0.4700	1.1016	1.0422	0.115*
C12A	0.5000	1.1010	1.0105	0.113
U12A	0.5393 (7)	1.0014(3)	1.1541 (5)	0.073(2) 0.100*
III2A III2D	0.0179	0.0721	1.1382	0.109*
П12Б	0.5804	0.9731	1.0870	0.109*
HI2C	0.5307	0.9001	1.1/29	0.109°
CI3A	0.2666 (6)	1.0980 (4)	1.04/5 (4)	0.0514 (17)
HI3A	0.2860	1.1296	1.0007	0.062*
CI4A	0.2652 (7)	1.1533 (5)	1.1203 (5)	0.072 (2)*
H14A	0.2125	1.1916	1.1117	0.107*
H14B	0.3310	1.1783	1.1259	0.107*
H14C	0.2512	1.1245	1.1688	0.107*
C15A	0.1574 (6)	1.0723 (5)	1.0309 (5)	0.069 (2)*
H15A	0.1152	1.1167	1.0190	0.104*
H15B	0.1315	1.0462	1.0778	0.104*
H15C	0.1557	1.0377	0.9852	0.104*
C16A	0.1368 (4)	0.9448 (3)	0.8358 (4)	0.0383 (13)
H16A	0.1669	0.9045	0.8653	0.046*
C17A	0.0373 (5)	0.9358 (4)	0.8068 (4)	0.0420 (14)
H17A	-0.0006	0.8914	0.8178	0.050*
C18A	-0.0032 (4)	0.9943 (4)	0.7615 (4)	0.0426 (15)
H18A	-0.0693	0.9896	0.7390	0.051*
C19A	0.0515 (4)	1.0587 (3)	0.7490 (3)	0.0373 (13)
H19A	0.0233	1.0995	0.7192	0.045*
C20A	0.1511 (4)	1.0638 (3)	0.7812 (3)	0.0297 (11)
C21A	0.2143 (4)	1.1347 (3)	0.7703 (4)	0.0348 (13)
H21A	0.2190	1.1459	0.7127	0.042*
H21B	0.1799	1.1781	0.7957	0.042*
C22A	0.4036 (5)	1.2254 (4)	0.6770 (3)	0.0423(14)
H22A	0.3369	1.2495	0.6642	0.051*
C23A	0 4814 (6)	1 2841 (4)	0.6574(5)	0.062(2)
H23A	0 4769	1 2962	0.6005	0.094*
H23R	0 5489	1 2644	0.6704	0.094*
H23C	0 4691	1 3302	0.6885	0.094*
C24A	0.4098(5)	1 1557 (4)	0.6000	0.0495 (16)
H^{24}	0.3026	1 1708	0.5672	0.074*
H24R	0.3520	1 1169	0.5072	0.074*
$H_2 H_2 H_2 H_2 H_2 H_2 H_2 H_2 H_2 H_2 $	0.785	1 1352	0.6230	0.074*
11270	0.7/05	1.1332	0.0457	0.0/ -

C25A	0.3677 (4)	1.2927 (3)	0.8403 (4)	0.0367 (13)
H25A	0.4292	1.3256	0.8395	0.044*
C26A	0.2796 (5)	1.3372 (4)	0.8017 (4)	0.0509 (17)
H26A	0.2645	1.3818	0.8343	0.076*
H26B	0.2198	1.3046	0.7980	0.076*
H26C	0.2984	1.3536	0.7482	0.076*
C27A	0.3398 (6)	1.2793 (4)	0.9278 (4)	0.0547 (18)
H27A	0.3273	1.3282	0.9536	0.082*
H27B	0.3956	1.2533	0.9555	0.082*
H27C	0.2788	1.2480	0.9299	0.082*
C28A	0.5249(4)	1.1648 (3)	0.8311 (4)	0.0374(13)
H28A	0.5080	1 1331	0.8785	0.045*
C29A	0.5978 (5)	1.1351	0.8620 (5)	0.019 0.0540(17)
H29A	0.6590	1.2205 (4)	0.8841	0.081*
H29R	0.5651	1.2560	0.9036	0.081*
H29C	0.5051	1.2500	0.9050	0.081*
C20A	0.0100	1.2333	0.0101 0.7742(4)	0.081°
	0.5805 (4)	1.1120 (4)	0.7742 (4)	0.04/4(10) 0.071*
	0.0443	1.0939	0.7990	0.071*
	0.5940	1.1391	0.7240	0.071*
H50C	0.5380	1.00/4	0.7627	$0.0/1^{+}$
Znib	0.16087(4)	0.72556 (3)	0.03234(3)	0.02721(15)
SIIB	-0.07885(10)	0.70358 (8)	0.59155 (9)	0.0284 (3)
S12B	0.31298 (11)	0.59836 (8)	0.55680 (9)	0.0267 (3)
NIB	0.1792 (3)	0.7365 (3)	0.7578 (3)	0.0299 (10)
N2B	0.0210 (3)	0.7084 (3)	0.6574 (2)	0.0290 (9)
N3B	0.2153 (3)	0.8345 (2)	0.5982 (3)	0.0288 (10)
N4B	0.2689 (3)	0.6892 (2)	0.5682 (3)	0.0297 (10)
C1B	0.2603 (4)	0.7557 (3)	0.8026 (4)	0.0355 (13)
H1BA	0.3217	0.7666	0.7765	0.043*
C2B	0.2587 (5)	0.7604 (3)	0.8840 (4)	0.0371 (13)
H2BA	0.3174	0.7745	0.9137	0.045*
C3B	0.1681 (5)	0.7437 (3)	0.9223 (3)	0.0392 (14)
H3BA	0.1643	0.7454	0.9787	0.047*
C4B	0.0845 (4)	0.7249 (3)	0.8760 (3)	0.0376 (13)
H4BA	0.0219	0.7144	0.9007	0.045*
C5B	0.0914 (4)	0.7212 (3)	0.7940 (3)	0.0291 (11)
C6B	0.0033 (4)	0.6978 (4)	0.7420 (3)	0.0363 (13)
H6BA	-0.0118	0.6436	0.7519	0.044*
H6BB	-0.0567	0.7277	0.7570	0.044*
C7B	-0.1881 (4)	0.7642 (4)	0.6286 (4)	0.0413 (14)
H7BA	-0.2121	0.7393	0.6783	0.050*
C8B	-0.1581 (5)	0.8458 (4)	0.6520 (4)	0.0527 (17)*
H8BA	-0.2125	0.8690	0.6824	0.079*
H8BB	-0.1465	0.8756	0.6038	0.079*
H8BC	-0.0960	0.8446	0.6847	0.079*
C9B	-0.2801 (5)	0.7663 (4)	0.5705 (4)	0.0527 (18)
H9BA	-0.3357	0.7939	0.5951	0.079*
H9BB	-0.3018	0.7143	0.5583	0.079*

H9BC	-0.2612	0.7919	0.5212	0.079*
C10B	-0.1340 (4)	0.6054 (3)	0.5768 (4)	0.0374 (13)
H10B	-0.1807	0.6078	0.5295	0.045*
C11B	-0.1974 (5)	0.5763 (4)	0.6491 (4)	0.0550 (18)
H11D	-0.2194	0.5240	0.6390	0.083*
H11E	-0.2567	0.6089	0.6555	0.083*
H11F	-0.1556	0.5780	0.6978	0.083*
C12B	-0.0527 (5)	0.5458 (4)	0.5587 (5)	0.0550 (18)
H12D	-0.0838	0.4954	0.5558	0.082*
H12E	-0.0008	0.5463	0.6010	0.082*
H12F	-0.0216	0.5577	0.5076	0.082*
C13B	-0.0279 (4)	0.7355 (3)	0.4918 (3)	0.0360 (13)
H13B	0.0408	0.7123	0.4882	0.043*
C14B	-0.0876(5)	0.7070 (5)	0.4170 (4)	0.059 (2)
H14D	-0.0551	0.7257	0.3688	0.088*
H14E	-0.1571	0.7259	0.4185	0.088*
H14F	-0.0882	0.6514	0.4165	0.088*
C15B	-0.0116 (6)	0.8202 (4)	0.4862 (4)	0.0530 (17)
H15D	0.0324	0.8313	0.4413	0.080*
H15E	0.0202	0.8386	0.5356	0.080*
H15F	-0.0768	0.8456	0.4779	0.080*
C16B	0.1863 (4)	0.9045 (3)	0.6184 (3)	0.0355 (13)
H16B	0.1341	0.9096	0.6560	0.043*
C17B	0.2275(5)	0.9694(3)	0.5879 (4)	0.0409 (14)
H17B	0.2054	1.0181	0.6046	0.049*
C18B	0.3011 (5)	0.9615 (3)	0.5330 (4)	0.0389 (14)
H18B	0 3307	1 0052	0 5098	0.047*
C19B	0.3321(4)	0.8916 (3)	0.5115 (4)	0.0382(14)
H19B	0.3838	0.8860	0.4734	0.046*
C20B	0 2881 (4)	0.8274(3)	0.5451 (3)	0.0301(11)
C21B	0.3212(5)	0.7483(3)	0.5743(4)	0.0301(11) 0.0395(14)
H21C	0.3094	0 7401	0.4666	0.047*
H21D	0 3947	0.7437	0.5351	0.047*
C22B	0.3917 0.4494 (4)	0.5906 (3)	0.5924(4)	0.0370(13)
H22B	0.4925	0.6080	0.5476	0.0370 (13)
C23B	0.4923 0.4844 (4)	0.5098(4)	0.6148 (4)	0.044
H23D	0.5572	0.5101	0.6264	0.0430 (13)
H23E	0.4699	0.4752	0.5704	0.067*
H23E	0.4483	0.4927	0.5704	0.067*
C24B	0.4734(5)	0.4927 0.6428 (4)	0.6644(4)	0.007
U24D	0.4754 (5)	0.6415	0.6764	0.073*
1124D 1124D	0.1363	0.6250	0.0704	0.073*
H24E	0.4500	0.0250	0.7109	0.073*
C25B	0.3109 (4)	0.5733 (3)	0.0517	0.073 0.0346 (12)
U25D	0.3103 (4)	0.5755 (5)	0.4409 (3)	0.0340 (12)
1123D C26D	0.3441 0.2025 (5)	0.01/4	0.4112 (4)	0.041
	0.2055 (5)	0.5727 (4)	0.4112(4)	0.0303 (10)
	0.2039	0.3///	0.3331	0.076*
П20E	0.1033	0.0132	0.4332	0.0/0.

H26F	0.1705	0.5248	0.4249	0.076*	
C27B	0.3714 (5)	0.5042 (4)	0.4192 (4)	0.0442 (15)	
H27D	0.3622	0.4972	0.3616	0.066*	
H27E	0.3476	0.4589	0.4470	0.066*	
H27F	0.4432	0.5122	0.4318	0.066*	
C28B	0.2242 (4)	0.5346 (3)	0.6128 (3)	0.0360 (12)	
H28B	0.1547	0.5518	0.5969	0.043*	
C29B	0.2278 (6)	0.4504 (4)	0.5911 (4)	0.0517 (18)	
H29D	0.1793	0.4224	0.6232	0.078*	
H29E	0.2960	0.4308	0.6015	0.078*	
H29F	0.2105	0.4442	0.5345	0.078*	
C30B	0.2298 (5)	0.5435 (4)	0.7044 (3)	0.0426 (14)	
H30D	0.1673	0.5246	0.7276	0.064*	
H30E	0.2386	0.5972	0.7180	0.064*	
H30F	0.2873	0.5144	0.7257	0.064*	
C1T	0.8334 (9)	0.9809 (6)	0.9947 (7)	0.095 (3)	
C2T	0.7634 (8)	0.9935 (6)	0.9348 (8)	0.107 (4)	
H2TC	0.7270	1.0399	0.9334	0.128*	
C3T	0.7458 (7)	0.9415 (6)	0.8788 (8)	0.098 (3)	
H3TC	0.7008	0.9534	0.8359	0.118*	
C4T	0.7899 (8)	0.8722 (6)	0.8807 (7)	0.090 (3)	
H4TC	0.7739	0.8353	0.8413	0.108*	
C5T	0.8572 (8)	0.8567 (6)	0.9402 (7)	0.095 (3)	
H5TA	0.8879	0.8080	0.9439	0.114*	
C6T	0.8809 (7)	0.9125 (5)	0.9957 (7)	0.083 (3)	
H6TA	0.9314	0.9027	1.0352	0.099*	
C7T	0.8582 (12)	1.0359 (10)	1.0535 (10)	0.160 (6)*	
H7TA	0.9090	1.0150	1.0905	0.240*	
H7TB	0.7974	1.0494	1.0828	0.240*	
H7TC	0.8853	1.0813	1.0278	0.240*	
O1TH	0.8602 (9)	1.2011 (7)	0.7855 (7)	0.079 (3)*	0.50
C1TH	0.9455 (13)	1.2052 (10)	0.8595 (11)	0.075 (5)*	0.50
H1TA	0.9120	1.2047	0.9117	0.090*	0.50
H1TB	0.9925	1.1616	0.8570	0.090*	0.50
C2TH	0.9996 (9)	1.2771 (7)	0.8477 (8)	0.042 (3)*	0.50
H2TA	1.0724	1.2669	0.8404	0.050*	0.50
H2TB	0.9924	1.3097	0.8952	0.050*	0.50
C3TH	0.9563 (9)	1.3178 (7)	0.7743 (7)	0.040 (3)*	0.50
H3TA	0.9321	1.3686	0.7903	0.048*	0.50
НЗТВ	1.0108	1.3250	0.7355	0.048*	0.50
C4TH	0.8709 (11)	1.2755 (8)	0.7348 (9)	0.057 (4)*	0.50
H4TA	0.8868	1.2634	0.6789	0.068*	0.50
H4TB	0.8079	1.3057	0.7357	0.068*	0.50

Atomic displacement parameters $(Å^2)$

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1A	0.0295 (3)	0.0243 (3)	0.0270 (3)	0.0012 (3)	-0.0014 (2)	0.0016 (3)

Acta Cryst. (2008). E64, m664

Sila	0.0493(9)	0 0249 (8)	0 0266 (8)	0.0012(7)	-0.0030(7)	-0.0014(6)
Si2A	0.0499(7)	0.0249(0)	0.0200(8)	-0.0012(7)	0.0030(7)	0.0014(0)
N1A	0.0300(7)	0.0240(7)	0.0333(0)	0.0003(0)	-0.0027(0)	-0.0021(0)
N2A	0.027(2)	0.031(2)	0.028(2)	0.0009(19)	-0.0021(19)	0.0012(19)
N2A	0.030(2)	0.027(2)	0.028(2)	0.0079(19)	-0.0021(19)	0.0007(18)
NJA NJA	0.030(2)	0.020(2)	0.028(2)	-0.0000(19)	-0.0018(10)	0.0020(18)
C1A	0.027(2)	0.028(2)	0.034(3)	-0.0008(18)	0.0018(19)	-0.0000(13)
C1A C2A	0.039(3)	0.041(3)	0.028(3)	-0.002(3)	0.004(2)	-0.012(3)
C2A C2A	0.039(3)	0.041(3)	0.030(3)	-0.002(3)	0.005(3)	-0.012(3)
C3A C4A	0.037(3)	0.037(3)	0.043(4)	0.008(3)	0.003(3)	-0.010(3)
C4A	0.042(3)	0.030(3)	0.040(3)	0.009(3)	-0.003(3)	-0.003(2)
CSA	0.025(2)	0.029(3)	0.034(3)	-0.001(2)	-0.002(2)	-0.003(2)
COA	0.042(3)	0.031(3)	0.030(3)	0.007(3)	-0.004(3)	-0.001(2)
C/A	0.056 (4)	0.035(3)	0.027(3)	0.002(3)	0.001(3)	-0.003(2)
C8A	0.141 (8)	0.053 (5)	0.029 (4)	-0.006 (5)	0.012 (4)	0.000 (3)
C9A	0.060 (4)	0.040 (4)	0.044 (4)	-0.008(3)	0.013 (3)	0.007 (3)
CIOA	0.076 (5)	0.041 (4)	0.038 (4)	-0.009(3)	-0.009 (3)	-0.002 (3)
CIIA	0.086 (6)	0.056 (5)	0.087 (6)	-0.026 (4)	-0.037 (5)	0.009 (4)
C12A	0.083 (5)	0.058 (4)	0.076 (6)	-0.005(4)	-0.051 (5)	-0.003(4)
C13A	0.074 (5)	0.038 (3)	0.042 (4)	0.009 (3)	0.015 (3)	0.005 (3)
C16A	0.038 (3)	0.033 (3)	0.044 (3)	0.002 (3)	0.000 (3)	0.003 (3)
C17A	0.043 (3)	0.037 (3)	0.046 (4)	-0.007(3)	0.003 (3)	-0.001 (3)
C18A	0.031 (3)	0.054 (4)	0.043 (4)	0.001 (3)	-0.005 (3)	-0.008(3)
C19A	0.035 (3)	0.042 (3)	0.035 (3)	0.003 (3)	-0.003 (2)	0.001 (3)
C20A	0.029 (3)	0.031 (3)	0.029 (3)	0.001 (2)	-0.001 (2)	-0.004 (2)
C21A	0.031 (3)	0.030 (3)	0.043 (3)	-0.001 (2)	-0.003 (2)	0.009 (3)
C22A	0.047 (3)	0.042 (3)	0.038 (3)	0.003 (3)	0.011 (3)	0.006 (3)
C23A	0.070 (5)	0.061 (5)	0.057 (5)	-0.008 (4)	0.023 (4)	0.020 (4)
C24A	0.053 (4)	0.057 (4)	0.038 (4)	0.009 (3)	0.009 (3)	0.003 (3)
C25A	0.038 (3)	0.028 (3)	0.043 (3)	0.004 (2)	0.007 (3)	-0.002 (3)
C26A	0.054 (4)	0.031 (3)	0.067 (5)	0.016 (3)	-0.001 (3)	-0.004 (3)
C27A	0.076 (5)	0.038 (3)	0.051 (4)	-0.001 (3)	0.019 (4)	-0.010 (3)
C28A	0.033 (3)	0.029 (3)	0.050 (4)	-0.002 (2)	-0.005 (3)	-0.004 (3)
C29A	0.042 (3)	0.047 (4)	0.073 (5)	-0.003 (3)	-0.007 (3)	-0.009(4)
C30A	0.033 (3)	0.039 (3)	0.069 (5)	0.009 (3)	-0.002 (3)	-0.005(3)
Zn1B	0.0253 (3)	0.0298 (3)	0.0266 (3)	0.0001 (3)	0.0029 (2)	0.0016 (3)
Si1B	0.0259 (7)	0.0312 (8)	0.0282 (7)	-0.0013 (6)	0.0005 (6)	-0.0007 (6)
Si2B	0.0284 (7)	0.0260 (7)	0.0259 (7)	-0.0008 (6)	0.0032 (6)	0.0002 (6)
N1B	0.028 (2)	0.034 (2)	0.028 (2)	0.0022 (19)	0.0008 (18)	0.0016 (19)
N2B	0.028 (2)	0.039 (3)	0.020 (2)	0.0020 (19)	0.0014 (17)	-0.0004 (19)
N3B	0.030 (2)	0.027 (2)	0.029 (2)	0.0024 (19)	-0.0003 (19)	0.0031 (19)
N4B	0.031 (2)	0.030 (2)	0.029 (2)	-0.0008 (19)	0.0068 (19)	0.0062 (19)
C1B	0.029 (3)	0.034 (3)	0.043 (3)	-0.004 (2)	-0.001(2)	-0.004(3)
C2B	0.040 (3)	0.034 (3)	0.037 (3)	-0.003(3)	-0.010(3)	-0.005(3)
C3B	0.051 (3)	0.039(3)	0.028 (3)	-0.006(3)	0.001 (3)	-0.001(2)
C4B	0.040(3)	0.041(3)	0.032(3)	0.002 (3)	0.013 (2)	0.007(3)
C5B	0.029(2)	0.029(3)	0.030(3)	-0.002(2)	0.001(2)	0.005(2)
C6B	0.021(2)	0.029(3)	0.024(3)	-0.002(2)	-0.001(2)	0.000(2)
C7B	0.031(3)	0.049(4)	0.027(3)	0.007(0)	-0.005(3)	-0.012(3)
0,0	0.000 (0)	עד) לו טוט	0.010(3)	0.010 (0)	0.000 (3)	0.012 (3)

C9B	0.038 (3)	0.060 (4)	0.060 (4)	0.017 (3)	-0.006 (3)	-0.012 (4)
C10B	0.030 (3)	0.045 (3)	0.037 (3)	-0.007 (3)	-0.005 (2)	0.000 (3)
C11B	0.057 (4)	0.060 (4)	0.047 (4)	-0.024 (4)	-0.001 (3)	0.011 (3)
C12B	0.045 (4)	0.041 (4)	0.078 (5)	0.003 (3)	-0.004 (3)	-0.014 (3)
C13B	0.039 (3)	0.040 (3)	0.029 (3)	-0.006 (3)	0.001 (2)	-0.004 (3)
C14B	0.059 (4)	0.086 (6)	0.031 (3)	-0.019 (4)	-0.001 (3)	0.001 (4)
C15B	0.062 (4)	0.049 (4)	0.047 (4)	-0.004 (3)	-0.010 (3)	0.011 (3)
C16B	0.037 (3)	0.034 (3)	0.035 (3)	0.004 (3)	0.003 (2)	-0.006 (2)
C17B	0.054 (4)	0.032 (3)	0.037 (3)	0.007 (3)	-0.003 (3)	-0.004 (3)
C18B	0.046 (3)	0.030 (3)	0.041 (3)	-0.004 (3)	0.001 (3)	0.007 (3)
C19B	0.041 (3)	0.035 (3)	0.038 (3)	0.002 (3)	0.009 (3)	0.007 (3)
C20B	0.026 (3)	0.031 (3)	0.033 (3)	0.002 (2)	-0.001 (2)	0.002 (2)
C21B	0.041 (3)	0.036 (3)	0.042 (3)	0.001 (3)	0.010 (3)	0.007 (3)
C22B	0.034 (3)	0.035 (3)	0.042 (3)	0.003 (2)	-0.001 (3)	-0.005 (3)
C23B	0.036 (3)	0.052 (4)	0.047 (4)	0.019 (3)	-0.004 (3)	0.005 (3)
C24B	0.041 (3)	0.045 (4)	0.059 (4)	0.003 (3)	-0.016 (3)	-0.017 (3)
C25B	0.040 (3)	0.033 (3)	0.030 (3)	0.003 (2)	0.003 (2)	-0.006 (2)
C26B	0.060 (4)	0.058 (4)	0.033 (3)	-0.003 (3)	-0.011 (3)	-0.008 (3)
C27B	0.056 (4)	0.041 (3)	0.036 (3)	-0.002 (3)	0.002 (3)	-0.007 (3)
C28B	0.032 (3)	0.042 (3)	0.035 (3)	-0.009 (3)	-0.001 (2)	0.009 (3)
C29B	0.062 (4)	0.049 (4)	0.044 (4)	-0.028 (3)	0.006 (3)	0.001 (3)
C30B	0.055 (4)	0.039 (3)	0.034 (3)	0.001 (3)	0.007 (3)	0.003 (3)
C1T	0.104 (8)	0.078 (7)	0.104 (8)	-0.032 (6)	0.027 (7)	-0.021 (6)
C2T	0.081 (7)	0.068 (6)	0.169 (12)	-0.011 (5)	-0.042 (8)	-0.005 (7)
C3T	0.066 (6)	0.083 (7)	0.145 (11)	-0.016 (5)	-0.022 (6)	0.007 (7)
C4T	0.084 (7)	0.093 (8)	0.092 (8)	-0.003 (6)	0.009 (6)	0.003 (6)
C5T	0.093 (7)	0.078 (7)	0.114 (9)	0.012 (6)	0.032 (7)	-0.005 (6)
C6T	0.081 (6)	0.066 (6)	0.102 (8)	0.011 (5)	0.016 (5)	0.016 (5)

Geometric parameters (Å, °)

Zn1A—N4A	1.920 (4)	N2B—C6B	1.456 (6)	
Zn1A—N2A	1.924 (4)	N3B—C20B	1.330 (7)	
Zn1A—N3A	2.127 (4)	N3B—C16B	1.338 (7)	
Zn1A—N1A	2.143 (4)	N4B—C21B	1.458 (7)	
Si1A—N2A	1.709 (5)	C1B—C2B	1.371 (8)	
Si1A—C13A	1.891 (7)	C1B—H1BA	0.9500	
Si1A—C7A	1.902 (6)	C2B—C3B	1.402 (8)	
Si1A—C10A	1.911 (7)	C2B—H2BA	0.9500	
Si2A—N4A	1.708 (4)	C3B—C4B	1.382 (8)	
Si2A—C25A	1.904 (6)	СЗВ—НЗВА	0.9500	
Si2A—C28A	1.906 (6)	C4B—C5B	1.385 (7)	
Si2A—C22A	1.926 (6)	C4B—H4BA	0.9500	
N1A—C5A	1.341 (7)	C5B—C6B	1.503 (7)	
N1A—C1A	1.352 (7)	C6B—H6BA	0.9900	
N2A—C6A	1.438 (7)	C6B—H6BB	0.9900	
N3A—C16A	1.317 (7)	C7B—C8B	1.541 (9)	
N3A—C20A	1.335 (7)	C7B—C9B	1.547 (8)	

N4A—C21A	1 449 (7)	C7B—H7BA	1 0000
C1A - C2A	1 357 (8)	C8B—H8BA	0.9800
C1A—H1AA	0.9500	C8B—H8BB	0.9800
$C^2A - C^3A$	1 379 (8)	C8B—H8BC	0.9800
C_{2A} H_{2AA}	0.9500	COB HOBA	0.9000
$C_{2A} = C_{4A}$	1 202 (8)		0.9800
$C_{2A} = U_{2A}$	1.392 (0)		0.9600
	0.9300	CIOD CIOD	0.9600
C4A—C5A	1.387 (8)	CIOB—CI2B	1.539 (9)
C4A—H4AA	0.9500	CIOB—CIIB	1.574 (9)
С5А—С6А	1.508 (8)	CI0B—HI0B	1.0000
С6А—Н6АА	0.9900	C11B—H11D	0.9800
С6А—Н6АВ	0.9900	C11B—H11E	0.9800
C7A—C9A	1.528 (9)	C11B—H11F	0.9800
C7A—C8A	1.539 (9)	C12B—H12D	0.9800
С7А—Н7АА	1.0000	C12B—H12E	0.9800
C8A—H8AA	0.9800	C12B—H12F	0.9800
C8A—H8AB	0.9800	C13B—C15B	1.511 (9)
C8A—H8AC	0.9800	C13B—C14B	1.556 (8)
С9А—Н9АА	0.9800	C13B—H13B	1.0000
С9А—Н9АВ	0.9800	C14B—H14D	0.9800
С9А—Н9АС	0.9800	C14B—H14E	0.9800
C10A—C11A	1.522 (10)	C14B—H14F	0.9800
C10A—C12A	1 525 (10)	C15B—H15D	0 9800
C10A - H10A	1 0000	C15B—H15E	0.9800
	0.9800	C15B—H15F	0.9800
C11A H11B	0.9800	C16B C17B	1 368 (0)
	0.9800	C_{16} C	0.0500
	0.9800	C17D $C19D$	1 262 (0)
CI2A—III2A	0.9800	C17D U17D	1.302 (9)
	0.9800		0.9300
CI2A—HI2C	0.9800	CI8B—CI9B	1.351 (8)
CI3A—CI5A	1.537 (11)	CI8B—HI8B	0.9500
CI3A—CI4A	1.563 (11)	C19B—C20B	1.396 (8)
С13А—Н13А	1.0000	C19B—H19B	0.9500
C14A—H14A	0.9800	C20B—C21B	1.504 (8)
C14A—H14B	0.9800	C21B—H21C	0.9900
C14A—H14C	0.9800	C21B—H21D	0.9900
C15A—H15A	0.9800	C22B—C23B	1.543 (8)
C15A—H15B	0.9800	C22B—C24B	1.549 (8)
C15A—H15C	0.9800	C22B—H22B	1.0000
C16A—C17A	1.406 (9)	C23B—H23D	0.9800
C16A—H16A	0.9500	С23В—Н23Е	0.9800
C17A—C18A	1.384 (9)	C23B—H23F	0.9800
C17A—H17A	0.9500	C24B—H24D	0.9800
C18A—C19A	1.364 (9)	C24B—H24E	0.9800
C18A—H18A	0.9500	C24B—H24F	0.9800
C19A—C20A	1.419 (8)	C25B—C26B	1.526 (9)
С19А—Н19А	0.9500	C25B—C27B	1.530 (8)
C20A—C21A	1.516 (8)	C25B—H25B	1.0000
	\-/		

C21A—H21A	0.9900	C26B—H26D	0.9800
C21A—H21B	0.9900	С26В—Н26Е	0.9800
C22A—C23A	1.502 (9)	C26B—H26F	0.9800
C22A—C24A	1.540 (9)	C27B—H27D	0.9800
C22A—H22A	1.0000	С27В—Н27Е	0.9800
C23A—H23A	0.9800	C27B—H27F	0.9800
C23A—H23B	0.9800	C28B—C29B	1.529 (9)
$C_{23}A - H_{23}C$	0.9800	C28B—C30B	1.548 (8)
C_{24A} H24A	0.9800	C28B—H28B	1 0000
C24A - H24B	0.9800	C29B—H29D	0.9800
C^{24A} H ^{24C}	0.9800	C_{29B} H29E	0.9800
C_{25A} C_{26A}	1 539 (8)	C_{29B} H29E	0.9800
C_{25A} C_{27A}	1 540 (9)	C_{30B} H30D	0.9800
$C_{25A} = C_{27A}$	1.0000	C30B—H30E	0.9800
C26A H26A	0.0800	C30B H30E	0.9800
C_{20A} H_{20A}	0.9800		1.361(14)
$C_{20}A = H_{20}C_{20}$	0.9800		1.301(14) 1.272(15)
$C_{20}A = H_{20}C$	0.9800	C1T = C2T	1.373(13)
$C_2/A = H_2/A$	0.9800		1.410(17) 1.222(15)
$C_2/A = H_2/B$	0.9800		1.332 (13)
$C_2/A = H_2/C$	0.9800	C2T—H2TC	0.9500
C28A—C30A	1.532 (8)		1.352 (14)
C28A—C29A	1.537 (8)	C31—H31C	0.9500
C28A—H28A	1.0000	C4T—C5T	1.356 (14)
С29А—Н29А	0.9800	C4T—H4TC	0.9500
C29A—H29B	0.9800	С5Т—С6Т	1.386 (14)
C29A—H29C	0.9800	C5T—H5TA	0.9500
C30A—H30A	0.9800	С6Т—Н6ТА	0.9500
C30A—H30B	0.9800	С7Т—Н7ТА	0.9800
C30A—H30C	0.9800	C7T—H7TB	0.9800
Zn1B—N4B	1.916 (4)	C7T—H7TC	0.9800
Zn1B—N2B	1.930 (4)	O1TH—C4TH	1.570 (18)
Zn1B—N1B	2.126 (4)	O1TH—C1TH	1.67 (2)
Zn1B—N3B	2.133 (4)	C1TH—C2TH	1.47 (2)
Si1B—N2B	1.711 (4)	C1TH—H1TA	0.9900
Si1B—C10B	1.893 (6)	C1TH—H1TB	0.9900
Si1B—C13B	1.903 (6)	С2ТН—С3ТН	1.528 (17)
Si1B—C7B	1.911 (6)	С2ТН—Н2ТА	0.9900
Si2B—N4B	1.716 (5)	C2TH—H2TB	0.9900
Si2B—C28B	1.889 (5)	C3TH—C4TH	1.499 (18)
Si2B—C22B	1.898 (6)	СЗТН—НЗТА	0.9900
Si2B—C25B	1.914 (6)	СЗТН—НЗТВ	0.9900
N1B—C1B	1.343 (7)	C4TH—H4TA	0.9900
N1B—C5B	1.348 (6)	C4TH—H4TB	0.9900
	- (-)		
N4A—Zn1A—N2A	145.65 (19)	C20B—N3B—Zn1B	110.4 (3)
N4A - Zn1A - N3A	84.49 (17)	C16B - N3B - Zn1B	131.5 (4)
N2A— $Zn1A$ — $N3A$	117.97 (18)	C21B - N4B - Si2B	116.4 (4)
N4A - Zn1A - N1A	119.56 (18)	$C_{21B} M_{4B} T_{n1B}$	114.3 (3)

N2A—Zn1A—N1A	83.91 (17)	Si2B—N4B—Zn1B	129.3 (2)
N3A—Zn1A—N1A	99.84 (16)	N1B—C1B—C2B	123.1 (5)
N2A—Si1A—C13A	107.1 (3)	N1B—C1B—H1BA	118.4
N2A—Si1A—C7A	109.6 (2)	C2B—C1B—H1BA	118.4
C13A—Si1A—C7A	114.0 (3)	C1B—C2B—C3B	118.3 (5)
N2A—Si1A—C10A	112.8 (3)	C1B—C2B—H2BA	120.8
C13A—Si1A—C10A	107.6 (3)	C3B—C2B—H2BA	120.8
C7A—Si1A—C10A	105.9 (3)	C4B—C3B—C2B	118.4 (5)
N4A—Si2A—C25A	115.0 (2)	С4В—С3В—Н3ВА	120.8
N4A—Si2A—C28A	105.5 (2)	С2В—С3В—Н3ВА	120.8
C25A—Si2A—C28A	107.6 (3)	C3B—C4B—C5B	120.3 (5)
N4A—Si2A—C22A	110.8 (3)	C3B—C4B—H4BA	119.9
C25A—Si2A—C22A	104.7 (3)	C5B—C4B—H4BA	119.9
C28A—Si2A—C22A	113.5 (3)	N1B—C5B—C4B	120.8 (5)
C5A—N1A—C1A	118.9 (5)	N1B—C5B—C6B	117.4 (4)
C5A—N1A—Zn1A	109.9 (4)	C4B—C5B—C6B	121.8 (5)
C1A—N1A—Zn1A	131.2 (4)	N2B—C6B—C5B	113.3 (4)
C6A—N2A—Si1A	118.1 (4)	N2B—C6B—H6BA	108.9
C6A—N2A—Zn1A	114.2 (3)	C5B—C6B—H6BA	108.9
Si1A—N2A—Zn1A	127.7 (2)	N2B—C6B—H6BB	108.9
C16A—N3A—C20A	120.2 (5)	C5B—C6B—H6BB	108.9
C16A—N3A—Zn1A	130.4 (4)	H6BA—C6B—H6BB	107.7
C20A—N3A—Zn1A	109.4 (3)	C8B—C7B—C9B	109.6 (5)
C21A—N4A—Si2A	118.3 (3)	C8B—C7B—Si1B	114.2 (4)
C21A—N4A—Zn1A	114.4 (3)	C9B—C7B—Si1B	113.6 (4)
Si2A—N4A—Zn1A	127.3 (2)	C8B—C7B—H7BA	106.2
N1A—C1A—C2A	122.4 (6)	C9B—C7B—H7BA	106.2
N1A—C1A—H1AA	118.8	Si1B—C7B—H7BA	106.2
C2A—C1A—H1AA	118.8	C7B—C8B—H8BA	109.5
C1A—C2A—C3A	119.5 (5)	C7B—C8B—H8BB	109.5
C1A—C2A—H2AA	120.3	H8BA—C8B—H8BB	109.5
C3A—C2A—H2AA	120.3	C7B—C8B—H8BC	109.5
C2A—C3A—C4A	118.9 (5)	H8BA—C8B—H8BC	109.5
С2А—С3А—НЗАА	120.5	H8BB—C8B—H8BC	109.5
С4А—С3А—НЗАА	120.5	С7В—С9В—Н9ВА	109.5
C5A—C4A—C3A	118.8 (5)	C7B—C9B—H9BB	109.5
C5A—C4A—H4AA	120.6	H9BA—C9B—H9BB	109.5
СЗА—С4А—Н4АА	120.6	С7В—С9В—Н9ВС	109.5
N1A—C5A—C4A	121.5 (5)	Н9ВА—С9В—Н9ВС	109.5
N1A—C5A—C6A	117.0 (5)	H9BB—C9B—H9BC	109.5
C4A—C5A—C6A	121.5 (5)	C12B—C10B—C11B	108.4 (6)
N2A—C6A—C5A	114.8 (5)	C12B—C10B—Si1B	112.4 (4)
N2A—C6A—H6AA	108.6	C11B—C10B—Si1B	114.0 (4)
С5А—С6А—Н6АА	108.6	C12B—C10B—H10B	107.3
N2A—C6A—H6AB	108.6	C11B—C10B—H10B	107.3
С5А—С6А—Н6АВ	108.6	Si1B—C10B—H10B	107.3
Н6АА—С6А—Н6АВ	107.5	C10B—C11B—H11D	109.5
C9A—C7A—C8A	111.3 (6)	C10B—C11B—H11E	109.5

C9A—C7A—Si1A	113.6 (4)	H11D—C11B—H11E	109.5
C8A—C7A—Si1A	114.2 (4)	C10B—C11B—H11F	109.5
С9А—С7А—Н7АА	105.6	H11D—C11B—H11F	109.5
С8А—С7А—Н7АА	105.6	H11E—C11B—H11F	109.5
Si1A—C7A—H7AA	105.6	C10B—C12B—H12D	109.5
С7А—С8А—Н8АА	109.5	C10B—C12B—H12E	109.5
C7A—C8A—H8AB	109.5	H12D—C12B—H12E	109.5
Н8АА—С8А—Н8АВ	109.5	C10B—C12B—H12F	109.5
C7A—C8A—H8AC	109.5	H12D— $C12B$ — $H12F$	109.5
H8AA—C8A—H8AC	109.5	H12E— $C12B$ — $H12F$	109.5
H8AB—C8A—H8AC	109.5	C15B—C13B—C14B	109.9 (6)
С7А—С9А—Н9АА	109.5	C15B—C13B—Si1B	113.6 (4)
C7A—C9A—H9AB	109.5	C14B— $C13B$ — $Si1B$	115.6 (4)
Н9АА—С9А—Н9АВ	109.5	C15B—C13B—H13B	105.6
C7A—C9A—H9AC	109.5	C14B—C13B—H13B	105.6
Н9АА—С9А—Н9АС	109.5	Si1B—C13B—H13B	105.6
H9AB—C9A—H9AC	109.5	C13B— $C14B$ — $H14D$	109 5
C11A - C10A - C12A	109.7 (7)	C13B— $C14B$ — $H14E$	109.5
C11A - C10A - Si1A	111.6(5)	H_{14D} $-C_{14B}$ $-H_{14E}$	109.5
C12A - C10A - Si1A	114 9 (5)	C13B— $C14B$ — $H14F$	109.5
C11A - C10A - H10A	106.7	H14D— $C14B$ — $H14F$	109.5
C12A - C10A - H10A	106.7	H14E— $C14B$ — $H14F$	109.5
SilA—Cl0A—H10A	106.7	C13B— $C15B$ — $H15D$	109.5
C10A - C11A - H11A	109.5	C13B— $C15B$ — $H15E$	109.5
C10A - C11A - H11B	109.5	H15D—C15B—H15E	109.5
H11A—C11A—H11B	109.5	C13B— $C15B$ — $H15F$	109.5
C10A - C11A - H11C	109.5	H15D— $C15B$ — $H15F$	109.5
H11A—C11A—H11C	109.5	H15E— $C15B$ — $H15F$	109.5
H11B—C11A—H11C	109.5	N3B-C16B-C17B	123.9 (5)
C10A - C12A - H12A	109.5	N3B-C16B-H16B	118.1
C10A - C12A - H12B	109.5	C17B-C16B-H16B	118.1
H12A— $C12A$ — $H12B$	109.5	C18B— $C17B$ — $C16B$	117.6 (6)
C10A - C12A - H12C	109.5	C18B— $C17B$ — $H17B$	121.2
H12A— $C12A$ — $H12C$	109.5	C16B - C17B - H17B	121.2
H12B— $C12A$ — $H12C$	109.5	C19B— $C18B$ — $C17B$	119.9 (6)
C15A - C13A - C14A	107.5 (6)	C19B—C18B—H18B	120.1
C15A - C13A - Si1A	115.9 (5)	C17B $C18B$ $H18B$	120.1
C14A - C13A - Si1A	114.8 (5)	C18B— $C19B$ — $C20B$	120.1 120.0(5)
C15A - C13A - H13A	105.9	C18B— $C19B$ — $H19B$	120.0 (3)
C14A - C13A - H13A	105.9	C_{20B} C_{19B} H_{19B}	120.0
SilA—Cl3A—Hl3A	105.9	N3B-C20B-C19B	120.0 120.5(5)
C13A - C14A - H14A	109.5	N3B C20B C17B	120.5(5) 1174(5)
C13A - C14A - H14B	109.5	C19B - C20B - C21B	1221(5)
H14A— $C14A$ — $H14B$	109 5	N4B-C21B-C20B	1137(5)
C13A - C14A - H14C	109 5	N4B-C21B-H21C	108.8
H_{14A} C_{14A} H_{14C}	109.5	C20B-C21B-H21C	108.8
H14B— $C14A$ — $H14C$	109.5	N4B-C21B-H21D	108.8
C13A - C15A - H15A	109.5	C20B-C21B-H21D	108.8

C13A—C15A—H15B	109.5	H21C—C21B—H21D	107.7
H15A—C15A—H15B	109.5	C23B—C22B—C24B	107.5 (5)
C13A—C15A—H15C	109.5	C23B—C22B—Si2B	115.1 (4)
H15A—C15A—H15C	109.5	C24B—C22B—Si2B	112.6 (4)
H15B—C15A—H15C	109.5	C23B—C22B—H22B	107.1
N3A—C16A—C17A	122.9 (6)	C24B—C22B—H22B	107.1
N3A—C16A—H16A	118.6	Si2B—C22B—H22B	107.1
C17A—C16A—H16A	118.6	C22B—C23B—H23D	109.5
C18A—C17A—C16A	117.3 (6)	C22B—C23B—H23E	109.5
C18A—C17A—H17A	121.4	H23D—C23B—H23E	109.5
С16А—С17А—Н17А	121.4	C22B—C23B—H23F	109.5
C19A—C18A—C17A	120.1 (6)	H23D—C23B—H23F	109.5
C19A—C18A—H18A	119.9	H23E—C23B—H23F	109.5
C17A—C18A—H18A	119.9	C22B—C24B—H24D	109.5
C18A—C19A—C20A	119.2 (5)	C22B—C24B—H24E	109.5
C18A - C19A - H19A	120.4	H_24D — C_24B — H_24E	109.5
C_{20A} C_{19A} H_{19A}	120.4	C22B— $C24B$ — $H24F$	109.5
N3A—C20A—C19A	120.3 (5)	H_24D — C_24B — H_24F	109.5
N3A - C20A - C21A	1184(5)	H24E— $C24B$ — $H24F$	109.5
C19A - C20A - C21A	121.2 (5)	C_{26B} C_{25B} C_{27B}	109.0 111.8(5)
N4A - C21A - C20A	1130(4)	$C_{26B} = C_{25B} = S_{12B}$	111.8(2)
N4A—C21A—H21A	109.0	C27B $C25B$ $Si2B$	1181(4)
C_{20A} C_{21A} H_{21A}	109.0	$C_{26B} = C_{25B} = H_{25B}$	104.6
N4A—C21A—H21B	109.0	C_{27B} C_{25B} H_{25B}	104.6
C_{20A} C_{21A} H_{21B}	109.0	Si2B—C25B—H25B	104.6
H_{21A} C_{21A} H_{21B}	107.8	$C_{25B} - C_{26B} - H_{26D}$	109.5
C23A—C22A—C24A	111.9 (5)	C25B—C26B—H26E	109.5
C23A—C22A—Si2A	114.1 (5)	H26D—C26B—H26E	109.5
C24A— $C22A$ — $Si2A$	113.6 (4)	C25B—C26B—H26F	109.5
C23A—C22A—H22A	105.4	H26D—C26B—H26F	109.5
C24A—C22A—H22A	105.4	H26E—C26B—H26F	109.5
Si2A—C22A—H22A	105.4	C25B—C27B—H27D	109.5
C22A—C23A—H23A	109.5	C25B—C27B—H27E	109.5
C22A—C23A—H23B	109.5	H27D—C27B—H27E	109.5
H23A—C23A—H23B	109.5	C25B—C27B—H27F	109.5
С22А—С23А—Н23С	109.5	H27D—C27B—H27F	109.5
H23A—C23A—H23C	109.5	H27E—C27B—H27F	109.5
H23B—C23A—H23C	109.5	C29B—C28B—C30B	109.6 (5)
C22A—C24A—H24A	109.5	C29B—C28B—Si2B	115.8 (4)
C22A—C24A—H24B	109.5	C30B—C28B—Si2B	114.5 (4)
H24A—C24A—H24B	109.5	C29B—C28B—H28B	105.3
C22A—C24A—H24C	109.5	C30B—C28B—H28B	105.3
H24A—C24A—H24C	109.5	Si2B—C28B—H28B	105.3
H24B—C24A—H24C	109.5	C28B—C29B—H29D	109.5
C26A—C25A—C27A	106.7 (5)	C28B—C29B—H29E	109.5
C26A—C25A—Si2A	114.5 (4)	H29D—C29B—H29E	109.5
C27A—C25A—Si2A	111.3 (4)	C28B—C29B—H29F	109.5
C26A—C25A—H25A	108.0	H29D—C29B—H29F	109.5

	4.0.0.0		
C27A—C25A—H25A	108.0	H29E—C29B—H29F	109.5
Si2A—C25A—H25A	108.0	C28B—C30B—H30D	109.5
C25A—C26A—H26A	109.5	C28B—C30B—H30E	109.5
C25A—C26A—H26B	109.5	H30D-C30B-H30E	109.5
H26A—C26A—H26B	109.5	C28B—C30B—H30F	109.5
C25A—C26A—H26C	109.5	H30D-C30B-H30F	109.5
H26A—C26A—H26C	109.5	H30E—C30B—H30F	109.5
H26B—C26A—H26C	109.5	C6T—C1T—C2T	117.2 (10)
С25А—С27А—Н27А	109.5	C6T—C1T—C7T	119.8 (13)
C25A—C27A—H27B	109.5	C2T-C1T-C7T	123.0(12)
H27A—C27A—H27B	109.5	C3T - C2T - C1T	120.9(11)
C_{25A} C_{27A} H_{27C}	109.5	C3T - C2T - H2TC	119.5
$H_{27A} = C_{27A} = H_{27C}$	109.5	CIT C2T H2TC	119.5
$\frac{1127A}{27A} = \frac{127C}{1127C}$	109.5	$C_{11} = C_{21} = H_{21}C$	119.5 122.3(11)
$\Pi Z/D = CZ/A = \Pi Z/C$	109.5	$C_2T = C_3T = C_4T$	122.3 (11)
$C_{20A} = C_{28A} = C_{29A}$	109.4 (3)		110.0
C30A - C28A - S12A	112.8 (4)	C41 - C31 - H31C	118.8
C29A—C28A—S12A	116.1 (4)	C31—C41—C51	118.5 (11)
C30A—C28A—H28A	105.9	C31—C41—H41C	120.7
C29A—C28A—H28A	105.9	C5T—C4T—H4TC	120.7
Si2A—C28A—H28A	105.9	C4T—C5T—C6T	119.3 (10)
С28А—С29А—Н29А	109.5	C4T—C5T—H5TA	120.3
C28A—C29A—H29B	109.5	C6T—C5T—H5TA	120.3
H29A—C29A—H29B	109.5	C1T—C6T—C5T	121.5 (11)
C28A—C29A—H29C	109.5	С1Т—С6Т—Н6ТА	119.3
H29A—C29A—H29C	109.5	С5Т—С6Т—Н6ТА	119.3
H29B—C29A—H29C	109.5	С1Т—С7Т—Н7ТА	109.5
C28A—C30A—H30A	109.5	C1T—C7T—H7TB	109.5
C28A—C30A—H30B	109.5	Н7ТА—С7Т—Н7ТВ	109.5
H30A—C30A—H30B	109.5	C1T—C7T—H7TC	109.5
C28A—C30A—H30C	109.5	H7TA—C7T—H7TC	109.5
H30A—C30A—H30C	109.5	Н7ТВ—С7Т—Н7ТС	109.5
H30B—C30A—H30C	109.5	C4TH—O1TH—C1TH	107.6 (11)
N4B—Zn1B—N2B	143.02 (19)	C2TH—C1TH—O1TH	1052(12)
N4B = 7n1B = N1B	120.98(18)	C2TH_C1TH_H1TA	110.7
$N2B_7n1B_N1B$	83 74 (16)		110.7
N/B Zn1B N3B	83.69 (17)	C2TH C1TH H1TB	110.7
N2P = 7n1P = N2P	122 00 (18)		110.7
N1D 7-1D N2D	122.00(18)		100.7
NID—ZIIID—IN3D	96.75 (17)		108.8
N2B—SIIB—CI0B	114.9 (2)	CITH-C2IH-C3IH	109.7 (11)
N2B—S11B—C13B	105.8 (2)	CIIH—C2IH—H2IA	109.7
Clob—SilB—Cl3B	107.3 (3)	C31H—C21H—H21A	109.7
N2B—S11B—C7B	110.1 (2)	C1TH—C2TH—H2TB	109.7
C10B—Si1B—C7B	105.1 (3)	C3TH—C2TH—H2TB	109.7
C13B—Si1B—C7B	113.8 (3)	Н2ТА—С2ТН—Н2ТВ	108.2
N4B—Si2B—C28B	106.5 (2)	C4TH—C3TH—C2TH	113.1 (11)
N4B—Si2B—C22B	110.9 (2)	С4ТН—С3ТН—Н3ТА	109.0
C28B—Si2B—C22B	113.4 (3)	С2ТН—С3ТН—Н3ТА	109.0
N4B—Si2B—C25B	108.9 (2)	С4ТН—С3ТН—Н3ТВ	109.0

C28B—Si2B—C25B	110.4 (3)	C2TH—C3TH—H3TB	109.0	
C22B—Si2B—C25B	106.7 (3)	НЗТА—СЗТН—НЗТВ	107.8	
C1B—N1B—C5B	119.1 (5)	C3TH—C4TH—O1TH	104.5 (11)	
C1B—N1B—Zn1B	130.8 (4)	C3TH—C4TH—H4TA	110.9	
C5B—N1B—Zn1B	110.1 (3)	O1TH—C4TH—H4TA	110.9	
C6B—N2B—Si1B	119.2 (3)	C3TH—C4TH—H4TB	110.9	
C6B—N2B—Zn1B	113.9 (3)	O1TH—C4TH—H4TB	110.9	
Si1B—N2B—Zn1B	126.9 (2)	H4TA—C4TH—H4TB	108.9	
C20B—N3B—C16B	118.1 (5)			