

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## Ammonium 1-hydroxy-2-naphthoate

Ye Bi and Cheng-Li Han\*

College of Chemistry and Chemical Engineering, Qiqihar University, Qiqihar 161006, People's Republic of China

Correspondence e-mail: chengli\_han@126.com


Received 18 March 2008; accepted 3 April 2008

 Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.080;  $wR$  factor = 0.227; data-to-parameter ratio = 12.9.

The title compound,  $\text{NH}_4^+\cdot\text{C}_{11}\text{H}_7\text{O}_3^-$ , was obtained by slow evaporation of a 30% ammonia solution of 1-hydroxy-2-naphthoic acid. The crystal structure is stabilized by intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds, forming layers parallel to the  $bc$  plane.

## Related literature

For related literature, see: Kickelbick & Schubert (1999); Ohki *et al.* (1986); Song *et al.* (2008).



## Experimental

## Crystal data

 $\text{NH}_4^+\cdot\text{C}_{11}\text{H}_7\text{O}_3^-$   
 $M_r = 205.21$ 

 Monoclinic,  $C2/c$   
 $a = 30.883$  (5) Å  
 $b = 3.880$  (1) Å  
 $c = 15.777$  (3) Å  
 $\beta = 95.567$  (2)°

 $V = 1881.6$  (7) Å<sup>3</sup>  
 $Z = 8$ 

 Mo  $K\alpha$  radiation  
 $\mu = 0.11$  mm<sup>-1</sup>  
 $T = 298$  (2) K  
 $0.23 \times 0.23 \times 0.20$  mm

## Data collection

 Bruker SMART CCD area-detector diffractometer  
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.976$ ,  $T_{\max} = 0.979$ 

 6728 measured reflections  
 1915 independent reflections  
 1351 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.040$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.080$   
 $wR(F^2) = 0.226$   
 $S = 1.04$   
 1915 reflections  
 149 parameters  
 10 restraints

 H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.55$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$                                      | $D-H$    | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|--|----------|-------------|-------------|---------------|
| $\text{O1}-\text{H1}\cdots\text{O2}$               | 0.82     | 1.73        | 2.463 (3)   | 148           |
| $\text{N1}-\text{H1A}\cdots\text{O1}^{\text{i}}$   | 0.89 (2) | 2.07 (3)    | 2.920 (3)   | 161 (3)       |
| $\text{N1}-\text{H1B}\cdots\text{O2}^{\text{ii}}$  | 0.89 (3) | 1.88 (3)    | 2.756 (3)   | 167 (3)       |
| $\text{N1}-\text{H1C}\cdots\text{O3}^{\text{iii}}$ | 0.89 (2) | 2.04 (2)    | 2.789 (3)   | 141 (3)       |
| $\text{N1}-\text{H1D}\cdots\text{O3}^{\text{iv}}$  | 0.88 (3) | 2.08 (2)    | 2.821 (3)   | 140 (3)       |

Symmetry codes: (i)  $x, -y + 1, z - \frac{1}{2}$ ; (ii)  $-x + 1, -y + 2, -z + 1$ ; (iii)  $x, y - 1, z - 1$ ; (iv)  $x, y, z - 1$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors acknowledge Qiqihar University for a research grant.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2202).

## References

- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Kickelbick, G. & Schubert, U. (1999). *J. Chem. Soc. Dalton Trans.* pp. 1301–1306.  
 Ohki, Y., Suzuki, Y., Takeuchi, T., Shimoi, M. & Ouchi, A. (1986). *Bull. Chem. Soc. Jpn.*, **59**, 1015–1019.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Song, W.-D., Yan, J.-B., Wang, H. & Ji, L.-L. (2008). *Acta Cryst.* **E64**, m5.

**supplementary materials**

*Acta Cryst.* (2008). E64, o813 [ doi:10.1107/S1600536808009082 ]

## Ammonium 1-hydroxy-2-naphthoate

Y. Bi and C.-L. Han

### Comment

1-Hydroxynaphthalene-2-carboxylic acid is a widely used ligand for the synthesis of metal complexes (Kickelbick & Schubert, 1999; Ohki *et al.*, 1986; Song *et al.*, 2008). We report herein the crystal structure of the title compound, which was obtained by slow evaporation of a 30% ammonium solution of 1-hydroxynaphthalene-2-carboxylic acid in air.

The compound consists of discrete 1-hydroxynaphthalene-2-carboxylate anions and ammonium cations (Fig. 1). The anion is substantially planar with a mean deviation of 0.015 (3) Å. The crystal structure is stabilized by intermolecular N–H···O hydrogen bonds (Table 1), forming layers parallel to the *bc* plane (Fig. 2).

### Experimental

Single crystals of the title compound were obtained by slow evaporation of a 30% ammonia solution of 1-hydroxynaphthalene-2-carboxylic acid in air.

### Refinement

Ammonium H atoms were located from a difference Fourier map and refined isotropically, with N–H distances restrained to 0.90 (1) Å, H···H distances restrained to 1.43 (2) Å, and with  $U_{\text{iso}}(\text{H})$  values fixed at 0.08 Å<sup>2</sup>. All other H atoms were placed in idealized positions and constrained to ride on their parent atoms with C–H distances of 0.93 Å, O–H distance of 0.82 Å, and with  $U_{\text{iso}}(\text{H})$  set at 1.2 $U_{\text{eq}}(\text{C})$  or 1.5 $U_{\text{eq}}(\text{O})$ .

### Figures




Fig. 1. The structure of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme. The intramolecular hydrogen bond is shown as a dashed line.





Fig. 2. A perspective view of crystal packing of the title compound, viewed along the *a* axis. Hydrogen bonds are shown as dashed lines.

## Ammonium 1-hydroxy-2-naphthoate

### Crystal data



$$M_r = 205.21$$

Monoclinic,  $C2/c$

Hall symbol:  $-C\ 2yc$

$$a = 30.883\ (5)\ \text{\AA}$$

$$b = 3.880\ (1)\ \text{\AA}$$

$$c = 15.777\ (3)\ \text{\AA}$$

$$\beta = 95.567\ (2)^\circ$$

$$V = 1881.6\ (7)\ \text{\AA}^3$$

$$Z = 8$$

$$F_{000} = 864$$

$$D_x = 1.449\ \text{Mg m}^{-3}$$

Mo  $K\alpha$  radiation

$$\lambda = 0.71073\ \text{\AA}$$

Cell parameters from 1379 reflections

$$\theta = 2.5\text{--}24.1^\circ$$

$$\mu = 0.11\ \text{mm}^{-1}$$

$$T = 298\ (2)\ \text{K}$$

Block, colourless

$$0.23 \times 0.23 \times 0.20\ \text{mm}$$

### Data collection

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$$T = 298(2)\ \text{K}$$

$\omega$  scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)

$$T_{\min} = 0.976, T_{\max} = 0.979$$

6728 measured reflections

1915 independent reflections

1351 reflections with  $I > 2\sigma(I)$

$$R_{\text{int}} = 0.040$$

$$\theta_{\max} = 27.0^\circ$$

$$\theta_{\min} = 2.6^\circ$$

$$h = -38 \rightarrow 38$$

$$k = -4 \rightarrow 4$$

$$l = -20 \rightarrow 19$$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.080$$

$$wR(F^2) = 0.226$$

$$S = 1.04$$

1915 reflections

149 parameters

10 restraints

Primary atom site location: structure-invariant direct  
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring  
sites

H atoms treated by a mixture of  
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.1461P)^2 + 0.0944P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.55\ \text{e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.24\ \text{e \AA}^{-3}$$

Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

|     | <i>x</i>     | <i>y</i>   | <i>z</i>     | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|------------|--------------|----------------------------------|
| O1  | 0.42976 (6)  | 1.0366 (6) | 0.70025 (11) | 0.0383 (6)                       |
| H1  | 0.4496       | 1.0981     | 0.7354       | 0.057*                           |
| O2  | 0.46982 (6)  | 1.1294 (6) | 0.84111 (13) | 0.0472 (6)                       |
| O3  | 0.44313 (7)  | 0.9570 (6) | 0.95912 (12) | 0.0486 (7)                       |
| N1  | 0.46529 (8)  | 0.4511 (7) | 0.08424 (16) | 0.0414 (7)                       |
| C1  | 0.39678 (8)  | 0.9092 (6) | 0.74083 (16) | 0.0259 (6)                       |
| C2  | 0.39995 (8)  | 0.8798 (7) | 0.82806 (16) | 0.0281 (6)                       |
| C3  | 0.36469 (8)  | 0.7397 (7) | 0.86694 (16) | 0.0322 (6)                       |
| H3  | 0.3666       | 0.7214     | 0.9260       | 0.039*                           |
| C4  | 0.32803 (9)  | 0.6309 (8) | 0.82012 (17) | 0.0354 (7)                       |
| H4  | 0.3052       | 0.5392     | 0.8474       | 0.042*                           |
| C5  | 0.32420 (8)  | 0.6559 (7) | 0.73053 (17) | 0.0291 (6)                       |
| C6  | 0.28703 (9)  | 0.5425 (7) | 0.67931 (19) | 0.0380 (7)                       |
| H6  | 0.2640       | 0.4471     | 0.7050       | 0.046*                           |
| C7  | 0.28423 (9)  | 0.5700 (8) | 0.5933 (2)   | 0.0441 (8)                       |
| H7  | 0.2594       | 0.4931     | 0.5606       | 0.053*                           |
| C8  | 0.31858 (10) | 0.7139 (8) | 0.55356 (18) | 0.0424 (8)                       |
| H8  | 0.3165       | 0.7321     | 0.4945       | 0.051*                           |
| C9  | 0.35487 (9)  | 0.8268 (8) | 0.60050 (17) | 0.0358 (7)                       |
| H9  | 0.3773       | 0.9242     | 0.5733       | 0.043*                           |
| C10 | 0.35900 (8)  | 0.7985 (7) | 0.68970 (16) | 0.0268 (6)                       |
| C11 | 0.43967 (9)  | 0.9947 (7) | 0.88135 (17) | 0.0321 (7)                       |
| H1A | 0.4533 (9)   | 0.347 (7)  | 0.1262 (14)  | 0.080*                           |
| H1B | 0.4891 (7)   | 0.565 (8)  | 0.1041 (18)  | 0.080*                           |
| H1C | 0.4720 (10)  | 0.291 (6)  | 0.0471 (16)  | 0.080*                           |
| H1D | 0.4467 (8)   | 0.597 (7)  | 0.0576 (18)  | 0.080*                           |

*Atomic displacement parameters ( $\text{\AA}^2$ )*

|    | $U^{11}$    | $U^{22}$    | $U^{33}$    | $U^{12}$    | $U^{13}$    | $U^{23}$     |
|----|-------------|-------------|-------------|-------------|-------------|--------------|
| O1 | 0.0294 (10) | 0.0542 (14) | 0.0317 (10) | −0.0088 (9) | 0.0045 (8)  | 0.0017 (9)   |
| O2 | 0.0307 (10) | 0.0587 (15) | 0.0509 (13) | −0.0139 (9) | −0.0027 (9) | −0.0017 (11) |

## supplementary materials

---

|     |             |             |             |              |              |              |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| O3  | 0.0537 (13) | 0.0560 (15) | 0.0326 (11) | -0.0071 (10) | -0.0135 (10) | -0.0059 (10) |
| N1  | 0.0445 (14) | 0.0356 (14) | 0.0439 (14) | -0.0050 (11) | 0.0037 (12)  | -0.0057 (11) |
| C1  | 0.0240 (12) | 0.0254 (13) | 0.0293 (13) | 0.0014 (9)   | 0.0070 (10)  | -0.0002 (10) |
| C2  | 0.0273 (13) | 0.0253 (14) | 0.0310 (13) | 0.0001 (10)  | 0.0003 (10)  | -0.0035 (10) |
| C3  | 0.0371 (14) | 0.0330 (15) | 0.0269 (13) | -0.0017 (12) | 0.0057 (11)  | 0.0026 (11)  |
| C4  | 0.0329 (14) | 0.0391 (17) | 0.0352 (14) | -0.0060 (11) | 0.0085 (12)  | 0.0052 (12)  |
| C5  | 0.0246 (12) | 0.0264 (14) | 0.0359 (14) | 0.0022 (10)  | 0.0010 (10)  | 0.0016 (11)  |
| C6  | 0.0306 (14) | 0.0337 (16) | 0.0488 (16) | -0.0029 (11) | -0.0010 (12) | -0.0043 (13) |
| C7  | 0.0338 (15) | 0.0469 (19) | 0.0480 (17) | 0.0026 (12)  | -0.0146 (13) | -0.0136 (14) |
| C8  | 0.0440 (16) | 0.0515 (19) | 0.0289 (14) | 0.0107 (14)  | -0.0100 (12) | -0.0052 (13) |
| C9  | 0.0343 (14) | 0.0436 (17) | 0.0300 (14) | 0.0060 (12)  | 0.0049 (11)  | -0.0019 (12) |
| C10 | 0.0263 (12) | 0.0239 (13) | 0.0297 (13) | 0.0024 (10)  | 0.0000 (10)  | -0.0007 (10) |
| C11 | 0.0346 (14) | 0.0271 (14) | 0.0331 (14) | -0.0008 (11) | -0.0044 (11) | -0.0041 (11) |

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

|            |            |           |           |
|------------|------------|-----------|-----------|
| O1—C1      | 1.349 (3)  | C3—H3     | 0.9300    |
| O1—H1      | 0.8200     | C4—C5     | 1.410 (4) |
| O2—C11     | 1.287 (3)  | C4—H4     | 0.9300    |
| O3—C11     | 1.230 (3)  | C5—C6     | 1.409 (4) |
| N1—H1A     | 0.89 (2)   | C5—C10    | 1.418 (4) |
| N1—H1B     | 0.89 (3)   | C6—C7     | 1.356 (4) |
| N1—H1C     | 0.89 (2)   | C6—H6     | 0.9300    |
| N1—H1D     | 0.88 (3)   | C7—C8     | 1.400 (4) |
| C1—C2      | 1.375 (4)  | C7—H7     | 0.9300    |
| C1—C10     | 1.419 (3)  | C8—C9     | 1.355 (4) |
| C2—C3      | 1.410 (3)  | C8—H8     | 0.9300    |
| C2—C11     | 1.487 (3)  | C9—C10    | 1.405 (4) |
| C3—C4      | 1.358 (4)  | C9—H9     | 0.9300    |
| C1—O1—H1   | 109.5      | C6—C5—C10 | 118.2 (3) |
| H1A—N1—H1B | 110.8 (19) | C4—C5—C10 | 119.3 (2) |
| H1A—N1—H1C | 108 (2)    | C7—C6—C5  | 121.3 (3) |
| H1B—N1—H1C | 110.0 (19) | C7—C6—H6  | 119.4     |
| H1A—N1—H1D | 110 (2)    | C5—C6—H6  | 119.4     |
| H1B—N1—H1D | 109 (2)    | C6—C7—C8  | 120.1 (3) |
| H1C—N1—H1D | 108.3 (19) | C6—C7—H7  | 119.9     |
| O1—C1—C2   | 121.5 (2)  | C8—C7—H7  | 119.9     |
| O1—C1—C10  | 117.3 (2)  | C9—C8—C7  | 120.5 (3) |
| C2—C1—C10  | 121.2 (2)  | C9—C8—H8  | 119.8     |
| C1—C2—C3   | 119.0 (2)  | C7—C8—H8  | 119.8     |
| C1—C2—C11  | 121.1 (2)  | C8—C9—C10 | 120.8 (3) |
| C3—C2—C11  | 120.0 (2)  | C8—C9—H9  | 119.6     |
| C4—C3—C2   | 121.4 (2)  | C10—C9—H9 | 119.6     |
| C4—C3—H3   | 119.3      | C9—C10—C5 | 119.1 (2) |
| C2—C3—H3   | 119.3      | C9—C10—C1 | 122.4 (2) |
| C3—C4—C5   | 120.5 (2)  | C5—C10—C1 | 118.5 (2) |
| C3—C4—H4   | 119.7      | O3—C11—O2 | 122.9 (2) |
| C5—C4—H4   | 119.7      | O3—C11—C2 | 121.0 (3) |
| C6—C5—C4   | 122.5 (3)  | O2—C11—C2 | 116.1 (2) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H··· <i>A</i>    | <i>D</i> —H | H··· <i>A</i> | <i>D</i> ··· <i>A</i> | <i>D</i> —H··· <i>A</i> |
|----------------------------|-------------|---------------|-----------------------|-------------------------|
| O1—H1···O2                 | 0.82        | 1.73          | 2.463 (3)             | 148                     |
| N1—H1A···O1 <sup>i</sup>   | 0.89 (2)    | 2.07 (3)      | 2.920 (3)             | 161 (3)                 |
| N1—H1B···O2 <sup>ii</sup>  | 0.89 (3)    | 1.88 (3)      | 2.756 (3)             | 167 (3)                 |
| N1—H1C···O3 <sup>iii</sup> | 0.89 (2)    | 2.04 (2)      | 2.789 (3)             | 141 (3)                 |
| N1—H1D···O3 <sup>iv</sup>  | 0.88 (3)    | 2.08 (2)      | 2.821 (3)             | 140 (3)                 |

Symmetry codes: (i)  $x, -y+1, z-1/2$ ; (ii)  $-x+1, -y+2, -z+1$ ; (iii)  $x, y-1, z-1$ ; (iv)  $x, y, z-1$ .

Fig. 1




Fig. 2

