

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tetraaquabis(biuret- κ^2O,O')-gadolinium(III) trichloride

William T. A. Harrison

Department of Chemistry, University of Aberdeen, Meston Walk, Aberdeen AB24 3UE, Scotland
Correspondence e-mail: w.harrison@abdn.ac.uk


Received 27 March 2008; accepted 31 March 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(O-C) = 0.004$ Å; R factor = 0.034; wR factor = 0.088; data-to-parameter ratio = 31.0.

In the title compound, $[Gd(C_2H_5N_3O_2)_2(H_2O)_4]Cl_3$, which is isostructural with its yttrium analogue, the Gd^{3+} ion (site symmetry 2) is bonded to eight O atoms (arising from two O,O' -bidentate biuret molecules and four water molecules) in a distorted square-antiprismatic arrangement. A network of $N-H \cdots O$, $N-H \cdots Cl$ and $O-H \cdots Cl$ hydrogen bonds helps to establish the packing, leading to a three-dimensional network. One of the chloride ions has site symmetry 2.

Related literature

For related structures, see: Haddad (1987, 1988); Harrison (2008). For related literature, see: Bernstein *et al.* (1995). For valence-sum calculations, see: Brese & O'Keeffe (1991).



Experimental

Crystal data

$[Gd(C_2H_5N_3O_2)_2(H_2O)_4]Cl_3$
 $M_r = 541.84$
Monoclinic, $C2/c$
 $a = 7.6501$ (3) Å
 $b = 13.2164$ (5) Å
 $c = 17.4557$ (6) Å
 $\beta = 100.961$ (1)°

$V = 1732.69$ (11) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 4.33$ mm⁻¹
 $T = 293$ (2) K
0.47 × 0.34 × 0.06 mm

Data collection

Bruker SMART1000 CCD diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 1999)
 $T_{min} = 0.235$, $T_{max} = 0.781$

8649 measured reflections
3134 independent reflections
2902 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.088$
 $S = 1.06$
3134 reflections

101 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 2.99$ e Å⁻³
 $\Delta\rho_{min} = -3.45$ e Å⁻³

Table 1

Selected bond lengths (Å).

Gd1—O1	2.350 (2)	Gd1—O4	2.407 (3)
Gd1—O2	2.375 (2)	Gd1—O3	2.414 (2)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N1—H1 ⁱ ···O1 ⁱ	0.86	2.10	2.910 (4)	157
N1—H2 ⁱⁱ ···Cl1 ⁱⁱ	0.86	2.40	3.194 (3)	154
N2—H3 ⁱⁱⁱ ···Cl1 ⁱⁱⁱ	0.86	2.53	3.315 (3)	153
N3—H4 ⁱⁱⁱⁱ ···Cl1 ⁱⁱⁱⁱ	0.86	2.54	3.363 (3)	161
N3—H5 ^v ···Cl1 ^v	0.86	2.53	3.311 (3)	151
O3—H6 ^{vi} ···Cl1 ^{vi}	0.81	2.39	3.163 (2)	162
O3—H7 ^{vii} ···Cl2 ^{vii}	0.79	2.28	3.059 (2)	167
O4—H8 ^{viii} ···Cl1	0.76	2.45	3.208 (2)	178
O4—H9 ^{ix} ···Cl2	0.80	2.41	3.127 (2)	150

Symmetry codes: (i) $-x + 1, y, -z + \frac{3}{2}$; (ii) $x + \frac{1}{2}, y - \frac{1}{2}, z$; (iii) $x - \frac{1}{2}, y - \frac{1}{2}, z$; (iv) $-x, -y, -z + 1$; (v) $-x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (vi) $x - 1, y, z$.

Data collection: SMART (Bruker, 1999); cell refinement: SAINT (Bruker, 1999); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SG2230).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Brese, N. E. & O'Keeffe, M. (1991). *Acta Cryst.* **B47**, 192–197.
Bruker (1999). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Haddad, S. F. (1987). *Acta Cryst.* **C43**, 1882–1885.
Haddad, S. F. (1988). *Acta Cryst.* **C44**, 815–818.
Harrison, W. T. A. (2008). *Acta Cryst.* **E64**, m619.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m620 [doi:10.1107/S1600536808008660]

Tetraaquabis(biuret- κ^2O,O')gadolinium(III) trichloride

W. T. A. Harrison

Comment

The title compound, (I), is isostructural with its recently described yttrium-containing analogue (Harrison, 2008).

The complete $[\text{Gd}(\text{biur})_2(\text{H}_2\text{O})_4]^{3+}$ (biur = biuret, $\text{C}_2\text{H}_5\text{N}_3\text{O}_2$) complex ion in (I) is generated by crystallographic 2-fold symmetry, with the metal atom lying on the rotation axis. Two uncoordinated chloride ions, one of which has site symmetry 2, complete the structure (Fig. 1) of (I).

The resulting GdO_8 polyhedral geometry in (I) (Table 1) is a distorted square antiprism. The nominal square face formed by atoms O1, O2, O1ⁱ and O2ⁱ ($i = -x, y, 3/2 - z$) is reasonably regular, but the second face formed by the four water molecules (O3, O4, O3ⁱ and O4ⁱ) is much more distorted, and the diagonal O3 \cdots O3ⁱ O4 \cdots O4ⁱ distances of 4.322 (4) Å and 3.551 (4) Å, respectively, are very different. Gd1 deviates from the mean planes of O1/O2/O1ⁱ/O2ⁱ and O3/O4/O3ⁱ/O4ⁱ by 1.1542 (16) Å and 1.3501 (19) Å, respectively. The gadolinium bond valence sum of 2.90, calculated by the Brese & O'Keffe (1991) method, indicates that it is slightly underbonded in (I), whereas in $[\text{Y}(\text{biur})_2(\text{H}_2\text{O})_4]\cdot 3\text{Cl}$ (Harrison, 2008), the yttrium cation was distinctly overbonded with a BVS of 3.34 (expected value = 3.00 in both cases).

The dihedral angle between the N1/C1/O1/N2 and N2/C2/O2/N3 fragments of the biuret molecule is 21.2 (2)°, which is far larger than the equivalent value of 5.06 (10)° in $[\text{Y}(\text{biur})_2(\text{H}_2\text{O})_4]\cdot 3\text{Cl}$ (Harrison, 2008). The gadolinium cation in (I) deviates from the N1/C1/O1/N2 and N2/C2/O2/N3 mean planes by -0.834 (6) Å and 0.530 (6) Å, respectively, indicating that the six-membered chelate ring is non-planar.

The component species in (I) are linked by a dense array of N—H \cdots O, N—H \cdots Cl and O—H \cdots Cl hydrogen bonds (Table 2) resulting in a three-dimensional network. Of note is the N—H \cdots O hydrogen bond, which results in [100] chains of cations, linked by $R^2_2(8)$ loops (Bernstein *et al.*, 1995), as also seen in the yttrium phase (Harrison, 2008).

For related rare-earth–biuret complexes, see: Haddad (1987 and 1988).

Experimental

0.1 M Aqueous solutions of GdCl_3 (10 ml) and biuret (10 ml) were mixed and a small quantity of dilute hydrochloric acid was added, to result in a colourless solution. Colourless blocks of (I) grew over several days as the water slowly evaporated.

Refinement

The crystal quality was rather poor, with smeared and split peaks evident in the diffraction pattern.

supplementary materials

The N-bound hydrogen atoms were geometrically placed (N—H = 0.88 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$. The water H atoms were located in difference maps and refined as riding in their as-found relative positions with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$.

The highest difference peak and deepest difference hole are 0.71 Å and 0.75 Å from Gd1, respectively.

Figures




Fig. 1. View of the molecular structure of (I) showing 50% displacement ellipsoids (arbitrary spheres for the H atoms). Symmetry code: (i) $-x, y, 3/2 - z$.

Tetraaquabis(biuret- κ^2O,O')gadolinium(III) trichloride

Crystal data

$[\text{Gd}(\text{C}_2\text{H}_5\text{N}_3\text{O}_2)_2(\text{H}_2\text{O})_4]\text{Cl}_3$

$M_r = 541.84$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 7.6501\ (3)\ \text{\AA}$

$b = 13.2164\ (5)\ \text{\AA}$

$c = 17.4557\ (6)\ \text{\AA}$

$\beta = 100.961\ (1)^\circ$

$V = 1732.69\ (11)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 1052$

$D_x = 2.077\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 6423 reflections

$\theta = 2.4\text{--}32.5^\circ$

$\mu = 4.33\ \text{mm}^{-1}$

$T = 293\ (2)\ \text{K}$

Slab, colourless

$0.47 \times 0.34 \times 0.06\ \text{mm}$

Data collection

Bruker SMART1000 CCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293\ (2)\ \text{K}$

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 1999)

$T_{\text{min}} = 0.235, T_{\text{max}} = 0.781$

8649 measured reflections

3134 independent reflections

2902 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\text{max}} = 32.5^\circ$

$\theta_{\text{min}} = 2.4^\circ$

$h = -11 \rightarrow 9$

$k = -19 \rightarrow 18$

$l = -19 \rightarrow 26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difmap (O-H) and geom (N-H)
$R[F^2 > 2\sigma(F^2)] = 0.033$	H-atom parameters constrained
$wR(F^2) = 0.088$	$w = 1/[\sigma^2(F_o^2) + (0.0654P)^2]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
3134 reflections	$(\Delta/\sigma)_{\max} < 0.001$
101 parameters	$\Delta\rho_{\max} = 2.99 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -3.45 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Gd1	0.0000	0.106486 (13)	0.7500	0.02252 (7)
Cl1	-0.04157 (10)	0.33393 (7)	0.50790 (4)	0.03672 (16)
Cl2	0.5000	0.30307 (10)	0.7500	0.0389 (2)
O1	0.2531 (3)	0.02702 (18)	0.71963 (13)	0.0310 (4)
O2	-0.0822 (3)	0.01130 (17)	0.63327 (12)	0.0305 (4)
N1	0.4595 (4)	-0.0112 (3)	0.64845 (18)	0.0434 (7)
H1	0.5423	0.0169	0.6820	0.052*
H2	0.4845	-0.0387	0.6072	0.052*
N2	0.1720 (3)	-0.0598 (2)	0.60404 (15)	0.0295 (5)
H3	0.2119	-0.0977	0.5712	0.035*
N3	-0.1037 (4)	-0.1127 (2)	0.5433 (2)	0.0371 (6)
H4	-0.2181	-0.1098	0.5337	0.044*
H5	-0.0497	-0.1552	0.5186	0.044*
C1	0.2950 (4)	-0.0121 (2)	0.66032 (16)	0.0269 (5)
C2	-0.0103 (4)	-0.0513 (2)	0.59616 (16)	0.0265 (5)
O3	-0.2462 (3)	0.18787 (18)	0.66543 (13)	0.0360 (5)
H6	-0.2805	0.1748	0.6199	0.043*
H7	-0.3244	0.2160	0.6810	0.043*

supplementary materials

O4	0.1191 (3)	0.2294 (2)	0.67206 (16)	0.0502 (7)
H8	0.0833	0.2534	0.6327	0.060*
H9	0.2181	0.2507	0.6743	0.060*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Gd1	0.01907 (9)	0.02952 (10)	0.01740 (9)	0.000	-0.00047 (6)	0.000
Cl1	0.0337 (3)	0.0470 (4)	0.0268 (3)	0.0011 (3)	-0.0007 (2)	-0.0067 (3)
Cl2	0.0301 (4)	0.0496 (6)	0.0377 (5)	0.000	0.0081 (4)	0.000
O1	0.0217 (8)	0.0441 (12)	0.0257 (9)	0.0024 (8)	0.0006 (7)	-0.0071 (8)
O2	0.0251 (9)	0.0359 (10)	0.0275 (9)	0.0038 (8)	-0.0024 (7)	-0.0077 (8)
N1	0.0233 (11)	0.073 (2)	0.0345 (14)	-0.0075 (12)	0.0071 (10)	-0.0178 (14)
N2	0.0240 (10)	0.0363 (12)	0.0269 (11)	0.0004 (9)	0.0018 (8)	-0.0083 (9)
N3	0.0297 (13)	0.0416 (15)	0.0364 (15)	-0.0020 (9)	-0.0025 (11)	-0.0147 (11)
C1	0.0227 (11)	0.0322 (12)	0.0241 (11)	0.0010 (9)	0.0005 (9)	-0.0008 (9)
C2	0.0247 (11)	0.0296 (12)	0.0231 (11)	0.0000 (9)	-0.0010 (9)	-0.0016 (9)
O3	0.0337 (10)	0.0484 (13)	0.0228 (9)	0.0135 (9)	-0.0027 (7)	-0.0022 (9)
O4	0.0384 (13)	0.0637 (17)	0.0423 (14)	-0.0157 (12)	-0.0084 (10)	0.0257 (12)

Geometric parameters (\AA , $^\circ$)

Gd1—O1 ⁱ	2.350 (2)	N1—H2	0.8600
Gd1—O1	2.350 (2)	N2—C1	1.377 (4)
Gd1—O2	2.375 (2)	N2—C2	1.380 (4)
Gd1—O2 ⁱ	2.375 (2)	N2—H3	0.8600
Gd1—O4 ⁱ	2.407 (3)	N3—C2	1.330 (4)
Gd1—O4	2.407 (3)	N3—H4	0.8600
Gd1—O3 ⁱ	2.414 (2)	N3—H5	0.8600
Gd1—O3	2.414 (2)	O3—H6	0.8067
O1—C1	1.253 (4)	O3—H7	0.7949
O2—C2	1.242 (3)	O4—H8	0.7597
N1—C1	1.314 (4)	O4—H9	0.8018
N1—H1	0.8600		
O1 ⁱ —Gd1—O1	126.90 (12)	O4—Gd1—O3	71.88 (8)
O1 ⁱ —Gd1—O2	82.06 (8)	O3 ⁱ —Gd1—O3	127.08 (12)
O1—Gd1—O2	70.41 (7)	C1—O1—Gd1	136.27 (18)
O1 ⁱ —Gd1—O2 ⁱ	70.41 (7)	C2—O2—Gd1	136.83 (18)
O1—Gd1—O2 ⁱ	82.06 (8)	C1—N1—H1	120.0
O2—Gd1—O2 ⁱ	116.04 (11)	C1—N1—H2	120.0
O1 ⁱ —Gd1—O4 ⁱ	75.96 (10)	H1—N1—H2	120.0
O1—Gd1—O4 ⁱ	147.78 (8)	C1—N2—C2	125.1 (3)
O2—Gd1—O4 ⁱ	140.85 (7)	C1—N2—H3	117.4
O2 ⁱ —Gd1—O4 ⁱ	86.53 (9)	C2—N2—H3	117.4
O1 ⁱ —Gd1—O4	147.78 (8)	C2—N3—H4	120.0
O1—Gd1—O4	75.96 (10)	C2—N3—H5	120.0

O2—Gd1—O4	86.53 (9)	H4—N3—H5	120.0
O2 ⁱ —Gd1—O4	140.85 (8)	O1—C1—N1	122.0 (3)
O4 ⁱ —Gd1—O4	95.10 (17)	O1—C1—N2	122.1 (3)
O1 ⁱ —Gd1—O3 ⁱ	129.93 (8)	N1—C1—N2	115.9 (3)
O1—Gd1—O3 ⁱ	75.91 (8)	O2—C2—N3	122.4 (3)
O2—Gd1—O3 ⁱ	144.00 (8)	O2—C2—N2	122.7 (2)
O2 ⁱ —Gd1—O3 ⁱ	70.30 (7)	N3—C2—N2	114.8 (3)
O4 ⁱ —Gd1—O3 ⁱ	71.88 (9)	Gd1—O3—H6	125.2
O4—Gd1—O3 ⁱ	73.11 (8)	Gd1—O3—H7	123.3
O1 ⁱ —Gd1—O3	75.91 (8)	H6—O3—H7	108.2
O1—Gd1—O3	129.93 (8)	Gd1—O4—H8	133.1
O2—Gd1—O3	70.30 (7)	Gd1—O4—H9	131.7
O2 ⁱ —Gd1—O3	144.00 (8)	H8—O4—H9	94.1
O4 ⁱ —Gd1—O3	73.11 (8)		
O1 ⁱ —Gd1—O1—C1	81.6 (3)	O4—Gd1—O2—C2	85.5 (3)
O2—Gd1—O1—C1	18.4 (3)	O3 ⁱ —Gd1—O2—C2	30.9 (4)
O2 ⁱ —Gd1—O1—C1	139.7 (3)	O3—Gd1—O2—C2	157.5 (3)
O4 ⁱ —Gd1—O1—C1	-149.9 (3)	Gd1—O1—C1—N1	149.2 (3)
O4—Gd1—O1—C1	-72.9 (3)	Gd1—O1—C1—N2	-31.6 (5)
O3 ⁱ —Gd1—O1—C1	-148.7 (3)	C2—N2—C1—O1	15.1 (5)
O3—Gd1—O1—C1	-21.8 (3)	C2—N2—C1—N1	-165.6 (3)
O1 ⁱ —Gd1—O2—C2	-124.7 (3)	Gd1—O2—C2—N3	161.3 (2)
O1—Gd1—O2—C2	9.2 (3)	Gd1—O2—C2—N2	-21.7 (5)
O2 ⁱ —Gd1—O2—C2	-61.1 (3)	C1—N2—C2—O2	9.9 (5)
O4 ⁱ —Gd1—O2—C2	179.3 (3)	C1—N2—C2—N3	-172.8 (3)

Symmetry codes: (i) $-x, y, -z+3/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O1 ⁱⁱ	0.86	2.10	2.910 (4)	157
N1—H2 \cdots Cl1 ⁱⁱⁱ	0.86	2.40	3.194 (3)	154
N2—H3 \cdots Cl1 ⁱⁱⁱ	0.86	2.53	3.315 (3)	153
N3—H4 \cdots Cl1 ^{iv}	0.86	2.54	3.363 (3)	161
N3—H5 \cdots Cl1 ^v	0.86	2.53	3.311 (3)	151
O3—H6 \cdots Cl1 ^{vi}	0.81	2.39	3.163 (2)	162
O3—H7 \cdots Cl2 ^{vii}	0.79	2.28	3.059 (2)	167
O4—H8 \cdots Cl1	0.76	2.45	3.208 (2)	178
O4—H9 \cdots Cl2	0.80	2.41	3.127 (2)	150

Symmetry codes: (ii) $-x+1, y, -z+3/2$; (iii) $x+1/2, y-1/2, z$; (iv) $x-1/2, y-1/2, z$; (v) $-x, -y, -z+1$; (vi) $-x-1/2, -y+1/2, -z+1$; (vii) $x-1, y, z$.

Fig. 1

