

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Poly[bis(μ_2 -benzyloxyacetato- $\kappa^3 O, O', O''$)cadmium(II)]Ji-Wei Liu^a and Seik Weng Ng^{b*}

^aCollege of Chemistry and Chemical Technology, Daqing Petroleum Institute, Daqing 163318, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 9 April 2008; accepted 18 April 2008

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(C-C) = 0.011$ Å; disorder in main residue; R factor = 0.044; wR factor = 0.125; data-to-parameter ratio = 15.2.

The title cadmium derivative of benzyloxyacetic acid, $[Cd(C_9H_9O_3)_2]_n$, exists as a μ_2 -carboxylate-bridged layer network. Two benzyloxyacetate units each chelate the metal through a carboxylate as well as through the ether O atoms; the metal is also coordinated by the double-bond carbonyl O atom of two adjacent benzyloxyacetate units in an octahedral geometry. The metal atom lies on a special position of 2 site symmetry. The phenyl group is disordered equally over two positions.

Related literature

There are no crystallographic examples of metal benzyloxyacetates although there are many examples of metal aryloxyacetates. For mononuclear diaquadi(phenoxyacetato)cadmium, see: Mak *et al.* (1985).



Experimental

Crystal data

$[Cd(C_9H_9O_3)_2]$
 $M_r = 442.72$
 Orthorhombic, $P2_12_12$
 $a = 6.7430$ (2) Å
 $b = 8.9449$ (2) Å
 $c = 15.4736$ (4) Å
 $V = 933.30$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.20$ mm⁻¹
 $T = 295$ (2) K
 $0.33 \times 0.13 \times 0.04$ mm

Data collection

Bruker APEXII diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{min} = 0.693$, $T_{max} = 0.954$
 5780 measured reflections
 1639 independent reflections
 1483 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.124$
 $S = 1.09$
 1639 reflections
 108 parameters
 37 restraints
 H-atom parameters constrained
 $\Delta\rho_{max} = 2.22$ e Å⁻³
 $\Delta\rho_{min} = -0.88$ e Å⁻³
 Absolute structure: Flack (1983),
 501 Friedel pairs
 Flack parameter: 0.03 (8)

Table 1

Selected geometric parameters (Å, °).

Cd1—O1	2.268 (4)	Cd1—O2 ⁱⁱⁱ	2.226 (5)
Cd1—O1 ⁱ	2.268 (4)	Cd1—O3	2.379 (4)
Cd1—O2 ⁱⁱ	2.226 (5)	Cd1—O3 ⁱ	2.379 (4)
O1—Cd1—O1 ⁱ	161.2 (3)	O1 ⁱ —Cd1—O3	95.5 (2)
O1—Cd1—O2 ⁱⁱ	105.9 (2)	O2 ⁱⁱ —Cd1—O2 ⁱⁱⁱ	93.3 (3)
O1—Cd1—O2 ⁱⁱⁱ	87.2 (2)	O2 ⁱⁱ —Cd1—O3	98.3 (2)
O1—Cd1—O3	69.6 (2)	O2 ⁱⁱ —Cd1—O3 ⁱ	156.2 (2)
O1—Cd1—O3 ⁱ	95.5 (2)	O2 ⁱⁱⁱ —Cd1—O3	156.2 (2)
O1 ⁱ —Cd1—O2 ⁱⁱ	87.2 (2)	O2 ⁱⁱⁱ —Cd1—O3 ⁱ	98.3 (2)
O1 ⁱ —Cd1—O2 ⁱⁱⁱ	105.9 (2)	O3—Cd1—O3 ⁱ	79.1 (2)
O1 ⁱ —Cd1—O3 ⁱ	69.6 (2)		

Symmetry codes: (i) $-x + 2, -y + 2, z$; (ii) $x - \frac{1}{2}, -y + \frac{3}{2}, -z$; (iii) $-x + \frac{5}{2}, y + \frac{1}{2}, -z$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); OLEX (Dolomanov *et al.*, 2003); software used to prepare material for publication: SHELXL97.

The authors thank Daqing Petroleum Institute and the University of Malaya for generously supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SG2235).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2006). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Dolomanov, O. V., Blake, A. J., Champness, N. R. & Schröder, M. (2003). *J. Appl. Cryst.* **36**, 1283–1284.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Mak, T. C. W., Yip, H.-W., O'Reilly, E. J., Smith, G. & Kennard, C. H. L. (1985). *Inorg. Chim. Acta*, **100**, 267–273.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, m708 [doi:10.1107/S1600536808010799]

Poly[bis(μ_2 -benzyloxyacetato- $\kappa^3O,O':O''$)cadmium(II)]

J.-W. Liu and S. W. Ng

Comment

The crystal structures of a large number of metal derivatives of aryloxyacetic acids have been reported; in some structures, the ether oxygen also engages in bonding so that the carboxylate unit functions both as a chelate as well as a bridge. The cadmium derivative of phenoxyacetic acid exists as a diaqua, carboxylate-chelated compound. The carboxyl $-\text{CO}_2$ portion engages in chelation instead (Mak *et al.*, 1985). The title cadmium analog has a benzyl group in place of the phenyl group, which is probably less crowded; this feature permits the ether linkage to bind to the metal atom. The compound (Scheme 1) is an anhydrous compound; the carboxylate group chelates to the metal atom. It also bridges adjacent metal atoms (Fig. 1); the bridges lead to the formation of a layer motif (Fig. 2).

Experimental

Cadmium dinitrate tetrahydrate (0.31 g, 1 mmol) and 2,2'-bipyridine (0.16 g, 1 mmol) were added to a hot aqueous solution of benzyloxyacetic acid (0.17 g, 1 mmol). The pH of the solution was adjusted to 6 with 0.1 M sodium hydroxide. The solution was allowed to evaporate at room temperature. Colorless single crystals are separated from the filtered solution after several days.

Refinement

Hydrogen atoms were treated as riding, with C–H = 0.93 to 0.97 Å and were included in the refinement with $U(\text{H})$ set to 1.2 times $U_{\text{eq}}(\text{C})$. The phenyl ring is disordered over two sites; the occupancy could not be refined, and each component was arbitrarily assigned 0.5 occupancy. The ring was refined as a rigid hexagon; the temperature factors of the primed atoms were constrained to those of the unprimed ones. The anisotropic temperature factors of the ring were restrained to be nearly isotropic. The C3–C4 and C3–C4' distances were restrained to within 0.01 Å of each other.

The final difference Fourier map had a large peak at about 1 Å from Cd1.

Figures




Fig. 1. Thermal displacement ellipsoid plot (Barbour, 2001) illustrating the coordination geometry of Cd in $[\text{Cd}(\text{C}_9\text{H}_9\text{O}_3)_2]_n$. Displacement ellipsoids are drawn at the 50% probability level and H atoms as spheres of arbitrary radii.




Fig. 2. OLEX (Dolomanov *et al.*, 2003) representation of the layer structure.

Poly[bis(μ_2 -benzyloxyacetato- κ^3 O,O':\langle i \rangleO'')cadmium(II)]

Crystal data

[Cd(C ₉ H ₉ O ₃) ₂]	$F_{000} = 444$
$M_r = 442.72$	$D_x = 1.575 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12$	Mo $K\alpha$ radiation
Hall symbol: P 2 2ab	$\lambda = 0.71073 \text{ \AA}$
$a = 6.7430 (2) \text{ \AA}$	Cell parameters from 2270 reflections
$b = 8.9449 (2) \text{ \AA}$	$\theta = 2.6\text{--}23.6^\circ$
$c = 15.4736 (4) \text{ \AA}$	$\mu = 1.20 \text{ mm}^{-1}$
$V = 933.30 (4) \text{ \AA}^3$	$T = 295 (2) \text{ K}$
$Z = 2$	Block, colorless
	$0.33 \times 0.13 \times 0.04 \text{ mm}$

Data collection

Bruker APEXII diffractometer	1639 independent reflections
Radiation source: fine-focus sealed tube	1483 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.039$
$T = 295(2) \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.3^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -7 \rightarrow 7$
$T_{\text{min}} = 0.693$, $T_{\text{max}} = 0.954$	$k = -10 \rightarrow 9$
5780 measured reflections	$l = -18 \rightarrow 17$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.0833P)^2]$
$wR(F^2) = 0.125$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\text{max}} = 0.001$
1639 reflections	$\Delta\rho_{\text{max}} = 2.22 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.88 \text{ e \AA}^{-3}$

108 parameters
 37 restraints
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map
 Extinction correction: none
 Absolute structure: Flack (1983), 501 Friedel pairs
 Flack parameter: 0.03 (8)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cd1	1.0000	1.0000	0.01184 (4)	0.0405 (3)	
O1	1.2315 (7)	0.8208 (5)	0.0358 (3)	0.0481 (12)	
O2	1.2948 (7)	0.5938 (5)	0.0869 (3)	0.0495 (12)	
O3	0.9048 (6)	0.8466 (5)	0.1304 (3)	0.0443 (11)	
C1	1.1913 (10)	0.7127 (7)	0.0819 (4)	0.0385 (15)	
C2	1.0100 (14)	0.7103 (6)	0.1367 (4)	0.0427 (13)	
H2A	0.9248	0.6287	0.1186	0.051*	
H2B	1.0472	0.6931	0.1964	0.051*	
C3	0.7017 (11)	0.8363 (9)	0.1615 (5)	0.059 (2)	
H3A	0.6423	0.7463	0.1381	0.070*	
H3B	0.6276	0.9209	0.1392	0.070*	
C4	0.680 (3)	0.834 (2)	0.2552 (5)	0.060 (3)	0.50
C5	0.567 (2)	0.7180 (17)	0.2895 (9)	0.100 (5)	0.50
H5	0.5061	0.6494	0.2529	0.119*	0.50
C6	0.545 (3)	0.705 (2)	0.3785 (10)	0.124 (6)	0.50
H6	0.4689	0.6276	0.4014	0.149*	0.50
C7	0.636 (3)	0.807 (2)	0.4332 (5)	0.127 (8)	0.50
H7	0.6206	0.7986	0.4927	0.152*	0.50
C8	0.749 (4)	0.923 (2)	0.3989 (9)	0.135 (5)	0.50
H8	0.8096	0.9915	0.4355	0.162*	0.50
C9	0.771 (4)	0.936 (2)	0.3099 (10)	0.108 (5)	0.50
H9	0.8468	1.0133	0.2870	0.130*	0.50
C4'	0.704 (3)	0.8030 (19)	0.2550 (5)	0.060 (3)	0.50
C5'	0.682 (3)	0.6579 (16)	0.2856 (9)	0.100 (5)	0.50
H5'	0.6602	0.5798	0.2471	0.119*	0.50
C6'	0.691 (3)	0.6296 (17)	0.3739 (10)	0.124 (6)	0.50
H6'	0.6755	0.5325	0.3943	0.149*	0.50
C7'	0.723 (3)	0.746 (2)	0.4315 (6)	0.127 (8)	0.50
H7'	0.7288	0.7273	0.4905	0.152*	0.50
C8'	0.745 (4)	0.891 (2)	0.4009 (9)	0.135 (5)	0.50
H8'	0.7667	0.9695	0.4395	0.162*	0.50
C9'	0.736 (4)	0.9198 (16)	0.3127 (10)	0.108 (5)	0.50
H9'	0.7514	1.0168	0.2922	0.130*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.0336 (4)	0.0237 (3)	0.0641 (4)	0.0012 (3)	0.000	0.000
O1	0.041 (3)	0.033 (3)	0.070 (3)	0.010 (2)	0.012 (2)	0.006 (3)
O2	0.048 (3)	0.030 (2)	0.070 (3)	0.010 (2)	0.006 (2)	0.003 (2)

supplementary materials

O3	0.037 (2)	0.028 (2)	0.068 (3)	0.0084 (19)	0.012 (2)	0.008 (2)
C1	0.036 (4)	0.027 (3)	0.053 (4)	0.006 (3)	-0.003 (3)	-0.005 (3)
C2	0.037 (3)	0.028 (3)	0.063 (3)	0.007 (4)	-0.006 (6)	0.004 (2)
C3	0.033 (4)	0.056 (4)	0.087 (6)	0.016 (4)	0.015 (4)	0.008 (4)
C4	0.054 (6)	0.054 (6)	0.072 (5)	0.009 (6)	0.014 (4)	-0.012 (4)
C5	0.109 (10)	0.098 (9)	0.092 (7)	0.006 (7)	0.031 (7)	0.016 (7)
C6	0.126 (11)	0.126 (9)	0.120 (8)	-0.006 (8)	0.019 (8)	0.020 (8)
C7	0.132 (11)	0.134 (11)	0.114 (9)	0.012 (9)	0.018 (8)	-0.004 (8)
C8	0.134 (9)	0.152 (9)	0.119 (8)	-0.005 (8)	-0.001 (7)	-0.031 (7)
C9	0.091 (9)	0.122 (8)	0.112 (7)	-0.008 (7)	0.007 (6)	-0.025 (6)
C4'	0.054 (6)	0.054 (6)	0.072 (5)	0.009 (6)	0.014 (4)	-0.012 (4)
C5'	0.109 (10)	0.098 (9)	0.092 (7)	0.006 (7)	0.031 (7)	0.016 (7)
C6'	0.126 (11)	0.126 (9)	0.120 (8)	-0.006 (8)	0.019 (8)	0.020 (8)
C7'	0.132 (11)	0.134 (11)	0.114 (9)	0.012 (9)	0.018 (8)	-0.004 (8)
C8'	0.134 (9)	0.152 (9)	0.119 (8)	-0.005 (8)	-0.001 (7)	-0.031 (7)
C9'	0.091 (9)	0.122 (8)	0.112 (7)	-0.008 (7)	0.007 (6)	-0.025 (6)

Geometric parameters (\AA , $^\circ$)

Cd1—O1	2.268 (4)	C5—C6	1.3900
Cd1—O1 ⁱ	2.268 (4)	C5—H5	0.9300
Cd1—O2 ⁱⁱ	2.226 (5)	C6—C7	1.3900
Cd1—O2 ⁱⁱⁱ	2.226 (5)	C6—H6	0.9300
Cd1—O3	2.379 (4)	C7—C8	1.3900
Cd1—O3 ⁱ	2.379 (4)	C7—H7	0.9300
O1—C1	1.232 (8)	C8—C9	1.3900
O2—C1	1.274 (7)	C8—H8	0.9300
O2—Cd1 ^{iv}	2.226 (5)	C9—H9	0.9300
O3—C2	1.414 (7)	C4'—C5'	1.3900
O3—C3	1.455 (8)	C4'—C9'	1.3900
C1—C2	1.488 (11)	C5'—C6'	1.3900
C2—H2A	0.9700	C5'—H5'	0.9300
C2—H2B	0.9700	C6'—C7'	1.3900
C3—C4	1.457 (11)	C6'—H6'	0.9300
C3—C4'	1.477 (11)	C7'—C8'	1.3900
C3—H3A	0.9700	C7'—H7'	0.9300
C3—H3B	0.9700	C8'—C9'	1.3900
C4—C5	1.3900	C8'—H8'	0.9300
C4—C9	1.3900	C9'—H9'	0.9300
O1—Cd1—O1 ⁱ	161.2 (3)	H3A—C3—H3B	107.5
O1—Cd1—O2 ⁱⁱ	105.9 (2)	C5—C4—C9	120.0
O1—Cd1—O2 ⁱⁱⁱ	87.2 (2)	C5—C4—C3	116.6 (12)
O1—Cd1—O3	69.6 (2)	C9—C4—C3	123.4 (12)
O1—Cd1—O3 ⁱ	95.5 (2)	C6—C5—C4	120.0
O1 ⁱ —Cd1—O2 ⁱⁱ	87.2 (2)	C6—C5—H5	120.0
O1 ⁱ —Cd1—O2 ⁱⁱⁱ	105.9 (2)	C4—C5—H5	120.0

O1 ⁱ —Cd1—O3 ⁱ	69.6 (2)	C5—C6—C7	120.0
O1 ⁱ —Cd1—O3	95.5 (2)	C5—C6—H6	120.0
O2 ⁱⁱ —Cd1—O2 ⁱⁱⁱ	93.3 (3)	C7—C6—H6	120.0
O2 ⁱⁱ —Cd1—O3	98.3 (2)	C6—C7—C8	120.0
O2 ⁱⁱ —Cd1—O3 ⁱ	156.2 (2)	C6—C7—H7	120.0
O2 ⁱⁱⁱ —Cd1—O3	156.2 (2)	C8—C7—H7	120.0
O2 ⁱⁱⁱ —Cd1—O3 ⁱ	98.3 (2)	C9—C8—C7	120.0
O3—Cd1—O3 ⁱ	79.1 (2)	C9—C8—H8	120.0
C1—O1—Cd1	119.9 (4)	C7—C8—H8	120.0
C1—O2—Cd1 ^{iv}	127.8 (5)	C8—C9—C4	120.0
C2—O3—C3	113.3 (6)	C8—C9—H9	120.0
C2—O3—Cd1	114.5 (4)	C4—C9—H9	120.0
C3—O3—Cd1	123.1 (4)	C5'—C4'—C9'	120.0
O1—C1—O2	124.7 (6)	C5'—C4'—C3	121.4 (12)
O1—C1—C2	121.5 (5)	C9'—C4'—C3	118.6 (12)
O2—C1—C2	113.8 (6)	C6'—C5'—C4'	120.0
O3—C2—C1	111.1 (5)	C6'—C5'—H5'	120.0
O3—C2—H2A	109.4	C4'—C5'—H5'	120.0
C1—C2—H2A	109.4	C5'—C6'—C7'	120.0
O3—C2—H2B	109.4	C5'—C6'—H6'	120.0
C1—C2—H2B	109.4	C7'—C6'—H6'	120.0
H2A—C2—H2B	108.0	C6'—C7'—C8'	120.0
O3—C3—C4	115.1 (10)	C6'—C7'—H7'	120.0
O3—C3—C4'	109.0 (11)	C8'—C7'—H7'	120.0
O3—C3—H3A	108.5	C9'—C8'—C7'	120.0
C4—C3—H3A	108.5	C9'—C8'—H8'	120.0
C4'—C3—H3A	101.8	C7'—C8'—H8'	120.0
O3—C3—H3B	108.5	C8'—C9'—C4'	120.0
C4—C3—H3B	108.5	C8'—C9'—H9'	120.0
C4'—C3—H3B	120.9	C4'—C9'—H9'	120.0
O2 ⁱⁱ —Cd1—O1—C1	-78.1 (6)	Cd1—O3—C2—C1	15.2 (7)
O2 ⁱⁱⁱ —Cd1—O1—C1	-170.7 (5)	O1—C1—C2—O3	-2.7 (10)
O1 ⁱ —Cd1—O1—C1	54.5 (5)	O2—C1—C2—O3	178.2 (6)
O3 ⁱ —Cd1—O1—C1	91.2 (6)	C2—O3—C3—C4	76.0 (10)
O3—Cd1—O1—C1	15.0 (5)	Cd1—O3—C3—C4	-139.0 (9)
O2 ⁱⁱ —Cd1—O3—C2	88.3 (5)	C2—O3—C3—C4'	64.4 (10)
O2 ⁱⁱⁱ —Cd1—O3—C2	-29.9 (7)	Cd1—O3—C3—C4'	-150.7 (8)
O1 ⁱ —Cd1—O3—C2	176.3 (5)	O3—C3—C4—C5	-128.2 (12)
O1—Cd1—O3—C2	-15.6 (4)	C4'—C3—C4—C5	-65 (6)
O3 ⁱ —Cd1—O3—C2	-115.7 (5)	O3—C3—C4—C9	49.7 (12)
O2 ⁱⁱ —Cd1—O3—C3	-56.3 (6)	C4'—C3—C4—C9	113 (7)
O2 ⁱⁱⁱ —Cd1—O3—C3	-174.5 (5)	C3—C4—C5—C6	178.0 (15)
O1 ⁱ —Cd1—O3—C3	31.7 (5)	C3—C4—C9—C8	-177.8 (16)
O1—Cd1—O3—C3	-160.2 (6)	O3—C3—C4'—C5'	-97.6 (15)

supplementary materials

O3 ⁱ —Cd1—O3—C3	99.7 (5)	C4—C3—C4'—C5'	141 (8)
Cd1—O1—C1—O2	166.5 (5)	O3—C3—C4'—C9'	80.5 (10)
Cd1—O1—C1—C2	-12.5 (9)	C4—C3—C4'—C9'	-41 (7)
Cd1 ^{iv} —O2—C1—O1	-25.2 (10)	C3—C4'—C5'—C6'	178.1 (17)
Cd1 ^{iv} —O2—C1—C2	153.9 (4)	C3—C4'—C9'—C8'	-178.1 (17)
C3—O3—C2—C1	163.3 (6)		

Symmetry codes: (i) $-x+2, -y+2, z$; (ii) $x-1/2, -y+3/2, -z$; (iii) $-x+5/2, y+1/2, -z$; (iv) $x+1/2, -y+3/2, -z$.

Fig. 1




Fig. 2

