

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-[5-(4-Pyridyl)-1,3,4-oxadiazol-2-yl]-pyridine *N*-oxide–isophthalic acid (1/1)

Gui-Ge Hou, Li-Li Liu, Jian-Ping Ma, Ru-Qi Huang and Yu-Bin Dong*

 College of Chemistry, Chemical Engineering and Materials Science, Shandong Normal University, Jinan 250014, People's Republic of China
 Correspondence e-mail: yubindong@sdnu.edu.cn


Received 6 March 2008; accepted 30 April 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.052; wR factor = 0.132; data-to-parameter ratio = 11.7.

The title compound, $\text{C}_{12}\text{H}_8\text{N}_4\text{O}_2 \cdot \text{C}_8\text{H}_6\text{O}_4$, was synthesized from 4-[5-(4-pyridyl)-1,3,4-oxadiazol-2-yl]pyridine *N*-oxide and isophthalic acid. The two molecules are linked through $\text{O}-\text{H} \cdots \text{O}$ and $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds. Weak intramolecular $\pi-\pi$ interactions between the two hydrogen-bonded chains result in the formation a one-dimensional supramolecular curved tape (the face-to-face distance between the pyridine *N*-oxide ring and the benzene ring is 3.7 Å).

Related literature

For related literature, see: Beckmann & Jänicke (2006); Dong *et al.* (2003); Du *et al.* (2006); Hunter (1994); Kitagawa *et al.* (2004); Long *et al.* (2004); Lu *et al.* (1997); Ma *et al.* (2005); Ren *et al.* (1995); Tan *et al.* (2006).



Experimental

Crystal data

 $\text{C}_{12}\text{H}_8\text{N}_4\text{O}_2 \cdot \text{C}_8\text{H}_6\text{O}_4$
 $M_r = 406.35$
 Triclinic, $P\bar{1}$
 $a = 7.0993$ (18) Å
 $b = 7.1770$ (19) Å
 $c = 19.823$ (5) Å
 $\alpha = 93.003$ (4)°
 $\beta = 98.481$ (3)°

 $\gamma = 112.745$ (4)°
 $V = 914.6$ (4) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 298$ (2) K
 $0.30 \times 0.15 \times 0.06$ mm

Data collection

 Bruker SMART APEX
 diffractometer
 Absorption correction: none
 4661 measured reflections

 3189 independent reflections
 1972 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.027$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.132$
 $S = 0.98$
 3189 reflections

 272 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.30$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.15$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O1}-\text{H1} \cdots \text{N4}$	0.82	1.82	2.640 (2)	171
$\text{O3}-\text{H3} \cdots \text{O5}^i$	0.82	1.72	2.521 (3)	163

 Symmetry code: (i) $x + 1, y + 1, z + 1$.

Data collection: *SMART* (Bruker 2000); cell refinement: *SMART*; data reduction: *SAINT* (Bruker 2000); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (grant No. 20671060) and Shangdong Natural Science Foundation (grant Nos. J06D05 and 2006BS04040) for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BV2095).

References

- Beckmann, J. & Jänicke, S. L. (2006). *Eur. J. Inorg. Chem.* pp. 3351–3358.
 Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Dong, Y.-B., Ma, J.-P., Huang, R.-Q., Smith, M. D. & zur Loye, H.-C. (2003). *Inorg. Chem.* **42**, 294–300.
 Du, M., Zhang, Z.-H. & Zhao, X.-J. (2006). *Cryst. Growth Des.* **6**, 390–396.
 Hunter, C. A. (1994). *Chem. Soc. Rev.* **23**, 101–102.
 Kitagawa, S., Kitaura, R. & Noro, S.-I. (2004). *Angew. Chem. Int. Ed.* **43**, 2334–2375.
 Long, D.-L., Hill, R. J., Blake, A. J., Champness, N. R., Hubberstey, P., Proserpio, D. M., Wilson, C. & Schroder, M. (2004). *Angew. Chem. Int. Ed.* **43**, 1851–1854.
 Lu, J., Paliwala, T., Lim, S. C., Yu, C., Niu, T. & Jacobson, A. J. (1997). *Inorg. Chem.* **36**, 923–929.
 Ma, B.-Q., Sun, H.-L. & Gao, S. (2005). *Inorg. Chem.* **44**, 837–839.
 Ren, Z.-J., Jiang, E. & Zhou, H.-B. (1995). *Youji Huaxue*, **15**, 218–220.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Tan, T.-F., Han, J., Pang, M.-L., Song, H.-B., Ma, Y.-X. & Meng, J.-B. (2006). *Cryst. Growth Des.* **6**, 1186–1193.

supplementary materials

Acta Cryst. (2008). E64, o997 [doi:10.1107/S1600536808012622]

4-[5-(4-Pyridyl)-1,3,4-oxadiazol-2-yl]pyridine *N*-oxide-isophthalic acid (1/1)

G.-G. Hou, L.-L. Liu, J.-P. Ma, R.-Q. Huang and Y.-B. Dong

Comment

Rigid pyridyl compounds have been considered as good hydrogen bond acceptors. These compounds include 4,4-bipyridine (Lu *et al.*, 1997; Kitagawa *et al.*, 2004; Tan *et al.*, 2006), 1,2-bis(4-pyridyl)ethyne (Beckmann & Jänicke, 2006), 4,4'-bipyridine-*N,N'*-dioxide (Long *et al.*, 2004; Ma *et al.*, 2005), and other similar molecules. We recently prepared the rigid bent compound, 2-(4-pyridyl)-5-(4-pyridyl *N*-oxide)-1,3,4-oxadiazole, (L2). It has a rigid 144° angle because of the bridging five-membered oxadiazole ring. In addition, isophthalic acid is a good hydrogen-bonding donor and its two carboxylic acid moieties make an angle of 120°. Supramolecular systems with novel architecture may result because of their specific geometry. Accordingly, when L2 reacted with isophthalic acid in a mixed CH₂Cl₂/CH₃OH solution, yellow crystals formed and its structure is reported here.

The molecular structure of (1) is shown in Fig. 1. The asymmetric unit contains one L2 molecule and one isophthalic acid molecule. O··H—O and N··H—O hydrogen bonds (Fig. 3) are formed by the —COOH groups of the isophthalic acid and the pyridyl-*N*-oxide O atom and N_{pyridine} atom, (Table 1). The dihedral angles between the benzene ring of isophthalic acid and pyridyl-*N*-oxide ring and pyridyl ring are 1.4° and 7.0°, respectively, which are similar to previously reported experimental values (Du *et al.*, 2006; Tan *et al.*, 2006). In addition to the specific geometry of the oxadiazole-containing rigid curved organic ligand L2, the molecules in (1) connect with each other generating one-dimensional extended zigzag chains, which have not been obtained using normal linear rigid bidentate organic ligands (Dong & Ma, 2003).

In the solid state, the crystal packing view of (1) shows a pair of zigzag chains which stack *via* aromatic π - π interactions (Fig. 3). The face-to-face distance between the pyridine-*N*-oxide ring and benzene ring from isophthalic acid is 3.7 Å. The shortest close contact is 3.447 (6) Å. Although these values are typical for aromatic π - π stacking interactions, compared with the strong π - π interactions (Hunter, 1994) they are comparatively weak. Two adjacent chains are further linked *via* intramolecular π - π stacking interactions to construct a one-dimensional supramolecular curved tape (Fig. 3). These weak intramolecular π - π interactions and crucial hydrogen bonds enhance the stability of the compound (1).

Experimental

L1 (2,5-bis(4-pyridyl)-1,3,4-oxadiazole) was prepared according to literature methods (Ren *et al.*, 1995). L1 (1.1 g, 5 mmol), 1.0 ml hydrogen dioxide 30% solution and 3.0 ml acetic acid were mixed and refluxed at 343–353 K for 20 h. After removal of solvent under vacuum, the residue was purified on a silica gel column using CH₂Cl₂/CH₃OH (15:1, *v/v*) as the eluent to afford L2, Yield: 38%. Mp: 501–503 K. ¹H NMR (DMSO, 300 MHz, *p.p.m.*): δ 8.89–8.91 (d, 2H, 2C₅H₄N), 8.34–8.37 (d, 2H, 2C₅H₄NO), 7.99–8.06 (m, 4H, 2C₅H₄N, 2C₅H₄NO). IR (KBr pellet cm⁻¹): 3423(*s*), 1620(*m*), 1563(*m*), 1537(*m*), 1475(*s*), 1439(*s*), 1407(*s*), 1272(*s*), 1174(*s*), 1111(*m*), 965(*m*), 848(*m*), 740(*m*), 703(*s*), 644(*s*), 509(*m*).

A CH₂Cl₂ and CH₃OH solution (10 ml, 1:1, *v/v*) of L2 (24 mg, 0.1 mmol) and isophthalic acid (16.6 mg, 0.1 mmol) kept at room temperature. After a few days yellow block crystals (1) were obtained (36 mg). Yield: 89%.

Refinement

H atoms on O atoms were located in a difference Fourier map and refined as riding in their as-found relative positions, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. Other H atoms were placed in calculated positions, with C—H = 0.93 Å, and refined in riding mode, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ (aromatic).

Figures




Fig. 1. : The structure of (1), with 30% probability displacement ellipsoids (arbitrary spheres for H atoms).




Fig. 2. : A view of the hydrogen-bonded chains observed in the crystal structure of (1).




Fig. 3. : The crystal packing of (1) via weak π - π interactions, and the face-to-face distance is 3.7 Å.



Fig. 4. The formation of L2.

4-[5-(4-pyridyl)-1,3,4-oxadiazol-2-yl]pyridine *N*-oxide–isophthalic acid (1/1)

Crystal data

$\text{C}_{12}\text{H}_8\text{N}_4\text{O}_2 \cdot \text{C}_8\text{H}_6\text{O}_4$

$Z = 2$

$M_r = 406.35$

$F_{000} = 420$

Triclinic, $P\bar{1}$

$D_x = 1.475 \text{ Mg m}^{-3}$

Hall symbol: -P 1

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

$a = 7.0993 (18) \text{ \AA}$

Cell parameters from 893 reflections

$b = 7.1770 (19) \text{ \AA}$

$\theta = 3.2\text{--}21.1^\circ$

$c = 19.823 (5) \text{ \AA}$

$\mu = 0.11 \text{ mm}^{-1}$

$\alpha = 93.003 (4)^\circ$

$T = 298 (2) \text{ K}$

$\beta = 98.481 (3)^\circ$

Bar, colourless

$\gamma = 112.745 (4)^\circ$

$0.30 \times 0.15 \times 0.06 \text{ mm}$

$V = 914.6 (4) \text{ \AA}^3$

Data collection

Bruker SMART APEX
diffractometer

1972 reflections with $I > 2\sigma(I)$

Radiation source: fine-focus sealed tube

$R_{\text{int}} = 0.027$

Monochromator: graphite

$\theta_{\text{max}} = 25.0^\circ$

$T = 298(2) \text{ K}$

$\theta_{\text{min}} = 3.1^\circ$

φ and ω scans

$h = -8 \rightarrow 8$

Absorption correction: none $k = -8 \rightarrow 8$
 4661 measured reflections $l = -17 \rightarrow 23$
 3189 independent reflections

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.052$	H-atom parameters constrained
$wR(F^2) = 0.132$	$w = 1/[\sigma^2(F_o^2) + (0.0637P)^2]$
$S = 0.98$	where $P = (F_o^2 + 2F_c^2)/3$
3189 reflections	$(\Delta/\sigma)_{\max} = 0.001$
272 parameters	$\Delta\rho_{\max} = 0.30 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.15 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.7708 (3)	0.9021 (3)	0.64281 (9)	0.0693 (6)
O2	1.0221 (3)	0.8709 (4)	0.59557 (10)	0.0861 (7)
O5	-0.2941 (3)	0.2221 (4)	-0.02763 (9)	0.0882 (8)
O3	0.8643 (3)	1.0843 (3)	0.88695 (9)	0.0738 (6)
H3	0.8309	1.1276	0.9203	0.111*
O4	1.1774 (3)	1.2181 (3)	0.95238 (9)	0.0726 (6)
C1	-0.4071 (4)	0.2792 (4)	0.06980 (13)	0.0592 (8)
H1A	-0.5408	0.2374	0.0446	0.071*
C2	-0.3697 (4)	0.3405 (4)	0.13821 (12)	0.0528 (7)
H2	-0.4781	0.3399	0.1595	0.063*
C3	-0.1721 (4)	0.4035 (4)	0.17649 (11)	0.0427 (6)
C4	-0.0155 (4)	0.4016 (4)	0.14322 (12)	0.0515 (7)
H4	0.1189	0.4429	0.1678	0.062*
C5	-0.0572 (4)	0.3385 (4)	0.07365 (12)	0.0558 (7)
H5A	0.0489	0.3375	0.0513	0.067*

supplementary materials

C6	-0.1345 (4)	0.4706 (4)	0.24973 (12)	0.0436 (6)
C7	0.0337 (4)	0.5784 (4)	0.35129 (11)	0.0431 (6)
C8	0.2083 (4)	0.6463 (4)	0.40831 (11)	0.0415 (6)
C9	0.1803 (4)	0.7064 (4)	0.47216 (12)	0.0496 (7)
H9	0.0533	0.7076	0.4782	0.059*
C10	0.3421 (4)	0.7639 (4)	0.52633 (12)	0.0520 (7)
H10	0.3205	0.8000	0.5694	0.062*
C11	0.5555 (4)	0.7154 (4)	0.45907 (13)	0.0551 (7)
H11	0.6856	0.7205	0.4542	0.066*
C12	0.4001 (4)	0.6505 (4)	0.40169 (12)	0.0490 (7)
H12	0.4245	0.6105	0.3596	0.059*
C13	0.9602 (4)	0.9161 (4)	0.64414 (14)	0.0521 (7)
C14	1.0945 (4)	0.9948 (4)	0.71368 (11)	0.0453 (6)
C15	1.0149 (4)	1.0271 (4)	0.77116 (12)	0.0451 (6)
H15	0.8738	0.9975	0.7670	0.054*
C16	1.1455 (4)	1.1037 (4)	0.83507 (11)	0.0436 (6)
C17	1.3553 (4)	1.1470 (4)	0.84013 (13)	0.0533 (7)
H17	1.4443	1.1995	0.8822	0.064*
C18	1.4324 (4)	1.1130 (4)	0.78334 (13)	0.0577 (8)
H18	1.5734	1.1419	0.7874	0.069*
C19	1.3043 (4)	1.0369 (4)	0.72065 (13)	0.0542 (7)
H19	1.3586	1.0134	0.6827	0.065*
C20	1.0666 (4)	1.1418 (4)	0.89757 (13)	0.0509 (7)
N1	-0.2515 (3)	0.2788 (3)	0.03830 (10)	0.0569 (6)
N2	-0.2670 (3)	0.4826 (3)	0.28576 (10)	0.0511 (6)
N3	-0.1552 (3)	0.5550 (3)	0.35274 (10)	0.0529 (6)
N4	0.5295 (3)	0.7709 (3)	0.52071 (10)	0.0525 (6)
H1	0.7016	0.8526	0.6043	0.079*
O6	0.0597 (2)	0.5275 (2)	0.28698 (7)	0.0451 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0608 (14)	0.0982 (17)	0.0411 (11)	0.0313 (12)	-0.0073 (9)	-0.0095 (10)
O2	0.0854 (16)	0.128 (2)	0.0449 (12)	0.0492 (15)	0.0062 (11)	-0.0239 (12)
O5	0.0759 (15)	0.157 (2)	0.0339 (12)	0.0592 (15)	-0.0082 (10)	-0.0223 (12)
O3	0.0545 (13)	0.1118 (18)	0.0450 (12)	0.0277 (12)	0.0025 (9)	-0.0144 (11)
O4	0.0630 (13)	0.1144 (18)	0.0341 (11)	0.0374 (12)	-0.0070 (9)	-0.0135 (11)
C1	0.0494 (17)	0.077 (2)	0.0459 (17)	0.0259 (15)	-0.0048 (13)	-0.0079 (14)
C2	0.0514 (17)	0.0658 (19)	0.0379 (16)	0.0239 (14)	0.0000 (12)	-0.0024 (13)
C3	0.0490 (16)	0.0424 (16)	0.0352 (14)	0.0190 (12)	0.0020 (12)	0.0006 (11)
C4	0.0498 (17)	0.0596 (19)	0.0377 (16)	0.0182 (14)	-0.0013 (12)	-0.0006 (12)
C5	0.0523 (18)	0.075 (2)	0.0405 (16)	0.0282 (15)	0.0051 (13)	0.0002 (14)
C6	0.0501 (17)	0.0421 (16)	0.0331 (14)	0.0167 (13)	-0.0027 (12)	0.0002 (11)
C7	0.0534 (17)	0.0443 (16)	0.0326 (15)	0.0210 (13)	0.0079 (12)	0.0010 (11)
C8	0.0474 (16)	0.0436 (16)	0.0313 (14)	0.0182 (12)	0.0019 (11)	0.0009 (11)
C9	0.0487 (16)	0.0590 (18)	0.0375 (15)	0.0193 (14)	0.0060 (12)	-0.0018 (12)
C10	0.0591 (19)	0.0614 (19)	0.0319 (15)	0.0225 (15)	0.0052 (13)	-0.0025 (12)

C11	0.0491 (17)	0.066 (2)	0.0508 (18)	0.0258 (15)	0.0033 (14)	0.0032 (14)
C12	0.0551 (18)	0.0575 (18)	0.0338 (15)	0.0228 (14)	0.0073 (13)	-0.0006 (12)
C13	0.0625 (19)	0.0471 (18)	0.0468 (18)	0.0236 (14)	0.0068 (14)	0.0024 (13)
C14	0.0574 (17)	0.0467 (17)	0.0315 (14)	0.0233 (13)	0.0020 (12)	-0.0004 (11)
C15	0.0472 (15)	0.0468 (16)	0.0417 (15)	0.0221 (13)	0.0003 (12)	0.0039 (12)
C16	0.0492 (16)	0.0482 (16)	0.0329 (14)	0.0220 (13)	-0.0005 (12)	0.0029 (11)
C17	0.0537 (18)	0.0604 (19)	0.0428 (16)	0.0251 (15)	-0.0042 (13)	-0.0001 (13)
C18	0.0508 (17)	0.073 (2)	0.0485 (18)	0.0270 (16)	0.0037 (14)	0.0004 (15)
C19	0.0609 (19)	0.0605 (19)	0.0458 (17)	0.0285 (15)	0.0126 (14)	0.0029 (13)
C20	0.0509 (18)	0.0600 (19)	0.0408 (16)	0.0251 (14)	-0.0007 (13)	0.0014 (13)
N1	0.0585 (16)	0.0779 (18)	0.0301 (13)	0.0295 (13)	-0.0054 (11)	-0.0088 (11)
N2	0.0505 (14)	0.0635 (16)	0.0366 (13)	0.0230 (12)	0.0021 (10)	-0.0024 (10)
N3	0.0525 (15)	0.0684 (16)	0.0351 (13)	0.0244 (12)	0.0016 (10)	-0.0013 (10)
N4	0.0582 (15)	0.0540 (15)	0.0382 (13)	0.0189 (12)	-0.0012 (11)	0.0011 (10)
O6	0.0486 (11)	0.0548 (11)	0.0293 (9)	0.0211 (9)	0.0009 (8)	-0.0035 (7)

Geometric parameters (Å, °)

O1—C13	1.306 (3)	C8—C12	1.377 (3)
O1—H1	0.8221	C8—C9	1.382 (3)
O2—C13	1.195 (3)	C9—C10	1.366 (3)
O5—N1	1.303 (2)	C9—H9	0.9300
O3—C20	1.312 (3)	C10—N4	1.333 (3)
O3—H3	0.8200	C10—H10	0.9300
O4—C20	1.205 (3)	C11—N4	1.326 (3)
C1—N1	1.347 (3)	C11—C12	1.380 (3)
C1—C2	1.358 (3)	C11—H11	0.9300
C1—H1A	0.9300	C12—H12	0.9300
C2—C3	1.381 (3)	C13—C14	1.497 (3)
C2—H2	0.9300	C14—C19	1.384 (3)
C3—C4	1.378 (3)	C14—C15	1.391 (3)
C3—C6	1.457 (3)	C15—C16	1.398 (3)
C4—C5	1.380 (3)	C15—H15	0.9300
C4—H4	0.9300	C16—C17	1.385 (3)
C5—N1	1.343 (3)	C16—C20	1.487 (3)
C5—H5A	0.9300	C17—C18	1.372 (3)
C6—N2	1.287 (3)	C17—H17	0.9300
C6—O6	1.358 (3)	C18—C19	1.372 (3)
C7—N3	1.290 (3)	C18—H18	0.9300
C7—O6	1.366 (3)	C19—H19	0.9300
C7—C8	1.454 (3)	N2—N3	1.400 (3)
C13—O1—H1	109.3	C8—C12—C11	118.4 (2)
C20—O3—H3	109.5	C8—C12—H12	120.8
N1—C1—C2	120.3 (2)	C11—C12—H12	120.8
N1—C1—H1A	119.8	O2—C13—O1	124.3 (3)
C2—C1—H1A	119.8	O2—C13—C14	122.8 (3)
C1—C2—C3	120.7 (2)	O1—C13—C14	112.8 (2)
C1—C2—H2	119.7	C19—C14—C15	119.2 (2)
C3—C2—H2	119.7	C19—C14—C13	118.7 (2)

supplementary materials

C4—C3—C2	118.0 (2)	C15—C14—C13	122.0 (2)
C4—C3—C6	122.1 (2)	C14—C15—C16	120.4 (2)
C2—C3—C6	119.9 (2)	C14—C15—H15	119.8
C3—C4—C5	120.3 (2)	C16—C15—H15	119.8
C3—C4—H4	119.8	C17—C16—C15	118.9 (2)
C5—C4—H4	119.8	C17—C16—C20	119.1 (2)
N1—C5—C4	119.7 (2)	C15—C16—C20	122.0 (2)
N1—C5—H5A	120.1	C18—C17—C16	120.3 (2)
C4—C5—H5A	120.1	C18—C17—H17	119.8
N2—C6—O6	113.3 (2)	C16—C17—H17	119.8
N2—C6—C3	127.5 (2)	C19—C18—C17	120.8 (3)
O6—C6—C3	119.1 (2)	C19—C18—H18	119.6
N3—C7—O6	112.1 (2)	C17—C18—H18	119.6
N3—C7—C8	127.9 (2)	C18—C19—C14	120.2 (2)
O6—C7—C8	120.0 (2)	C18—C19—H19	119.9
C12—C8—C9	118.4 (2)	C14—C19—H19	119.9
C12—C8—C7	122.7 (2)	O4—C20—O3	123.4 (2)
C9—C8—C7	118.8 (2)	O4—C20—C16	123.4 (2)
C10—C9—C8	119.0 (2)	O3—C20—C16	113.2 (2)
C10—C9—H9	120.5	O5—N1—C5	120.9 (2)
C8—C9—H9	120.5	O5—N1—C1	118.2 (2)
N4—C10—C9	123.4 (2)	C5—N1—C1	120.9 (2)
N4—C10—H10	118.3	C6—N2—N3	105.6 (2)
C9—C10—H10	118.3	C7—N3—N2	106.81 (19)
N4—C11—C12	123.7 (2)	C11—N4—C10	117.1 (2)
N4—C11—H11	118.2	C6—O6—C7	102.14 (18)
C12—C11—H11	118.2		
N1—C1—C2—C3	0.2 (4)	C14—C15—C16—C20	179.5 (2)
C1—C2—C3—C4	-0.2 (4)	C15—C16—C17—C18	-0.7 (4)
C1—C2—C3—C6	179.4 (2)	C20—C16—C17—C18	179.9 (2)
C2—C3—C4—C5	0.1 (4)	C16—C17—C18—C19	0.4 (4)
C6—C3—C4—C5	-179.5 (2)	C17—C18—C19—C14	0.6 (4)
C3—C4—C5—N1	0.0 (4)	C15—C14—C19—C18	-1.2 (4)
C4—C3—C6—N2	178.5 (3)	C13—C14—C19—C18	178.7 (2)
C2—C3—C6—N2	-1.1 (4)	C17—C16—C20—O4	3.7 (4)
C4—C3—C6—O6	-2.0 (4)	C15—C16—C20—O4	-175.7 (3)
C2—C3—C6—O6	178.4 (2)	C17—C16—C20—O3	-176.5 (2)
N3—C7—C8—C12	174.8 (2)	C15—C16—C20—O3	4.1 (4)
O6—C7—C8—C12	-4.2 (4)	C4—C5—N1—O5	178.7 (2)
N3—C7—C8—C9	-4.2 (4)	C4—C5—N1—C1	0.0 (4)
O6—C7—C8—C9	176.9 (2)	C2—C1—N1—O5	-178.8 (2)
C12—C8—C9—C10	-1.1 (4)	C2—C1—N1—C5	0.0 (4)
C7—C8—C9—C10	177.9 (2)	O6—C6—N2—N3	0.5 (3)
C8—C9—C10—N4	2.0 (4)	C3—C6—N2—N3	-179.9 (2)
C9—C8—C12—C11	-0.3 (4)	O6—C7—N3—N2	0.4 (3)
C7—C8—C12—C11	-179.3 (2)	C8—C7—N3—N2	-178.6 (2)
N4—C11—C12—C8	1.2 (4)	C6—N2—N3—C7	-0.5 (3)
O2—C13—C14—C19	6.8 (4)	C12—C11—N4—C10	-0.5 (4)
O1—C13—C14—C19	-173.4 (2)	C9—C10—N4—C11	-1.1 (4)

O2—C13—C14—C15	-173.3 (3)	N2—C6—O6—C7	-0.3 (3)
O1—C13—C14—C15	6.5 (4)	C3—C6—O6—C7	-179.9 (2)
C19—C14—C15—C16	0.9 (4)	N3—C7—O6—C6	0.0 (3)
C13—C14—C15—C16	-179.0 (2)	C8—C7—O6—C6	179.0 (2)
C14—C15—C16—C17	0.1 (4)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N4	0.82	1.82	2.640 (2)	171
O3—H3 \cdots O5 ⁱ	0.82	1.72	2.521 (3)	163

Symmetry codes: (i) $x+1, y+1, z+1$.

Fig. 1




Fig. 2




Fig. 3




Fig. 4

