

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dichlorido{2-[2-(piperidin-1-yl)ethyl- iminomethyl]phenolato}zinc(II) monohydrate

 Dong-Fang Zhang,^{a*} Mei-Huan Zhou^b and Chang-Ji Yuan^a

^aSchool of Pharmaceutical Sciences, China Medical University, Shenyang 110001, People's Republic of China, and ^bNortheast Pharmaceutical Group Co. Ltd, Shenyang 110026, People's Republic of China
Correspondence e-mail: Insyzdf@sohu.com


Received 11 May 2008; accepted 13 May 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.045; wR factor = 0.104; data-to-parameter ratio = 19.5.

In the title mononuclear zinc(II) complex, $[\text{ZnCl}_2(\text{C}_{14}\text{H}_{20}\text{N}_2\text{O})]\cdot\text{H}_2\text{O}$, the Zn^{II} atom is four-coordinated by the phenolate O and imine N atoms of the Schiff base ligand and by two Cl atoms in a tetrahedral geometry. In the crystal structure, $\text{O}-\text{H}\cdots\text{Cl}$, $\text{O}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds involving the water molecules bridge adjacent complexes into a ladder-like structure running along the c axis.

Related literature

For general background on Schiff base complexes, see: Kawamoto *et al.* (2008); Tomat *et al.* (2007). For biological properties of Schiff base compounds, see: Abd-Elzaher (2004); Iqbal *et al.* (2005); Osowole *et al.* (2005); Raman & Thangaraja (2005). For related structures, see: Ali *et al.* (2008); Li (2007); Tatar *et al.* (2002); Wang (2007).



Experimental

Crystal data

$[\text{ZnCl}_2(\text{C}_{14}\text{H}_{20}\text{N}_2\text{O})]\cdot\text{H}_2\text{O}$
 $M_r = 386.61$
Monoclinic, $P2_1/c$
 $a = 9.1860$ (18) Å
 $b = 19.875$ (4) Å
 $c = 9.966$ (2) Å
 $\beta = 110.20$ (3)°

$V = 1707.6$ (7) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 1.76$ mm⁻¹
 $T = 298$ (2) K
0.20 × 0.18 × 0.17 mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\text{min}} = 0.720$, $T_{\text{max}} = 0.754$
14151 measured reflections
3882 independent reflections
2685 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.104$
 $S = 0.97$
3882 reflections
199 parameters
4 restraints
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.49$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.58$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Zn1—O1	1.929 (2)	Zn1—Cl2	2.2066 (10)
Zn1—N1	2.024 (2)	Zn1—Cl1	2.2523 (10)
O1—Zn1—N1	95.83 (10)	O1—Zn1—Cl1	109.48 (8)
O1—Zn1—Cl2	113.80 (8)	N1—Zn1—Cl1	109.22 (8)
N1—Zn1—Cl2	111.04 (8)	Cl2—Zn1—Cl1	115.66 (4)

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2C}\cdots\text{O2}$	0.90 (4)	1.81 (4)	2.712 (3)	177 (4)
$\text{O2}-\text{H2B}\cdots\text{O1}^{\text{i}}$	0.84 (3)	1.91 (3)	2.741 (3)	168 (4)
$\text{O2}-\text{H2A}\cdots\text{Cl1}^{\text{ii}}$	0.85 (3)	2.44 (3)	3.272 (3)	168 (4)

 Symmetry codes: (i) $x, y, z - 1$; (ii) $-x + 1, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge China Medical University for support of this research.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2598).

References

- Abd-Elzaher, M. M. (2004). *Appl. Organomet. Chem.* **18**, 149–155.
Ali, H. M., Mohamed Mustafa, M. I., Rizal, M. R. & Ng, S. W. (2008). *Acta Cryst.* **E64**, m718–m719.
Bruker (2000). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Iqbal, M. S., Bukhari, I. H. & Arif, M. (2005). *Appl. Organomet. Chem.* **19**, 864–869.
Kawamoto, T., Nishiwaki, M., Tsunekawa, Y., Nozaki, K. & Konno, T. (2008). *Inorg. Chem.* **47**, 3095–3104.
Li, W.-H. (2007). *Acta Cryst.* **E63**, m1446–m1447.
Osowole, A. A., Kolawole, G. A. & Fagade, O. E. (2005). *Synth. React. Inorg. Met.-Org. Nano-Met. Chem.* **35**, 829–836.

Raman, N. & Thangaraja, C. (2005). *Pol. J. Chem.* **79**, 1123–1134.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Tatar, L., Atakol, O. & Ülkü, D. (2002). *Acta Cryst.* **E58**, m83–m85.

Tomat, E., Cuesta, L., Lynch, V. M. & Sessler, J. L. (2007). *Inorg. Chem.* **46**, 6224–6226.
Wang, S.-X. (2007). *Acta Cryst.* **E63**, m706–m707.

supplementary materials

Acta Cryst. (2008). E64, m825-m826 [doi:10.1107/S1600536808014311]

Dichlorido{2-[2-(piperidin-1-yl)ethyliminomethyl]phenolato}zinc(II) monohydrate

D.-F. Zhang, M.-H. Zhou and C.-J. Yuan

Comment

Zinc(II) complexes with Schiff base ligands have received much attention in recent years (Tomat *et al.*, 2007; Kawamoto *et al.*, 2008). Some of the complexes have been found to have biological properties (Oswole *et al.*, 2005; Iqbal *et al.*, 2005; Raman & Thangaraja, 2005; Abd-Elzaher, 2004). In this paper, the crystal structure of the title new zinc(II) complex with the Schiff base ligand 2-[(2-piperidin-1-ylethylimino)methyl]phenol is reported.

The title compound consists of a mononuclear Schiff base zinc(II) complex molecule and a water of hydration (Fig. 1). The Zn^{II} atom in the complex is four-coordinate in a tetrahedral geometry with one phenolate O and one imine N atoms of the Schiff base ligand, and with two Cl atoms. Bond lengths and angles (Table 1) about the Zn^{II} centre are comparable with the values observed in other Schiff base zinc(II) complexes (Wang, 2007; Ali *et al.*, 2008; Li, 2007; Tatar *et al.*, 2002). The crystal structure is stabilized by intermolecular O—H...Cl, O—H...O and N—H...O hydrogen bonds (Table 2 and Fig. 2).

Experimental

A mixture of salicylaldehyde (0.1 mmol, 12.2 mg), 2-piperidin-1-ylethylamine (0.1 mmol, 12.8 mg) and ZnCl₂ (0.1 mmol, 13.6 mg) in methanol was stirred for 30 min at room temperature to give a yellow solution. After keeping the solution in air for 12 d, yellow block-shaped crystals were formed.

Refinement

Atoms H2A, H2B and H2C were located from a difference Fourier map and refined isotropically, with O—H, N—H, and H...H distances restrained to 0.85 (1), 0.90 (1), and 1.37 (2) Å, respectively. The remaining H atoms were placed in calculated positions and constrained to ride on their parent atoms, with C—H = 0.93 or 0.97 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids.



Fig. 2. Crystal packing of the title compound. Intermolecular hydrogen bonds are shown as dashed lines. H atoms not involved in the interactions have been omitted for clarity.

Dichlorido{2-[2-(piperidin-1-yl)ethyliminomethyl]phenolato}zinc(II) monohydrate

Crystal data

[ZnCl₂(C₁₄H₂₀N₂O)]·H₂O

M_r = 386.61

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2ybc

a = 9.1860 (18) Å

b = 19.875 (4) Å

c = 9.966 (2) Å

β = 110.20 (3)°

V = 1707.6 (7) Å³

Z = 4

*F*₀₀₀ = 800

D_x = 1.504 Mg m⁻³

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 2461 reflections

θ = 2.4–25.0°

μ = 1.76 mm⁻¹

T = 298 (2) K

Block, yellow

0.20 × 0.18 × 0.17 mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 298(2) K

ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2000)

*T*_{min} = 0.720, *T*_{max} = 0.755

14151 measured reflections

3882 independent reflections

2685 reflections with *I* > 2σ(*I*)

*R*_{int} = 0.056

θ_{max} = 27.5°

θ_{min} = 2.1°

h = -11→11

k = -25→25

l = -12→12

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2σ(*F*²)] = 0.045

wR(*F*²) = 0.104

S = 0.97

3882 reflections

199 parameters

4 restraints

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H atoms treated by a mixture of
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0472P)^2]$$

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} = 0.001

Δρ_{max} = 0.49 e Å⁻³

Δρ_{min} = -0.58 e Å⁻³

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.30417 (4)	0.444609 (17)	0.67776 (4)	0.03531 (13)
C11	0.53819 (11)	0.40618 (5)	0.68728 (10)	0.0547 (3)
C12	0.13426 (12)	0.36691 (5)	0.68208 (10)	0.0566 (3)
O1	0.3309 (3)	0.51454 (10)	0.8185 (2)	0.0466 (6)
O2	0.2673 (3)	0.48851 (13)	0.0627 (3)	0.0540 (7)
N1	0.2172 (3)	0.50529 (13)	0.5054 (2)	0.0330 (6)
N2	0.1661 (3)	0.38887 (13)	0.1946 (3)	0.0322 (6)
C6	0.2940 (3)	0.60705 (15)	0.6525 (3)	0.0349 (7)
C1	0.3412 (4)	0.57904 (16)	0.7920 (3)	0.0360 (7)
C2	0.3972 (4)	0.62356 (17)	0.9080 (3)	0.0423 (8)
H2	0.4311	0.6063	1.0004	0.051*
C3	0.4035 (4)	0.69155 (18)	0.8893 (4)	0.0494 (9)
H3	0.4404	0.7197	0.9684	0.059*
C4	0.3549 (4)	0.71867 (18)	0.7520 (4)	0.0553 (10)
H4	0.3586	0.7649	0.7392	0.066*
C5	0.3019 (4)	0.67719 (17)	0.6368 (4)	0.0481 (9)
H5	0.2702	0.6956	0.5454	0.058*
C7	0.2314 (4)	0.56952 (15)	0.5204 (3)	0.0351 (7)
H7	0.1976	0.5948	0.4367	0.042*
C8	0.1417 (4)	0.48009 (16)	0.3607 (3)	0.0420 (8)
H8A	0.1519	0.5126	0.2919	0.050*
H8B	0.0321	0.4730	0.3428	0.050*
C9	0.2177 (4)	0.41441 (16)	0.3450 (3)	0.0379 (8)
H9A	0.3293	0.4206	0.3789	0.045*
H9B	0.1945	0.3807	0.4051	0.045*
C10	0.2450 (4)	0.32330 (17)	0.1916 (3)	0.0414 (8)
H10A	0.2136	0.2904	0.2482	0.050*
H10B	0.3564	0.3292	0.2339	0.050*
C11	0.2048 (4)	0.29752 (17)	0.0403 (3)	0.0454 (9)
H11A	0.2538	0.2542	0.0419	0.054*
H11B	0.2446	0.3284	-0.0141	0.054*
C12	0.0304 (4)	0.29045 (18)	-0.0318 (4)	0.0507 (9)
H12A	0.0064	0.2773	-0.1308	0.061*

supplementary materials

H12B	-0.0082	0.2558	0.0158	0.061*
C13	-0.0471 (4)	0.35675 (18)	-0.0247 (3)	0.0476 (9)
H13A	-0.1587	0.3514	-0.0667	0.057*
H13B	-0.0156	0.3900	-0.0805	0.057*
C14	-0.0063 (4)	0.38173 (17)	0.1267 (3)	0.0422 (8)
H14A	-0.0555	0.4249	0.1267	0.051*
H14B	-0.0442	0.3503	0.1814	0.051*
H2C	0.198 (5)	0.4214 (14)	0.148 (4)	0.080*
H2B	0.295 (4)	0.492 (2)	-0.009 (2)	0.080*
H2A	0.330 (4)	0.5124 (18)	0.128 (3)	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0448 (3)	0.0329 (2)	0.0281 (2)	0.00015 (17)	0.01246 (17)	0.00067 (15)
Cl1	0.0473 (6)	0.0665 (6)	0.0490 (5)	0.0112 (5)	0.0150 (4)	0.0005 (5)
Cl2	0.0635 (6)	0.0507 (5)	0.0599 (6)	-0.0155 (5)	0.0266 (5)	0.0007 (4)
O1	0.0800 (18)	0.0341 (13)	0.0296 (12)	-0.0072 (12)	0.0238 (12)	-0.0028 (10)
O2	0.0692 (19)	0.0562 (16)	0.0411 (14)	-0.0195 (13)	0.0248 (14)	-0.0026 (12)
N1	0.0398 (16)	0.0353 (14)	0.0255 (13)	0.0008 (12)	0.0132 (12)	-0.0010 (11)
N2	0.0331 (15)	0.0381 (14)	0.0252 (13)	0.0002 (12)	0.0099 (11)	0.0004 (11)
C6	0.0319 (18)	0.0368 (17)	0.0374 (18)	0.0024 (14)	0.0138 (15)	0.0022 (14)
C1	0.0355 (19)	0.0385 (17)	0.0386 (19)	0.0004 (14)	0.0187 (15)	-0.0065 (14)
C2	0.046 (2)	0.047 (2)	0.0370 (19)	-0.0021 (16)	0.0183 (16)	-0.0113 (15)
C3	0.037 (2)	0.045 (2)	0.068 (3)	-0.0056 (16)	0.0200 (19)	-0.0248 (19)
C4	0.050 (2)	0.0339 (19)	0.080 (3)	0.0040 (17)	0.020 (2)	-0.0045 (19)
C5	0.049 (2)	0.0342 (18)	0.060 (2)	0.0049 (16)	0.0171 (19)	0.0042 (17)
C7	0.0336 (18)	0.0404 (19)	0.0336 (18)	0.0073 (14)	0.0145 (15)	0.0088 (14)
C8	0.047 (2)	0.048 (2)	0.0294 (17)	0.0098 (17)	0.0113 (16)	0.0019 (15)
C9	0.043 (2)	0.0495 (19)	0.0190 (16)	0.0072 (16)	0.0082 (14)	-0.0001 (13)
C10	0.039 (2)	0.0447 (19)	0.0378 (19)	0.0078 (16)	0.0102 (16)	0.0010 (15)
C11	0.045 (2)	0.045 (2)	0.045 (2)	0.0031 (16)	0.0135 (17)	-0.0076 (16)
C12	0.049 (2)	0.053 (2)	0.047 (2)	-0.0069 (18)	0.0124 (18)	-0.0116 (17)
C13	0.036 (2)	0.063 (2)	0.0363 (19)	-0.0028 (17)	0.0029 (16)	-0.0081 (17)
C14	0.0313 (19)	0.050 (2)	0.041 (2)	0.0048 (16)	0.0077 (15)	-0.0008 (16)

Geometric parameters (\AA , $^\circ$)

Zn1—O1	1.929 (2)	C4—H4	0.93
Zn1—N1	2.024 (2)	C5—H5	0.93
Zn1—Cl2	2.2066 (10)	C7—H7	0.93
Zn1—Cl1	2.2523 (10)	C8—C9	1.514 (4)
O1—C1	1.319 (4)	C8—H8A	0.97
O2—H2B	0.84 (3)	C8—H8B	0.97
O2—H2A	0.85 (3)	C9—H9A	0.97
N1—C7	1.287 (4)	C9—H9B	0.97
N1—C8	1.457 (4)	C10—C11	1.512 (4)
N2—C9	1.496 (4)	C10—H10A	0.97
N2—C10	1.496 (4)	C10—H10B	0.97

N2—C14	1.499 (4)	C11—C12	1.520 (5)
N2—H2C	0.90 (4)	C11—H11A	0.97
C6—C5	1.407 (4)	C11—H11B	0.97
C6—C1	1.420 (4)	C12—C13	1.511 (5)
C6—C7	1.448 (4)	C12—H12A	0.97
C1—C2	1.404 (4)	C12—H12B	0.97
C2—C3	1.368 (5)	C13—C14	1.507 (4)
C2—H2	0.93	C13—H13A	0.97
C3—C4	1.393 (5)	C13—H13B	0.97
C3—H3	0.93	C14—H14A	0.97
C4—C5	1.360 (5)	C14—H14B	0.97
O1—Zn1—N1	95.83 (10)	C9—C8—H8A	110.0
O1—Zn1—C12	113.80 (8)	N1—C8—H8B	110.0
N1—Zn1—C12	111.04 (8)	C9—C8—H8B	110.0
O1—Zn1—C11	109.48 (8)	H8A—C8—H8B	108.3
N1—Zn1—C11	109.22 (8)	N2—C9—C8	113.5 (2)
C12—Zn1—C11	115.66 (4)	N2—C9—H9A	108.9
C1—O1—Zn1	123.71 (18)	C8—C9—H9A	108.9
H2B—O2—H2A	106 (2)	N2—C9—H9B	108.9
C7—N1—C8	116.8 (3)	C8—C9—H9B	108.9
C7—N1—Zn1	119.9 (2)	H9A—C9—H9B	107.7
C8—N1—Zn1	123.31 (19)	N2—C10—C11	111.2 (2)
C9—N2—C10	109.1 (2)	N2—C10—H10A	109.4
C9—N2—C14	113.8 (2)	C11—C10—H10A	109.4
C10—N2—C14	110.6 (2)	N2—C10—H10B	109.4
C9—N2—H2C	103 (3)	C11—C10—H10B	109.4
C10—N2—H2C	112 (3)	H10A—C10—H10B	108.0
C14—N2—H2C	108 (3)	C10—C11—C12	110.9 (3)
C5—C6—C1	119.1 (3)	C10—C11—H11A	109.4
C5—C6—C7	115.4 (3)	C12—C11—H11A	109.4
C1—C6—C7	125.4 (3)	C10—C11—H11B	109.4
O1—C1—C2	118.6 (3)	C12—C11—H11B	109.4
O1—C1—C6	123.9 (3)	H11A—C11—H11B	108.0
C2—C1—C6	117.4 (3)	C13—C12—C11	109.5 (3)
C3—C2—C1	122.1 (3)	C13—C12—H12A	109.8
C3—C2—H2	119.0	C11—C12—H12A	109.8
C1—C2—H2	119.0	C13—C12—H12B	109.8
C2—C3—C4	120.1 (3)	C11—C12—H12B	109.8
C2—C3—H3	120.0	H12A—C12—H12B	108.2
C4—C3—H3	120.0	C12—C13—C14	112.1 (3)
C5—C4—C3	119.7 (3)	C12—C13—H13A	109.2
C5—C4—H4	120.2	C14—C13—H13A	109.2
C3—C4—H4	120.2	C12—C13—H13B	109.2
C4—C5—C6	121.6 (3)	C14—C13—H13B	109.2
C4—C5—H5	119.2	H13A—C13—H13B	107.9
C6—C5—H5	119.2	N2—C14—C13	110.0 (3)
N1—C7—C6	127.5 (3)	N2—C14—H14A	109.7
N1—C7—H7	116.2	C13—C14—H14A	109.7
C6—C7—H7	116.2	N2—C14—H14B	109.7

supplementary materials

N1—C8—C9	108.6 (2)	C13—C14—H14B	109.7
N1—C8—H8A	110.0	H14A—C14—H14B	108.2

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N2—H2C···O2	0.90 (4)	1.81 (4)	2.712 (3)	177 (4)
O2—H2B···O1 ⁱ	0.84 (3)	1.91 (3)	2.741 (3)	168 (4)
O2—H2A···C11 ⁱⁱ	0.85 (3)	2.44 (3)	3.272 (3)	168 (4)

Symmetry codes: (i) $x, y, z-1$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

