

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dichloridobis[(1*S*,1*S'*,2*R*,2*R'*)-(+)-1,1'-di-*tert*-butyl-2,2'-diphospholane- κ^2 P,P']-ruthenium(II)

Chubei Wang,^a Haiyan Tao^{b*} and Baoming Ji^c

^aCollege of Chemistry, Central China Normal University, Wuhan, Hubei 430072, People's Republic of China, ^bCollege of Chemistry and Molecular Science, Wuhan University, Wuhan, Hubei 430072, People's Republic of China, and ^cCollege of Chemistry and Chemical Engineering, Luoyang Normal University, Luoyang, Henan 471022, People's Republic of China

Correspondence e-mail: haiyantao21@hotmail.com


Received 29 February 2008; accepted 27 March 2008

Key indicators: single-crystal X-ray study; $T = 108$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.024; wR factor = 0.058; data-to-parameter ratio = 22.8.

In the title compound, $[\text{RuCl}_2(\text{C}_{16}\text{H}_{32}\text{P}_2)_2]$, the Ru^{II} ion is situated on a twofold rotation axis, so the asymmetric unit contains one half-molecule. The slightly distorted octahedral environment of the Ru center is formed by four P atoms [$\text{Ru}-\text{P} = 2.4417$ (6) and 2.4544 (6) Å] from two different (1*S*,1*S'*,2*R*,2*R'*)-TangPhos ligands [(1*S*,1*S'*,2*R*,2*R'*)-TangPhos = (1*S*,1*S'*,2*R*,2*R'*)-(+)-1,1'-di-*tert*-butyl-2,2'-diphospholane] and two Cl atoms [$\text{Ru}-\text{Cl} = 2.4267$ (5) Å].

Related literature

For related literature, see: Ikariya *et al.* (1985); James & Fogg (1993); Stoop *et al.* (1999).



Experimental

Crystal data

 $[\text{RuCl}_2(\text{C}_{16}\text{H}_{32}\text{P}_2)_2]$ $M_r = 744.72$ Orthorhombic, $C222_1$ $a = 11.8640$ (14) Å $b = 20.669$ (3) Å $c = 14.2274$ (17) Å $V = 3488.8$ (8) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.81$ mm⁻¹ $T = 108$ (2) K $0.24 \times 0.15 \times 0.10$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan

(SADABS; Bruker, 2001)

 $T_{\min} = 0.863$, $T_{\max} = 0.921$

11542 measured reflections

4164 independent reflections

4018 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.024$ $wR(F^2) = 0.057$ $S = 1.06$

4164 reflections

183 parameters

H-atom parameters not refined

 $\Delta\rho_{\text{max}} = 0.83$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.29$ e Å⁻³

Absolute structure: Flack (1983),

1750 Friedel pairs

Flack parameter: 0.00 (2)

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the National Natural Science Foundation of China (grant No. 20702039).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2390).

References

- Bruker (1997). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2001). *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Ikariya, T., Ishii, Y., Kawano, H., Arai, T., Saburi, M., Yoshikawa, S. & Akutagawa, S. (1985). *J. Chem. Soc. Chem. Commun.* pp. 922–924.
 James, B. R. & Fogg, D. E. (1993). *J. Organomet. Chem.* **462**, c21–c24.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stoop, R. M., Bauer, C., Setz, P., Wörle, M., Wong, T. Y. H. & Mezzetti, A. (1999). *Organometallics*, **18**, 5691–5700.

supplementary materials

Acta Cryst. (2008). E64, m754 [doi:10.1107/S1600536808008301]

Dichloridobis[(1*S*,1*S'*,2*R*,2*R'*)-(+)-1,1'-di-*tert*-butyl-2,2'-diphospholane- κ^2 P,P']ruthenium(II)

C. Wang, H. Tao and B. Ji

Comment

Recently, some chiral diphosphino ruthenium complexes have been synthesized and used as catalysts for the asymmetric reactions (Stoop *et al.*, 1999; James *et al.*, 1993). Herein, we report the synthesis and crystal structure of the title compound - the ruthenium(II) complex containing the chiral TangPhos ligand.

As shown in Fig. 1, the crystals of the title complex contain discrete $[\text{RuCl}_2(\text{TangPhos})_2]$ units with the metal center in a slightly distorted octahedral environment. The *trans*-axial positions of Ru^{II} environment are occupied by Cl^{II} and Cl^I atoms, and the equatorial positions are occupied by P1, P2, P1ⁱ, P2ⁱ atoms, respectively [symmetry code: (i) $-x, y, -z + 1/2$], from two different TangPhos ligands, resulting in two chelate rings, which assume a half-chair conformation with the *tert*-butyl in the less hindered equatorial positions. Similar conformations were found in previously reported related structures (Stoop *et al.*, 1999; Ikariya *et al.*, 1985).

Experimental

To a solution of $[\text{RuCl}_2(\text{PPh})_3]$ (96 mg, 0.1 mmol) in 2 ml of deoxygenated CH_2Cl_2 was added dropwise a solution of (1*S*,1*S'*,2*R*,2*R'*)-TangPhos, (56 mg, 0.2 mmol) in CH_2Cl_2 (1 ml). The resulting mixture was stirred at ambient temperature for 6 h. Deoxygenated ether (12 ml) was added into the vigorously stirring solution and kept for 6 h at room temperature. The resulting light brown precipitate was filtered, washed with ether (3 times with 10 ml), and dried under vacuum. Yield: 45 mg (56%). Crystals suitable for X-ray diffraction were obtained by diffusion of hexane into a CD_2Cl_2 solution of the above compound at room temperature.

Refinement

All H atoms were positioned geometrically (C—H 0.98–1.00 Å), and treated as riding, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. View of the title compound with the atomic numbering and 40% probability displacement ellipsoids [symmetry code: (i) $-x, y, -z + 1/2$]. H atoms are omitted for clarity.

Dichloridobis[(1*S*,1*S'*,2*R*,2*R'*)-(+)-1,1'-di-*tert*-butyl-2,2'-diphospholane- κ^2P,P']ruthenium(II)

Crystal data

[RuCl ₂ (C ₁₆ H ₃₂ P ₂) ₂]	$F_{000} = 1576$
$M_r = 744.72$	$D_x = 1.418 \text{ Mg m}^{-3}$
Orthorhombic, $C222_1$	Mo $K\alpha$ radiation
Hall symbol: C 2c 2	$\lambda = 0.71073 \text{ \AA}$
$a = 11.8640 (14) \text{ \AA}$	Cell parameters from 5992 reflections
$b = 20.669 (3) \text{ \AA}$	$\theta = 2.4\text{--}28.2^\circ$
$c = 14.2274 (17) \text{ \AA}$	$\mu = 0.81 \text{ mm}^{-1}$
$V = 3488.8 (8) \text{ \AA}^3$	$T = 108 (2) \text{ K}$
$Z = 4$	Brick, orange
	$0.24 \times 0.15 \times 0.10 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	4164 independent reflections
Radiation source: fine-focus sealed tube	4018 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.025$
$T = 108(2) \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
phi and ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2001)	$h = -14 \rightarrow 15$
$T_{\text{min}} = 0.863$, $T_{\text{max}} = 0.921$	$k = -17 \rightarrow 27$
11542 measured reflections	$l = -18 \rightarrow 18$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters not refined
$R[F^2 > 2\sigma(F^2)] = 0.024$	$w = 1/[\sigma^2(F_o^2) + (0.0315P)^2 + 0.2939P]$
$wR(F^2) = 0.057$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} = 0.003$
4164 reflections	$\Delta\rho_{\text{max}} = 0.83 \text{ e \AA}^{-3}$
183 parameters	$\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1750 Friedel pairs
	Flack parameter: 0.00 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.02414 (19)	0.16346 (11)	-0.00952 (15)	0.0256 (5)
H1A	-0.0059	0.1290	-0.0497	0.038*
H1B	0.0806	0.1883	-0.0445	0.038*
H1C	-0.0373	0.1922	0.0098	0.038*
C2	-0.0038 (3)	0.08613 (11)	0.12059 (16)	0.0353 (6)
H2A	-0.0740	0.1086	0.1364	0.053*
H2B	0.0291	0.0677	0.1778	0.053*
H2C	-0.0198	0.0514	0.0756	0.053*
C3	0.1820 (2)	0.09489 (11)	0.04344 (16)	0.0257 (5)
H3A	0.2177	0.0737	0.0973	0.039*
H3B	0.2361	0.1243	0.0136	0.039*
H3C	0.1580	0.0621	-0.0021	0.039*
C4	0.07852 (18)	0.13369 (11)	0.07717 (15)	0.0188 (4)
C5	0.21860 (15)	0.15194 (10)	0.24348 (17)	0.0185 (4)
H5A	0.2097	0.1649	0.3100	0.022*
H5B	0.2031	0.1050	0.2384	0.022*
C6	0.33823 (17)	0.16652 (12)	0.21003 (18)	0.0262 (5)
H6A	0.3927	0.1616	0.2622	0.031*
H6B	0.3601	0.1369	0.1584	0.031*
C7	0.33514 (17)	0.23639 (12)	0.17560 (17)	0.0261 (5)
H7A	0.4066	0.2474	0.1434	0.031*
H7B	0.3250	0.2663	0.2293	0.031*
C8	0.23545 (18)	0.24229 (10)	0.10707 (15)	0.0179 (4)
H8	0.2549	0.2176	0.0489	0.021*
C9	0.20454 (18)	0.31083 (11)	0.07802 (15)	0.0185 (4)
H9	0.2760	0.3358	0.0695	0.022*
C10	0.13653 (19)	0.31551 (11)	-0.01378 (14)	0.0220 (5)
H10A	0.1839	0.3027	-0.0680	0.026*
H10B	0.0702	0.2865	-0.0112	0.026*
C11	0.09906 (19)	0.38550 (11)	-0.02376 (15)	0.0237 (5)
H11A	0.0465	0.3904	-0.0773	0.028*
H11B	0.1647	0.4142	-0.0338	0.028*
C12	0.04012 (19)	0.40157 (10)	0.06914 (15)	0.0203 (5)

supplementary materials

H12A	-0.0402	0.3887	0.0663	0.024*
H12B	0.0442	0.4486	0.0817	0.024*
C13	0.2153 (2)	0.42001 (11)	0.21194 (17)	0.0261 (5)
C14	0.3123 (2)	0.39022 (14)	0.26745 (19)	0.0414 (7)
H14A	0.3582	0.4247	0.2951	0.062*
H14B	0.3590	0.3640	0.2253	0.062*
H14C	0.2819	0.3628	0.3176	0.062*
C15	0.2678 (2)	0.46048 (11)	0.13264 (17)	0.0291 (5)
H15A	0.2078	0.4806	0.0954	0.044*
H15B	0.3134	0.4324	0.0921	0.044*
H15C	0.3158	0.4943	0.1598	0.044*
C16	0.1502 (3)	0.46594 (13)	0.2754 (2)	0.0505 (9)
H16A	0.1180	0.4417	0.3281	0.076*
H16B	0.0895	0.4864	0.2394	0.076*
H16C	0.2013	0.4993	0.2995	0.076*
Cl1	-0.13351 (4)	0.27771 (3)	0.12078 (3)	0.01700 (10)
P1	0.11419 (5)	0.35562 (3)	0.16379 (4)	0.01581 (11)
P2	0.11916 (4)	0.19819 (3)	0.16823 (4)	0.01347 (11)
Ru1	0.0000	0.277275 (10)	0.2500	0.01108 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0294 (13)	0.0265 (11)	0.0207 (11)	0.0048 (10)	-0.0060 (9)	-0.0074 (9)
C2	0.0464 (14)	0.0316 (12)	0.0279 (12)	-0.0175 (14)	0.0162 (14)	-0.0125 (10)
C3	0.0352 (13)	0.0209 (11)	0.0209 (11)	0.0099 (10)	0.0033 (10)	-0.0011 (9)
C4	0.0241 (11)	0.0182 (10)	0.0142 (10)	0.0018 (9)	0.0029 (8)	-0.0020 (8)
C5	0.0201 (10)	0.0212 (9)	0.0141 (9)	0.0048 (8)	0.0021 (9)	0.0025 (10)
C6	0.0168 (11)	0.0359 (14)	0.0260 (11)	0.0088 (10)	0.0036 (9)	0.0107 (10)
C7	0.0132 (10)	0.0342 (15)	0.0310 (12)	-0.0006 (9)	0.0012 (9)	0.0117 (10)
C8	0.0166 (10)	0.0211 (11)	0.0159 (9)	0.0012 (8)	0.0046 (8)	0.0031 (8)
C9	0.0181 (10)	0.0208 (11)	0.0166 (10)	0.0011 (8)	0.0045 (8)	0.0041 (9)
C10	0.0285 (12)	0.0242 (12)	0.0133 (10)	0.0052 (10)	0.0058 (9)	0.0023 (8)
C11	0.0304 (13)	0.0240 (12)	0.0166 (11)	0.0015 (10)	0.0039 (9)	0.0061 (9)
C12	0.0247 (11)	0.0162 (10)	0.0199 (11)	0.0009 (9)	0.0015 (8)	0.0037 (9)
C13	0.0326 (13)	0.0224 (12)	0.0234 (11)	-0.0126 (10)	0.0024 (10)	-0.0004 (10)
C14	0.0485 (16)	0.0439 (15)	0.0317 (17)	-0.0285 (13)	-0.0166 (12)	0.0120 (12)
C15	0.0348 (13)	0.0212 (11)	0.0315 (13)	-0.0111 (11)	0.0056 (11)	0.0024 (10)
C16	0.0582 (19)	0.0429 (17)	0.050 (2)	-0.0304 (15)	0.0255 (15)	-0.0289 (14)
Cl1	0.0168 (2)	0.0205 (2)	0.0137 (2)	0.0003 (2)	-0.00355 (16)	0.0000 (2)
P1	0.0188 (3)	0.0144 (3)	0.0143 (3)	-0.0034 (2)	0.0013 (2)	0.0013 (2)
P2	0.0146 (3)	0.0147 (2)	0.0111 (2)	0.0011 (2)	0.0020 (2)	0.0013 (2)
Ru1	0.01224 (10)	0.01165 (10)	0.00935 (9)	0.000	0.00015 (8)	0.000

Geometric parameters (\AA , $^\circ$)

C1—C4	1.522 (3)	C10—C11	1.520 (3)
C1—H1A	0.9800	C10—H10A	0.9900
C1—H1B	0.9800	C10—H10B	0.9900

C1—H1C	0.9800	C11—C12	1.532 (3)
C2—C4	1.517 (3)	C11—H11A	0.9900
C2—H2A	0.9800	C11—H11B	0.9900
C2—H2B	0.9800	C12—P1	1.868 (2)
C2—H2C	0.9800	C12—H12A	0.9900
C3—C4	1.543 (3)	C12—H12B	0.9900
C3—H3A	0.9800	C13—C16	1.520 (4)
C3—H3B	0.9800	C13—C14	1.525 (3)
C3—H3C	0.9800	C13—C15	1.537 (3)
C4—P2	1.920 (2)	C13—P1	1.918 (2)
C5—C6	1.527 (3)	C14—H14A	0.9800
C5—P2	1.858 (2)	C14—H14B	0.9800
C5—H5A	0.9900	C14—H14C	0.9800
C5—H5B	0.9900	C15—H15A	0.9800
C6—C7	1.525 (3)	C15—H15B	0.9800
C6—H6A	0.9900	C15—H15C	0.9800
C6—H6B	0.9900	C16—H16A	0.9800
C7—C8	1.538 (3)	C16—H16B	0.9800
C7—H7A	0.9900	C16—H16C	0.9800
C7—H7B	0.9900	Cl1—Ru1	2.4267 (5)
C8—C9	1.521 (3)	P1—Ru1	2.4417 (6)
C8—P2	1.869 (2)	P2—Ru1	2.4544 (6)
C8—H8	1.0000	Ru1—Cl1 ⁱ	2.4267 (5)
C9—C10	1.538 (3)	Ru1—P1 ⁱ	2.4417 (6)
C9—P1	1.870 (2)	Ru1—P2 ⁱ	2.4544 (6)
C9—H9	1.0000		
C4—C1—H1A	109.5	C12—C11—H11A	110.7
C4—C1—H1B	109.5	C10—C11—H11B	110.7
H1A—C1—H1B	109.5	C12—C11—H11B	110.7
C4—C1—H1C	109.5	H11A—C11—H11B	108.8
H1A—C1—H1C	109.5	C11—C12—P1	107.29 (15)
H1B—C1—H1C	109.5	C11—C12—H12A	110.3
C4—C2—H2A	109.5	P1—C12—H12A	110.3
C4—C2—H2B	109.5	C11—C12—H12B	110.3
H2A—C2—H2B	109.5	P1—C12—H12B	110.3
C4—C2—H2C	109.5	H12A—C12—H12B	108.5
H2A—C2—H2C	109.5	C16—C13—C14	109.1 (2)
H2B—C2—H2C	109.5	C16—C13—C15	107.6 (2)
C4—C3—H3A	109.5	C14—C13—C15	107.1 (2)
C4—C3—H3B	109.5	C16—C13—P1	109.12 (17)
H3A—C3—H3B	109.5	C14—C13—P1	112.12 (17)
C4—C3—H3C	109.5	C15—C13—P1	111.66 (16)
H3A—C3—H3C	109.5	C13—C14—H14A	109.5
H3B—C3—H3C	109.5	C13—C14—H14B	109.5
C2—C4—C1	108.6 (2)	H14A—C14—H14B	109.5
C2—C4—C3	107.60 (19)	C13—C14—H14C	109.5
C1—C4—C3	107.18 (17)	H14A—C14—H14C	109.5
C2—C4—P2	109.67 (14)	H14B—C14—H14C	109.5

supplementary materials

C1—C4—P2	111.87 (15)	C13—C15—H15A	109.5
C3—C4—P2	111.76 (15)	C13—C15—H15B	109.5
C6—C5—P2	108.00 (15)	H15A—C15—H15B	109.5
C6—C5—H5A	110.1	C13—C15—H15C	109.5
P2—C5—H5A	110.1	H15A—C15—H15C	109.5
C6—C5—H5B	110.1	H15B—C15—H15C	109.5
P2—C5—H5B	110.1	C13—C16—H16A	109.5
H5A—C5—H5B	108.4	C13—C16—H16B	109.5
C7—C6—C5	105.35 (17)	H16A—C16—H16B	109.5
C7—C6—H6A	110.7	C13—C16—H16C	109.5
C5—C6—H6A	110.7	H16A—C16—H16C	109.5
C7—C6—H6B	110.7	H16B—C16—H16C	109.5
C5—C6—H6B	110.7	C12—P1—C9	92.92 (10)
H6A—C6—H6B	108.8	C12—P1—C13	101.48 (10)
C6—C7—C8	107.28 (19)	C9—P1—C13	102.59 (10)
C6—C7—H7A	110.3	C12—P1—Ru1	116.00 (7)
C8—C7—H7A	110.3	C9—P1—Ru1	108.52 (7)
C6—C7—H7B	110.3	C13—P1—Ru1	128.89 (8)
C8—C7—H7B	110.3	C5—P2—C8	92.89 (9)
H7A—C7—H7B	108.5	C5—P2—C4	101.01 (10)
C9—C8—C7	115.57 (18)	C8—P2—C4	102.11 (9)
C9—C8—P2	113.76 (15)	C5—P2—Ru1	115.80 (8)
C7—C8—P2	103.53 (14)	C8—P2—Ru1	108.73 (7)
C9—C8—H8	107.9	C4—P2—Ru1	129.60 (7)
C7—C8—H8	107.9	C11 ⁱ —Ru1—Cl1	179.58 (3)
P2—C8—H8	107.9	C11 ⁱ —Ru1—P1	90.913 (19)
C8—C9—C10	114.56 (18)	C11—Ru1—P1	88.806 (19)
C8—C9—P1	114.98 (15)	C11 ⁱ —Ru1—P1 ⁱ	88.806 (19)
C10—C9—P1	102.85 (14)	C11—Ru1—P1 ⁱ	90.913 (19)
C8—C9—H9	108.0	P1—Ru1—P1 ⁱ	96.91 (3)
C10—C9—H9	108.0	C11 ⁱ —Ru1—P2 ⁱ	91.107 (19)
P1—C9—H9	108.0	C11—Ru1—P2 ⁱ	89.175 (19)
C11—C10—C9	107.00 (18)	P1—Ru1—P2 ⁱ	177.969 (19)
C11—C10—H10A	110.3	P1 ⁱ —Ru1—P2 ⁱ	83.341 (18)
C9—C10—H10A	110.3	C11 ⁱ —Ru1—P2	89.175 (19)
C11—C10—H10B	110.3	C11—Ru1—P2	91.107 (19)
C9—C10—H10B	110.3	P1—Ru1—P2	83.341 (18)
H10A—C10—H10B	108.6	P1 ⁱ —Ru1—P2	177.969 (19)
C10—C11—C12	105.02 (17)	P2 ⁱ —Ru1—P2	96.48 (3)
C10—C11—H11A	110.7		
P2—C5—C6—C7	33.0 (2)	C9—C8—P2—C4	110.84 (16)
C5—C6—C7—C8	-50.9 (2)	C7—C8—P2—C4	-122.93 (15)
C6—C7—C8—C9	169.48 (18)	C9—C8—P2—Ru1	-28.74 (16)
C6—C7—C8—P2	44.4 (2)	C7—C8—P2—Ru1	97.50 (14)
C7—C8—C9—C10	159.29 (18)	C2—C4—P2—C5	76.02 (19)
P2—C8—C9—C10	-81.1 (2)	C1—C4—P2—C5	-163.42 (15)

C7—C8—C9—P1	-81.9 (2)	C3—C4—P2—C5	-43.22 (17)
P2—C8—C9—P1	37.76 (19)	C2—C4—P2—C8	171.43 (17)
C8—C9—C10—C11	171.88 (18)	C1—C4—P2—C8	-68.01 (17)
P1—C9—C10—C11	46.39 (19)	C3—C4—P2—C8	52.19 (17)
C9—C10—C11—C12	-53.0 (2)	C2—C4—P2—Ru1	-61.4 (2)
C10—C11—C12—P1	33.8 (2)	C1—C4—P2—Ru1	59.14 (18)
C11—C12—P1—C9	-6.50 (16)	C3—C4—P2—Ru1	179.34 (11)
C11—C12—P1—C13	96.98 (16)	C12—P1—Ru1—Cl1 ⁱ	-160.02 (8)
C11—C12—P1—Ru1	-118.75 (14)	C9—P1—Ru1—Cl1 ⁱ	97.09 (7)
C8—C9—P1—C12	-147.28 (16)	C13—P1—Ru1—Cl1 ⁱ	-27.34 (10)
C10—C9—P1—C12	-22.07 (15)	C12—P1—Ru1—Cl1	19.66 (8)
C8—C9—P1—C13	110.27 (16)	C9—P1—Ru1—Cl1	-83.23 (7)
C10—C9—P1—C13	-124.52 (15)	C13—P1—Ru1—Cl1	152.34 (10)
C8—C9—P1—Ru1	-28.60 (16)	C12—P1—Ru1—P1 ⁱ	-71.11 (8)
C10—C9—P1—Ru1	96.61 (13)	C9—P1—Ru1—P1 ⁱ	-174.00 (8)
C16—C13—P1—C12	77.9 (2)	C13—P1—Ru1—P1 ⁱ	61.56 (10)
C14—C13—P1—C12	-161.05 (17)	C12—P1—Ru1—P2	110.92 (8)
C15—C13—P1—C12	-40.84 (19)	C9—P1—Ru1—P2	8.03 (7)
C16—C13—P1—C9	173.58 (18)	C13—P1—Ru1—P2	-116.41 (10)
C14—C13—P1—C9	-65.40 (19)	C5—P2—Ru1—Cl1 ⁱ	20.56 (7)
C15—C13—P1—C9	54.81 (19)	C8—P2—Ru1—Cl1 ⁱ	-82.33 (7)
C16—C13—P1—Ru1	-59.7 (2)	C4—P2—Ru1—Cl1 ⁱ	153.05 (9)
C14—C13—P1—Ru1	61.3 (2)	C5—P2—Ru1—Cl1	-159.76 (7)
C15—C13—P1—Ru1	-178.44 (13)	C8—P2—Ru1—Cl1	97.35 (7)
C6—C5—P2—C8	-6.72 (17)	C4—P2—Ru1—Cl1	-27.27 (9)
C6—C5—P2—C4	96.21 (17)	C8—P2—Ru1—P1	8.68 (7)
C6—C5—P2—Ru1	-119.15 (15)	C5—P2—Ru1—P2 ⁱ	-70.46 (7)
C9—C8—P2—C5	-147.25 (16)	C8—P2—Ru1—P2 ⁱ	-173.36 (8)
C7—C8—P2—C5	-21.02 (16)	C4—P2—Ru1—P2 ⁱ	62.03 (9)

Symmetry codes: (i) $-x, y, -z+1/2$.

Fig. 1

