

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bromido-1 κ Br-tricarbonyl-2 κ^3 C-(2 η^5 -cyclopentadienyl)molybdenum(I)-tungsten(I)(W—Mo)

Martin O. Onani,^a Jan-André Gertenbach^b and Muhammad D. Bala^{c*}

^aUniversity of the Western Cape, Modderdam Road, Bellville, Cape Town 7535, South Africa, ^bDepartment of Chemistry and Polymer Science, University of Stellenbosch, Private Bag X1, Matieland 7602, South Africa, and ^cSchool of Chemistry, University of KwaZulu-Natal, Westville Campus, Private Bag X54001, Durban 4000, South Africa

Correspondence e-mail: bala@ukzn.ac.za


Received 23 April 2008; accepted 30 April 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.017$ Å; R factor = 0.031; wR factor = 0.069; data-to-parameter ratio = 21.0.

The title compound, $[\text{WMoBr}(\text{C}_5\text{H}_5)(\text{CO})_3]$, is built up from a pseudo-square-pyramidal piano-stool coordination around the Mo atom, the important geometry being Mo—W = 2.6872 (7) Å, W—Br = 2.5591 (9) Å and Mo—W—Br = 158.35 (3)°.

Related literature

For related literature, see Albright *et al.* (1978); Bueno & Churchill (1981); Changamu *et al.* (2006); Friedrich *et al.* (2004).



Experimental

Crystal data

$[\text{WMoBr}(\text{C}_5\text{H}_5)(\text{CO})_3]$
 $M_r = 508.82$
 Tetragonal, $P4_21c$
 $a = 11.9375$ (9) Å
 $c = 15.546$ (2) Å
 $V = 2215.4$ (4) Å³

$Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 15.09$ mm⁻¹
 $T = 100$ (2) K
 $0.11 \times 0.10 \times 0.07$ mm

Data collection

Bruker APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2002)
 $T_{\min} = 0.251$, $T_{\max} = 0.347$

13298 measured reflections
 2673 independent reflections
 2497 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.069$
 $S = 1.02$
 2673 reflections
 127 parameters
 H-atom parameters constrained

$\Delta\rho_{\text{max}} = 1.31$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.72$ e Å⁻³
 Absolute structure: Flack (1983),
 1118 Friedel pairs
 Flack parameter: 0.00 (1)

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2003).

The authors are grateful to the NRF, WSU and UWC for funding, and to Miss Lungelwa Dyantyi for assistance with the experimental work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2343).

References

- Albright, M. J., Glick, D. M. & Oliver, J. P. (1978). *J. Organomet. Chem.* **161**, 221–231.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2002). SMART and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2003). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bueno, C. & Churchill, R. M. (1981). *Inorg. Chem.* **20**, 2197–2202.
 Changamu, E. O., Friedrich, H. B., Onani, M. O. & Rademeyer, M. (2006). *J. Organomet. Chem.* **691**, 4615–4625.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Friedrich, H. B., Howie, R. A., Laing, M. & Onani, M. O. (2004). *J. Organomet. Chem.* **689**, 181–193.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, m780 [doi:10.1107/S1600536808012828]

Bromido-1 κ Br-tricarbonyl-2 κ^3 C-(2 η^5 -cyclopentadienyl)molybdenum(I)tungsten(I)(W—Mo)

M. O. Onani, J.-A. Gertenbach and M. D. Bala

Comment

The compound **I** was a by-product of a study on the functionalization of paraffins using transition metals. The functionalized compounds have potential applications in catalysis and organic syntheses (Changamu *et al.*, 2006). The compound **I** is similar to the reported structure of $(\eta^5\text{-C}_5\text{H}_5(\text{CO})_3\text{MoHgCl}$ (Bueno *et al.*, 1981), Albright *et al.* (1978). The bond distances of W—Mo, 2.6872 (7) Å and W—Br, 2.5591 (9) Å are comparable to Hg—Mo, 2.693 (30) Å and Hg—Cl, 2.437 (8) Å respectively. The slight difference between the bond lengths involving the halides could be attributed to the difference in electronegativity and hence basicity between bromine and chlorine. The coordination around Mo is a pseudo-square pyramidal piano stool arrangement. (Fig. 1)

Experimental

The compound **I** was prepared according to a reported procedure (Friedrich *et al.*, 2004) and crystals were grown by slow evaporation of a mixture of dichloromethane and hexane at 263 K.

Refinement

Hydrogen atoms were treated as riding on their parent C atoms with C—H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure of the title complex showing the atom numbering scheme. Ellipsoids are drawn at the 50% probability level.

Bromido-1 κ Br-tricarbonyl-2 κ^3 C-(2 η^5 -cyclopentadienyl)molybdenum(I)tungsten(I)(W—Mo)

Crystal data

[WMoBr(C₅H₅)(CO)₃]

$M_r = 508.82$

Tetragonal, $P\bar{4}2_1c$

Hall symbol: P -4 2n

$a = 11.9375$ (9) Å

$Z = 8$

$F_{000} = 1824$

$D_x = 3.051$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2238 reflections

supplementary materials

$b = 11.9375 (9) \text{ \AA}$	$\theta = 2.2\text{--}25.5^\circ$
$c = 15.546 (2) \text{ \AA}$	$\mu = 15.09 \text{ mm}^{-1}$
$\alpha = 90^\circ$	$T = 100 (2) \text{ K}$
$\beta = 90^\circ$	Block, yellow
$\gamma = 90^\circ$	$0.11 \times 0.10 \times 0.07 \text{ mm}$
$V = 2215.4 (4) \text{ \AA}^3$	

Data collection

Bruker APEX CCD area-detector diffractometer	2673 independent reflections
Radiation source: fine-focus sealed tube	2497 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.048$
$T = 100(2) \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2002)	$h = -15 \rightarrow 14$
$T_{\text{min}} = 0.251, T_{\text{max}} = 0.347$	$k = -8 \rightarrow 15$
13298 measured reflections	$l = -20 \rightarrow 18$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.030$	$w = 1/[\sigma^2(F_o^2) + (0.0238P)^2]$
$wR(F^2) = 0.069$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.02$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2673 reflections	$\Delta\rho_{\text{max}} = 1.31 \text{ e \AA}^{-3}$
127 parameters	$\Delta\rho_{\text{min}} = -0.72 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 1118 Friedel pairs
	Flack parameter: 0.00 (1)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
W1	0.64476 (3)	0.39974 (3)	0.89630 (2)	0.01742 (9)
Mo1	0.73518 (6)	0.20182 (6)	0.85204 (5)	0.01833 (16)
Br1	0.62054 (7)	0.61253 (7)	0.90359 (5)	0.01974 (17)
O1	0.5124 (6)	0.1402 (6)	0.9503 (5)	0.0480 (19)
O2	0.6348 (5)	0.0161 (6)	0.7340 (4)	0.0402 (17)
O3	0.6948 (5)	0.3305 (6)	0.6794 (4)	0.0330 (16)
C1	0.5937 (8)	0.1680 (7)	0.9122 (6)	0.031 (2)
C2	0.6724 (7)	0.0833 (7)	0.7766 (6)	0.026 (2)
C3	0.7057 (7)	0.2868 (8)	0.7450 (6)	0.027 (2)
C4	0.8467 (10)	0.1507 (11)	0.9698 (7)	0.047 (3)
H4	0.8151	0.1245	1.0222	0.057*
C5	0.8729 (8)	0.0831 (8)	0.9024 (7)	0.038 (2)
H5	0.8622	0.0043	0.8996	0.046*
C6	0.9203 (7)	0.1535 (11)	0.8354 (6)	0.044 (3)
H6	0.9468	0.1308	0.7805	0.052*
C7	0.9181 (9)	0.2684 (10)	0.8715 (9)	0.059 (4)
H7	0.9431	0.3359	0.8455	0.071*
C8	0.8694 (10)	0.2533 (11)	0.9549 (7)	0.052 (3)
H8	0.8557	0.3126	0.9943	0.063*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
W1	0.01888 (17)	0.01447 (16)	0.01889 (15)	0.00532 (12)	0.00265 (14)	-0.00152 (14)
Mo1	0.0158 (3)	0.0170 (3)	0.0222 (3)	0.0033 (3)	0.0015 (3)	-0.0020 (3)
Br1	0.0214 (4)	0.0168 (4)	0.0209 (4)	0.0015 (3)	0.0032 (3)	0.0010 (3)
O1	0.036 (4)	0.032 (4)	0.076 (5)	0.006 (3)	0.029 (4)	0.011 (4)
O2	0.034 (4)	0.033 (4)	0.054 (4)	0.006 (3)	-0.008 (3)	-0.016 (3)
O3	0.035 (4)	0.045 (4)	0.019 (3)	0.007 (3)	-0.005 (3)	0.002 (3)
C1	0.032 (5)	0.015 (4)	0.045 (6)	0.009 (4)	0.004 (5)	0.003 (4)
C2	0.020 (5)	0.020 (5)	0.037 (5)	0.003 (4)	0.006 (4)	-0.012 (4)
C3	0.020 (5)	0.033 (5)	0.029 (5)	0.002 (4)	-0.004 (4)	-0.008 (4)
C4	0.042 (6)	0.069 (8)	0.030 (5)	-0.002 (7)	0.000 (5)	0.002 (6)
C5	0.033 (5)	0.030 (5)	0.053 (6)	0.015 (4)	-0.025 (5)	0.004 (5)
C6	0.016 (5)	0.091 (9)	0.024 (5)	0.028 (5)	-0.007 (4)	-0.007 (5)
C7	0.025 (6)	0.043 (7)	0.110 (11)	-0.014 (5)	-0.036 (6)	0.048 (7)
C8	0.037 (7)	0.058 (8)	0.061 (7)	0.015 (6)	-0.016 (5)	-0.035 (7)

Geometric parameters (\AA , $^\circ$)

W1—Br1	2.5591 (9)	O3—C3	1.152 (12)
W1—Mo1	2.6872 (7)	C4—C8	1.276 (17)
Mo1—C1	1.972 (10)	C4—C5	1.359 (15)
Mo1—C3	1.980 (10)	C4—H4	0.9500
Mo1—C2	1.985 (8)	C5—C6	1.453 (15)

supplementary materials

Mo1—C6	2.298 (8)	C5—H5	0.9500
Mo1—C5	2.307 (8)	C6—C7	1.482 (17)
Mo1—C7	2.344 (10)	C6—H6	0.9500
Mo1—C4	2.345 (12)	C7—C8	1.433 (16)
Mo1—C8	2.346 (10)	C7—H7	0.9500
O1—C1	1.184 (11)	C8—H8	0.9500
O2—C2	1.132 (10)		
Br1—W1—Mo1	158.35 (3)	C4—Mo1—W1	104.9 (3)
C1—Mo1—C3	110.5 (4)	C8—Mo1—W1	82.5 (3)
C1—Mo1—C2	79.1 (4)	O1—C1—Mo1	175.0 (8)
C3—Mo1—C2	78.5 (4)	O2—C2—Mo1	178.9 (8)
C1—Mo1—C6	145.6 (4)	O3—C3—Mo1	174.3 (8)
C3—Mo1—C6	101.8 (4)	C8—C4—C5	112.4 (11)
C2—Mo1—C6	96.8 (4)	C8—C4—Mo1	74.3 (7)
C1—Mo1—C5	108.9 (4)	C5—C4—Mo1	71.5 (6)
C3—Mo1—C5	136.6 (4)	C8—C4—H4	123.8
C2—Mo1—C5	91.8 (4)	C5—C4—H4	123.8
C6—Mo1—C5	36.8 (4)	Mo1—C4—H4	121.9
C1—Mo1—C7	143.7 (4)	C4—C5—C6	107.4 (10)
C3—Mo1—C7	95.8 (4)	C4—C5—Mo1	74.5 (6)
C2—Mo1—C7	132.1 (4)	C6—C5—Mo1	71.3 (5)
C6—Mo1—C7	37.2 (4)	C4—C5—H5	126.3
C5—Mo1—C7	60.0 (4)	C6—C5—H5	126.3
C1—Mo1—C4	93.6 (4)	Mo1—C5—H5	119.8
C3—Mo1—C4	152.9 (4)	C5—C6—C7	104.9 (9)
C2—Mo1—C4	119.4 (4)	C5—C6—Mo1	71.9 (5)
C6—Mo1—C4	58.5 (4)	C7—C6—Mo1	73.0 (5)
C5—Mo1—C4	34.0 (4)	C5—C6—H6	127.6
C7—Mo1—C4	57.2 (4)	C7—C6—H6	127.6
C1—Mo1—C8	108.4 (4)	Mo1—C6—H6	119.6
C3—Mo1—C8	124.1 (4)	C8—C7—C6	103.5 (9)
C2—Mo1—C8	148.0 (4)	C8—C7—Mo1	72.3 (6)
C6—Mo1—C8	59.1 (4)	C6—C7—Mo1	69.7 (5)
C5—Mo1—C8	56.1 (4)	C8—C7—H7	128.2
C7—Mo1—C8	35.6 (4)	C6—C7—H7	128.2
C4—Mo1—C8	31.6 (4)	Mo1—C7—H7	121.7
C1—Mo1—W1	73.4 (2)	C4—C8—C7	111.8 (10)
C3—Mo1—W1	72.1 (3)	C4—C8—Mo1	74.2 (7)
C2—Mo1—W1	128.8 (3)	C7—C8—Mo1	72.1 (6)
C6—Mo1—W1	129.5 (3)	C4—C8—H8	124.1
C5—Mo1—W1	137.7 (3)	C7—C8—H8	124.1
C7—Mo1—W1	92.5 (3)	Mo1—C8—H8	121.1
Br1—W1—Mo1—C1	-172.7 (3)	C8—Mo1—C6—C5	73.6 (7)
Br1—W1—Mo1—C3	-54.0 (3)	W1—Mo1—C6—C5	119.9 (6)
Br1—W1—Mo1—C2	-112.2 (3)	C1—Mo1—C6—C7	-115.9 (9)
Br1—W1—Mo1—C6	36.8 (3)	C3—Mo1—C6—C7	83.9 (7)
Br1—W1—Mo1—C5	87.2 (4)	C2—Mo1—C6—C7	163.6 (6)
Br1—W1—Mo1—C7	41.3 (3)	C5—Mo1—C6—C7	-112.4 (8)

Br1—W1—Mo1—C4	97.9 (3)	C4—Mo1—C6—C7	-76.0 (7)
Br1—W1—Mo1—C8	75.5 (3)	C8—Mo1—C6—C7	-38.9 (6)
C1—Mo1—C4—C8	-120.1 (8)	W1—Mo1—C6—C7	7.4 (7)
C3—Mo1—C4—C8	33.5 (13)	C5—C6—C7—C8	-0.4 (9)
C2—Mo1—C4—C8	160.4 (7)	Mo1—C6—C7—C8	65.0 (7)
C6—Mo1—C4—C8	81.2 (8)	C5—C6—C7—Mo1	-65.4 (6)
C5—Mo1—C4—C8	120.8 (11)	C1—Mo1—C7—C8	8.5 (10)
C7—Mo1—C4—C8	36.9 (7)	C3—Mo1—C7—C8	145.7 (7)
W1—Mo1—C4—C8	-46.3 (8)	C2—Mo1—C7—C8	-134.5 (8)
C1—Mo1—C4—C5	119.1 (7)	C6—Mo1—C7—C8	-112.3 (8)
C3—Mo1—C4—C5	-87.3 (11)	C5—Mo1—C7—C8	-72.6 (7)
C2—Mo1—C4—C5	39.6 (8)	C4—Mo1—C7—C8	-32.7 (6)
C6—Mo1—C4—C5	-39.6 (7)	W1—Mo1—C7—C8	73.4 (7)
C7—Mo1—C4—C5	-83.8 (8)	C1—Mo1—C7—C6	120.8 (8)
C8—Mo1—C4—C5	-120.8 (11)	C3—Mo1—C7—C6	-102.0 (6)
W1—Mo1—C4—C5	-167.1 (6)	C2—Mo1—C7—C6	-22.2 (8)
C8—C4—C5—C6	0.7 (13)	C5—Mo1—C7—C6	39.7 (6)
Mo1—C4—C5—C6	64.2 (6)	C4—Mo1—C7—C6	79.6 (7)
C8—C4—C5—Mo1	-63.4 (10)	C8—Mo1—C7—C6	112.3 (8)
C1—Mo1—C5—C4	-67.1 (8)	W1—Mo1—C7—C6	-174.3 (6)
C3—Mo1—C5—C4	138.6 (7)	C5—C4—C8—C7	-1.0 (14)
C2—Mo1—C5—C4	-146.2 (7)	Mo1—C4—C8—C7	-62.8 (8)
C6—Mo1—C5—C4	115.0 (10)	C5—C4—C8—Mo1	61.8 (9)
C7—Mo1—C5—C4	74.7 (8)	C6—C7—C8—C4	0.9 (12)
C8—Mo1—C5—C4	32.8 (7)	Mo1—C7—C8—C4	64.1 (9)
W1—Mo1—C5—C4	18.7 (9)	C6—C7—C8—Mo1	-63.2 (6)
C1—Mo1—C5—C6	177.9 (6)	C1—Mo1—C8—C4	65.5 (8)
C3—Mo1—C5—C6	23.6 (8)	C3—Mo1—C8—C4	-162.4 (7)
C2—Mo1—C5—C6	98.8 (6)	C2—Mo1—C8—C4	-33.5 (12)
C7—Mo1—C5—C6	-40.2 (6)	C6—Mo1—C8—C4	-79.1 (8)
C4—Mo1—C5—C6	-115.0 (10)	C5—Mo1—C8—C4	-35.3 (7)
C8—Mo1—C5—C6	-82.2 (7)	C7—Mo1—C8—C4	-119.8 (10)
W1—Mo1—C5—C6	-96.2 (7)	W1—Mo1—C8—C4	135.2 (7)
C4—C5—C6—C7	-0.1 (10)	C1—Mo1—C8—C7	-174.7 (7)
Mo1—C5—C6—C7	66.2 (6)	C3—Mo1—C8—C7	-42.6 (8)
C4—C5—C6—Mo1	-66.3 (7)	C2—Mo1—C8—C7	86.3 (11)
C1—Mo1—C6—C5	-3.5 (10)	C6—Mo1—C8—C7	40.7 (6)
C3—Mo1—C6—C5	-163.7 (6)	C5—Mo1—C8—C7	84.5 (7)
C2—Mo1—C6—C5	-84.0 (6)	C4—Mo1—C8—C7	119.8 (10)
C7—Mo1—C6—C5	112.4 (8)	W1—Mo1—C8—C7	-105.0 (7)
C4—Mo1—C6—C5	36.5 (6)		

Fig. 1

