

Retraction of articles

This article reports the retraction of 39 articles published in *Acta Crystallographica Section E* between 2004 and 2009.

After thorough investigation (see Harrison *et al.*, 2010), 39 additional articles are retracted as a result of problems with the data sets or incorrect atom assignments. Full details of all the articles are given in Table 1.

Table 1

Details of articles to be retracted, in order of publication.

Title	Reference	Retracted by	DOI	Refcode
<i>trans</i> -Bis[1-[3-(cyclohexylamino)propyliminomethyl]-2-naphtholato]copper(II) dichloride dihydrate	Zhang (2004)	Journal	10.1107/S1600536804028296	BIPDUA
Bis(4-bromo-2-formylphenolato- $\kappa^2 O, O'$)copper(II)	Sun & Gao (2005)	Author	10.1107/S160053680500187X	FEYSUY
Bis(salicylaldehyde)zinc(II)	Xiong & Liu (2005)	Journal	10.1107/S1600536805010913	GAMDUU
Bis(4-bromo-2-formylphenolato- $\kappa^2 O, O'$)zinc(II)	Chen (2006)	Journal	10.1107/S1600536805040432	SAZCUS
Bis(2-formylphenolato- $\kappa^2 O, O'$)nickel(II)	Li & Chen (2006)	Journal	10.1107/S1600536806012931	IDAZAP
Bis(2-formylphenolato)cobalt(II)	Qiu (2006)	Journal	10.1107/S1600536806015704	GEJDUV
Bis(2-formylphenolato- $\kappa^2 O, O'$)manganese(II)	Wang & Fang (2006)	Journal	10.1107/S1600536806021039	IDOVED
Tetraaqua(1,10-phenanthroline- $\kappa^2 N, N'$)copper(II) naphthalene-1,5-disulfonate dihydrate	Liu <i>et al.</i> (2006)	Author	10.1107/S1600536806030637	GENYOO
Tetraaqua(1,10-phenanthroline- $\kappa^2 N, N'$)nickel(II) naphthalene-1,5-disulfonate dihydrate	Liu & Fan (2006)	Author	10.1107/S1600536806035410	KERBEP
{6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-trinitratolutetium(III)copper(II)	Sui <i>et al.</i> (2006)	Journal	10.1107/S160053680604565X	HESPEB
Bis(2-formylphenolato- $\kappa^2 O, O'$)iron(II)	Yang <i>et al.</i> (2007)	Author	10.1107/S1600536807021721	PIFCAJ
2,6-Dimethoxybenzohydrazide	Qadeer <i>et al.</i> (2007a)	Journal	10.1107/S1600536807022593	PIFHES
2-(2,4-Dichlorophenylsulfanyl)acetohydrazide	Qadeer <i>et al.</i> (2007b)	Journal	10.1107/S1600536807022891	YIFSOW
{6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-trinitratoeuropium(III)zinc(II)	Hu <i>et al.</i> (2007)	Author	10.1107/S1600536807031121	WIHKEE
{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-trinitratocerium(III)zinc(II)	Sui, Zhang, Hu & Yin (2007)	Author	10.1107/S1600536807032564	WIHREL
{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-trinitratopraseodymium(III)zinc(II)	Chen <i>et al.</i> (2007)	Author	10.1107/S1600536807032540	WIHRIP
{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-trinitratopraseodymium(III)nickel(II)	Sui, Li <i>et al.</i> (2007)	Author	10.1107/S1600536807032618	UFACUA
{6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato-1 $\kappa^4 O^1, O^2, O^3, O^4, O^5, O^6, O^7, O^8, O^9, O^{10}$:2 $\kappa^2 O^1, N, N', N'', N'''$ }(methanol-1 κO)- μ -nitrate-1:2 $\kappa^2 O:O'$ -dinitrate-1 $\kappa^4 O, O'$ -cerium(III)zinc(II)	Sui, Fang, Hu & Lin (2007)	Author	10.1107/S1600536807033314	UDUYIC
{6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-methanol- μ -nitrate-dinitratosamarium(III)nickel(II)	Sui, Zhang, Hu & Jiang (2007)	Author	10.1107/S1600536807037130	AFECEU
{6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-methanol- μ -nitrate-dinitratopraseodymium(III)zinc(II)	Sui, Fang & Yuan (2007)	Author	10.1107/S1600536807037488	AFICEY
{6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-methanol- μ -nitrate-dinitratolutetium(III)zinc(II)	Sui, Sui <i>et al.</i> (2007)	Author	10.1107/S1600536807037737	AFEF0H
catena-Poly[[chloridonickel(II)]-di- μ -chlorido-[chloridonickel(II)]- μ -4,4'-methylenebis(3,5-dimethylpyrazole)- $\kappa^2 N^2, N^2$]	Huang & Chen (2007)	Author	10.1107/S1600536807039384	VIJYOD
[2,2'-[<i>o</i> -Phenylenebis(nitrilomethylidyne)]diphenolato]zinc(II)	Liu <i>et al.</i> (2007a)	Author	10.1107/S1600536807040640	DIKYUS
<i>trans</i> -Bis(ethylenediamine- $\kappa^2 N, N'$)bis(nitrate- κO)zinc(II)	Liu, Zeng & Chen (2007)	Author	10.1107/S1600536807042390	XIKYEW
[<i>N, N'</i> -(<i>o</i> -Phenylene)bis(picolinamido)- $\kappa^2 N, N', N'', N'''$]cobalt(II)	Liu & Zeng (2007a)	Author	10.1107/S1600536807044571	XILFII
[<i>N, N'</i> -(<i>o</i> -Phenylene)dipicolinamide- $\kappa^2 N$]nickel(II)	Liu & Zeng (2007b)	Author	10.1107/S1600536807048386	WINWEW
[2,2'-[<i>o</i> -Phenylenebis(nitrilomethylidyne)]diphenolato]manganese(II)	Liu <i>et al.</i> (2007b)	Author	10.1107/S1600536807052993	VIQPIV
<i>N</i> -(2-Amino-3-pyridyl)urea monohydrate	Li <i>et al.</i> (2007)	Author	10.1107/S1600536807047526	SIMFEA
<i>N</i> -(2-Fluorophenyl)carbamic acid monohydrate	Yang (2007)	Author	10.1107/S1600536807052464	WINMOW
Aqua(dimethylglyoxime- $\kappa^2 N, N'$)(3,5-dinitro-2-oxidobenzooato- $\kappa^2 O^1, O^2$)-copper(II)	Liu & Wen (2007)	Author	10.1107/S1600536807054244	HIQCAM
μ -Acetato-tri- μ -ferrocenecarboxylatobis[(<i>N, N</i> -dimethylformamide)-copper(II)]	Liu, Lin <i>et al.</i> (2007)	Journal	10.1107/S1600536807059041	HIQQEE

Table 1 (continued)

Title	Reference	Retracted by	DOI	Refcode
{ μ -6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}- μ -nitrate-dinitratoeuropium(III)zinc(II)	Hu <i>et al.</i> (2008)	Author	10.1107/S160053680706151X	MIRPAF
Bis(4-chloro-2-formylphenolato)nickel(II)	Li <i>et al.</i> (2008)	Author	10.1107/S1600536807056309	RISTET
{ μ -6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}- μ -nitrate-dinitratoerbium(III)zinc(II)	Chen <i>et al.</i> (2008)	Author	10.1107/S1600536808006958	QIXHIP
Bis(2-ethoxy-6-formylphenolato- $\kappa^2 O^1, O^6$)nickel(II)	Han (2008)	Journal	10.1107/S160053680800809X	QIXLIT
{ μ -6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}- μ -nitrate-dinitratoholmium(III)zinc(II)	Xiao, Sui <i>et al.</i> (2008)	Author	10.1107/S1600536808013743	BIZTUA
{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato}-trinitratoholmium(III)nickel(II)	Xiao, Fu <i>et al.</i> (2008)	Author	10.1107/S1600536808013755	BIZVAI
Hydrogen-bonding patterns in the cocrystal terephthalic acid-4,4'-bipyridine (2I)	Wang <i>et al.</i> (2009)	Journal	10.1107/S160053680903236X	DUCZEH
{6,6'-Dimethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidyne)]diphenolato-1 $\kappa^4 O^1, O^1, O^6, O^6$:2 $\kappa^4 O^1, N, N, O^1$ } (ethanol-1 κO)- μ -nitrate-1:2 $\kappa^2 O$:O'-dinitrato-1 $\kappa^2 O, O'$ -samarium(III)zinc(II)	Huang <i>et al.</i> (2009)	Journal	10.1107/S1600536809033558	YUCWAV

References

- Chen, Q. (2006). *Acta Cryst.* **E62**, m56–m57.
- Chen, J.-R., Sui, Y., Luo, Q.-Y. & Jiang, R.-Q. (2007). *Acta Cryst.* **E63**, m2091–m2092.
- Chen, J.-R., Sui, Y., Chen, L., Wen, J.-W. & Yin, L.-Y. (2008). *Acta Cryst.* **E64**, m562–m563.
- Han, Z.-Q. (2008). *Acta Cryst.* **E64**, m592.
- Harrison, W. T. A., Simpson, J. & Weil, M. (2010). *Acta Cryst.* **E66**, e1–e2.
- Hu, R.-H., Sui, Y., Chen, L. & He, C.-M. (2008). *Acta Cryst.* **E64**, m8–m9.
- Hu, R.-H., Sui, Y., Fang, X.-N. & Chen, H.-M. (2007). *Acta Cryst.* **E63**, m2039–m2040.
- Huang, C.-F. & Chen, H.-L. (2007). *Acta Cryst.* **E63**, m2356–m2357.
- Huang, Q., Sui, Y.-H. & Zhang, G.-X. (2009). *Acta Cryst.* **E65**, m1161–m1162.
- Li, Y.-G. & Chen, H.-J. (2006). *Acta Cryst.* **E62**, m1038–m1039.
- Li, N.-G., Tao, R.-M. & Fu, B.-F. (2007). *Acta Cryst.* **E63**, o4228.
- Li, Z., Zhang, X. & Pu, X. (2008). *Acta Cryst.* **E64**, m215.
- Liu, J.-T. & Fan, S.-D. (2006). *Acta Cryst.* **E62**, m2507–m2508.
- Liu, J.-T., Fan, S.-D. & Li, D.-Q. (2006). *Acta Cryst.* **E62**, m2165–m2166.
- Liu, D., Lin, J., Xu, Y., Huang, C. & Li, X. (2007). *Acta Cryst.* **E63**, m3094.
- Liu, Y.-Q. & Wen, H.-R. (2007). *Acta Cryst.* **E63**, m2928.
- Liu, Y.-Q. & Zeng, X.-R. (2007a). *Acta Cryst.* **E63**, m2547.
- Liu, Y.-Q. & Zeng, X.-R. (2007b). *Acta Cryst.* **E63**, m2684.
- Liu, Y.-Q., Zeng, X.-R. & Chen, W.-T. (2007). *Acta Cryst.* **E63**, m2462.
- Liu, Y.-Q., Zeng, X.-R., Luo, Q.-Y. & Xu, Y.-P. (2007a). *Acta Cryst.* **E63**, m2396.
- Liu, Y.-Q., Zeng, X.-R., Luo, Q.-Y. & Xu, Y.-P. (2007b). *Acta Cryst.* **E63**, m2854.
- Qadeer, G., Rama, N. H. & Chen, W.-T. (2007a). *Acta Cryst.* **E63**, o2892.
- Qadeer, G., Rama, N. H. & Chen, W.-T. (2007b). *Acta Cryst.* **E63**, o2932.
- Qiu, X.-Y. (2006). *Acta Cryst.* **E62**, m1190–m1191.
- Sui, Y., Fang, X.-N., Hu, P. & Lin, J. (2007). *Acta Cryst.* **E63**, m2135–m2136.
- Sui, Y., Fang, X.-N. & Yuan, M.-W. (2007). *Acta Cryst.* **E63**, m2275–m2276.
- Sui, Y., Li, X.-F., Huang, G.-S. & Wang, G.-J. (2007). *Acta Cryst.* **E63**, m2093–m2094.
- Sui, Y., Sui, Y.-H., Luo, Q.-Y. & Wang, Y.-D. (2007). *Acta Cryst.* **E63**, m2277–m2278.
- Sui, Y., Xiao, Y.-A., Fang, X.-N., Zeng, X.-R. & Li, M.-H. (2006). *Acta Cryst.* **E62**, m3205–m3207.
- Sui, Y., Zhang, J.-H., Hu, R.-H. & Jiang, R.-Q. (2007). *Acta Cryst.* **E63**, m2256–m2257.
- Sui, Y., Zhang, J.-H., Hu, R.-H. & Yin, L.-Y. (2007). *Acta Cryst.* **E63**, m2089–m2090.
- Sun, Y.-X. & Gao, G.-Z. (2005). *Acta Cryst.* **E61**, m354–m355.
- Wang, Q. & Fang, X.-N. (2006). *Acta Cryst.* **E62**, m1492–m1493.
- Wang, S., Yang, T., Li, Z. & Yu, X. (2009). *Acta Cryst.* **E65**, o2198.
- Xiao, Y.-A., Fu, X.-K., Sui, Y., Wu, Q. & Xiong, S.-H. (2008). *Acta Cryst.* **E64**, m806–m807.
- Xiao, Y.-A., Sui, Y., Yi, X.-G., Wu, J.-H. & Zhang, L.-P. (2008). *Acta Cryst.* **E64**, m804–m805.
- Xiong, Z.-Y. & Liu, L.-J. (2005). *Acta Cryst.* **E61**, m863–m864.
- Yang, X.-M. (2007). *Acta Cryst.* **E63**, o4453.
- Yang, Y.-M., Lu, P.-C., Zhu, T.-T. & Liu, C.-H. (2007). *Acta Cryst.* **E63**, m1613.
- Zhang, P. (2004). *Acta Cryst.* **E60**, m1808–m1810.

{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidene)]diphenolato}-trinitratoholmium(III)nickel(II)

Yi-An Xiao,^a Xiang-Kai Fu,^b Yan Sui,^{c*} Qing Wu^c and Shao-Hui Xiong^c

^aCollege of Life Sciences, JingGangShan University, 343009 Ji'an, JiangXi, People's Republic of China, ^bCollege of Chemistry & Chemical Engineering, Southwest University, 400715 Beibei, Chongqing, People's Republic of China, and ^cJiangXi Province Key Laboratory of Coordination Chemistry, College of Chemistry & Chemical Engineering, JingGangShan University, 343009 Ji'an, JiangXi, People's Republic of China

Correspondence e-mail: ysui@163.com


Received 6 May 2008; accepted 8 May 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.043; wR factor = 0.114; data-to-parameter ratio = 16.4.

In the title heteronuclear Ni^{II}-Ho^{III} complex (systematic name: { μ -6,6'-diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidene)]diphenolato-1 κ^4 O¹,O^{1'},O⁶,O^{6'}:2 κ^4 O¹,N,N',O^{1'}}trinitrato-1 κ^6 O,O'-holmium(III)nickel(II)), [HoNi(C₂₀H₂₂N₂O₄)(NO₃)₃], with the hexadentate Schiff base compartmental ligand *N,N'*-bis(3-ethoxysalicylidene)ethylenediamine (H₂L), the Ho and Ni atoms are doubly bridged by two phenolate O atoms of the Schiff base ligand. The coordination of Ni is square-planar with the donor centers of two imine N atoms and two phenolate O atoms. The holmium(III) center has a tenfold coordination environment of O atoms, involving the phenolate O atoms, two ethoxy O atoms and two O atoms each from the three nitrates. Weak C—H...O and O...Ni [3.383 (4) Å] interactions generate a two-dimensional zigzag sheet.

Related literature

For related literature, see: Baggio *et al.* (2000); Caravan *et al.* (1999); Edder *et al.* (2000); Knoer *et al.* (2005); Sui *et al.* (2006).



Experimental

Crystal data

[HoNi(C₂₀H₂₂N₂O₄)(NO₃)₃]
 $M_r = 764.07$
 Orthorhombic, $P2_12_12_1$
 $a = 8.5825$ (8) Å
 $b = 13.7028$ (14) Å
 $c = 21.203$ (2) Å
 $V = 2493.6$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 3.98$ mm⁻¹
 $T = 293$ (2) K
 $0.17 \times 0.16 \times 0.13$ mm

Data collection

Bruker APEXII area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2004)
 $T_{\min} = 0.559$, $T_{\max} = 0.625$
 18705 measured reflections
 5970 independent reflections
 4299 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.055$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.113$
 $S = 1.04$
 5970 reflections
 364 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 2.00$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.61$ e Å⁻³
 Absolute structure: Flack (1983), 2455 Friedel pairs
 Flack parameter: -0.003 (18)

Table 1

Selected bond lengths (Å).

Ho1—O1	2.390 (4)	Ho1—O9	2.492 (5)
Ho1—O2	2.343 (4)	Ho1—O11	2.531 (5)
Ho1—O3	2.540 (4)	Ho1—O12	2.425 (5)
Ho1—O4	2.594 (4)	Ni1—O1	1.846 (4)
Ho1—O5	2.430 (5)	Ni1—O2	1.847 (4)
Ho1—O6	2.468 (5)	Ni1—N1	1.837 (5)
Ho1—O8	2.460 (5)	Ni1—N2	1.837 (5)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C12—H12...O10 ⁱ	0.93	2.37	3.286 (8)	169
C10—H10A...O10 ⁱⁱ	0.97	2.42	3.301 (9)	150

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2*; data reduction: *APEX2*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *APEX2*; software used to prepare material for publication: *APEX2* and *publCIF* (Westrip, 2008).

We gratefully acknowledge financial support from the Department of Education, JiangXi Province (No. 2007317) and the Natural Science Foundation of JiangXi Province (No. 2007GZH1667).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2401).

References

- Baggio, R., Garland, M. T., Moreno, Y., Pena, O., Pereg, M. & Spodine, E. (2000). *J. Chem. Soc. Dalton Trans.* pp. 2061–2066.
- Bruker (2004). *APEX2* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Caravan, P., Ellison, J. J., McMurry, T. J. & Lauffer, R. B. (1999). *Chem. Rev.* **99**, 2293–2352.
- Edder, C., Piguet, C., Bernardinelli, G., Mareda, J., Bochet, C. G., Bunzli, J.-C. G. & Hopfgartner, G. (2000). *Inorg. Chem.* **39**, 5059–5073.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Knoer, R., Lin, H.-H., Wei, H.-H. & Mohanta, S. (2005). *Inorg. Chem.* **44**, 3524–3536.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sui, Y., Fang, X.-N., Xiao, Y.-A., Luo, Q.-Y. & Li, M.-H. (2006). *Acta Cryst.* **E62**, m2230–m2232.
- Westrip, S. P. (2008). *publCIF*. In preparation.

Article retracted

supplementary materials

Article retracted

Acta Cryst. (2008). E64, m806-m807 [doi:10.1107/S1600536808013755]

{ μ -6,6'-Diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidene)]diphenolato}trinitratoholmium(III)nickel(II)

Y.-A. Xiao, X.-K. Fu, Y. Sui, Q. Wu and S.-H. Xiong

Comment

The potential applications of trivalent lanthanide complexes as contrast agent for magnetic resonance imaging and stains for fluorescence imaging have prompted considerable interest in the preparation, magnetic and optical properties of 3 d-4f heterometallic dinuclear complexes (Baggio *et al.*, 2000; Caravan *et al.*, 1999; Edder *et al.*, 2000; Knoer *et al.*, 2005). As part of our investigations into the structure and applications of 3 d-4f heterometallic Schiff base complexes (Sui *et al.* 2006), we report here the synthesis and X-ray crystal structure analysis of the title complex, (I), a new Ni^{II}—Ho^{III} complex with salen-type Schiff base *N,N*-bis(3-ethoxysalicylidene) ethylenediamine(H₂L).

Complex (I) crystallizes in the space group *P*2₁2₁2₁, with nickel and holmium doubly bridged by two phenolate O atoms provided by a salen-type Schiff base ligand. The inner salen-type cavity is occupied by nickel(II), while holmium(III) is present in the open and larger portion of the dinucleating compartmental Schiff base ligand. The dihedral angles between the mean planes of Ni1/O1/O2 and Ho1/O1/O2 is 6.97 (26)° suggesting that the bridging moiety is almost planar; the deviation of atoms from the least squares Ni1/O1/O2/Ho1 plane being -0.0583 (2) Å for Ni, -0.0397 (3) Å for Ho, 0.0483 (2) Å for O1 and 0.0497 (2) Å for O2.

The holmium(III) center in (I) has a decacoordination environment of O atoms. In addition to the phenolate ligands, two ethoxy O atoms coordinate to this metal center, two O atoms from each of the three nitrates chelate to holmium to complete the decacoordination. The three kinds of Ho—O bond distances are significantly different, the shortest being the Ho—O(phenolate) and longest being the Ho—O(ethoxy) separations.

The coordination of nickel(II) is approximately square planar. The donor centers are alternatively above and below the mean N₂O₂ plane with an average deviation from the plane of 0.0698 (2) Å, while Ni1 is 0.0022 (2) Å above this square plane.

Adjacent molecules are held together by weak interactions (O13ⁱⁱⁱ⋯Ni1=3.383 (4) Å, C10—H10Aⁱ⋯O10ⁱ=3.301 (9), and C12—H12ⁱⁱ⋯O10ⁱⁱ=3.286 (8); symmetry codes:(i)x - 1, y, z; (ii)-x, 1/2 + y, 1/2 - z.) these link the molecules into a two-dimensional zigzag sheet (Fig 2).

Experimental

H₂L was prepared by the 2:1 condensation of 3-ethoxysalicylaldehyde and ethylenediamine in methanol. Complex (I) was obtained by the treatment of nickel(II) acetate tetrahydrate (0.217 g, 1 mmol) with H₂L(0.356 g, 1 mmol) in methanol solution (80 ml) under reflux for 3 h and then for another 3 h after the addition of holmium(III) nitrate hexahydrate (0.459 g, 1 mmol). The reaction mixture was cooled and the resulting precipitate was filtered off, washed with diethyl ether and dried *in vacuo*. Single crystals of (I) suitable for X-ray analysis were obtained by slow evaporation at room temperature of

supplementary materials

a methanol solution. Analysis calculated for $C_{20}H_{22}HoN_5NiO_{13}$: C 31.44, H 2.90, Ho 21.59, N 9.17, Ni 7.68%; found: C 31.85, H 2.95, Ho 21.55, N 9.24, Ni 7.78. IR(KBr, cm^{-1}): 1645(C=N), 1385,1491(nitrate).

Refinement

The H atoms were positioned geometrically and treated as riding on their parent atoms, with C—H distances of 0.97 (methylene) and 0.96 Å (methyl), and with $U_{iso}(H) = 1.5U_{eq}(C)$ for methyl H atoms and $1.2U_{eq}(C)$ for other H atoms.

Figures




Fig. 1. The molecular structure of (I), showing 30% probability displacement ellipsoids.




Fig. 2. The packing diagram of (I), viewed along the *b* axis; hydrogen bonds are shown as dashed lines.

{ μ -6,6'-diethoxy-2,2'-[ethane-1,2-diylbis(nitrilomethylidene)]diphenolato- $1\kappa^4O^1,O^1',O^6,O^6'$:
 $2\kappa^4O^1,N,N',O^1',O^1'$ }trinitrato- $1\kappa^6O,O'$ -holmium(III) nickel(II)

Crystal data

[HoNi(C₂₀H₂₂N₂O₄)(NO₃)₃]

$M_r = 764.07$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 8.5825$ (8) Å

$b = 13.7028$ (14) Å

$c = 21.203$ (2) Å

$V = 2493.6$ (4) Å³

$Z = 4$

$F_{000} = 1504$

$D_x = 2.035$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 6504 reflections

$\theta = 1.9$ – 28.3°

$\mu = 3.98$ mm⁻¹

$T = 293$ (2) K

Block, red

$0.17 \times 0.16 \times 0.13$ mm

Data collection

Bruker APEXII area-detector diffractometer	5970 independent reflections
Radiation source: fine-focus sealed tube	4299 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.055$
$T = 293(2)$ K	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2004)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.559$, $T_{\text{max}} = 0.626$	$k = -17 \rightarrow 18$
18705 measured reflections	$l = -27 \rightarrow 27$

Refinement

Refinement on F^2	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0473P)^2]$
$R[F^2 > 2\sigma(F^2)] = 0.043$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.113$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.04$	$\Delta\rho_{\text{max}} = 2.00 \text{ e } \text{\AA}^{-3}$
5970 reflections	$\Delta\rho_{\text{min}} = -0.61 \text{ e } \text{\AA}^{-3}$
364 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001x Fc^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0058 (4)
Secondary atom site location: difference Fourier map	Absolute structure: Flack (1983), 2455 Friedel pairs
Hydrogen site location: inferred from neighbouring sites	Flack parameter: -0.003 (18)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C5	-0.1509 (7)	0.3317 (5)	0.3044 (3)	0.0427 (15)
Ho1	0.25263 (3)	0.49982 (2)	0.400886 (12)	0.04143 (11)

supplementary materials

Ni1	-0.06665 (9)	0.56000 (6)	0.31450 (4)	0.03838 (19)
O2	0.0323 (4)	0.4496 (3)	0.34434 (19)	0.0399 (10)
O1	0.0808 (5)	0.6237 (3)	0.36270 (19)	0.0411 (10)
N1	-0.1998 (6)	0.4934 (4)	0.2621 (2)	0.0434 (12)
N2	-0.1729 (6)	0.6703 (4)	0.2905 (3)	0.0423 (13)
N3	0.1352 (8)	0.5100 (5)	0.5279 (3)	0.0588 (16)
O6	0.0496 (6)	0.5277 (4)	0.4813 (2)	0.0647 (14)
C12	-0.1460 (7)	0.7577 (5)	0.3087 (3)	0.0470 (16)
H12	-0.2126	0.8065	0.2945	0.056*
O3	0.1551 (5)	0.3283 (3)	0.4234 (2)	0.0422 (10)
C18	0.0875 (7)	0.7189 (4)	0.3736 (3)	0.0386 (13)
O4	0.3135 (5)	0.6799 (3)	0.4295 (2)	0.0473 (11)
O5	0.2787 (6)	0.4948 (4)	0.5150 (2)	0.0609 (13)
C15	0.1149 (8)	0.9161 (5)	0.4034 (4)	0.0556 (18)
H15	0.1222	0.9816	0.4147	0.067*
C4	-0.0255 (7)	0.3594 (4)	0.3442 (3)	0.0387 (14)
O7	0.0834 (8)	0.5055 (4)	0.5815 (3)	0.0541 (17)
C19	0.4500 (8)	0.7097 (5)	0.4648 (3)	0.0533 (17)
H19A	0.4917	0.6538	0.4872	0.064*
H19B	0.4203	0.7583	0.4957	0.064*
C1	0.3381 (9)	0.1918 (6)	0.4400 (4)	0.059 (2)
H1A	0.2848	0.1544	0.4084	0.089*
H1B	0.3793	0.1487	0.4716	0.089*
H1C	0.4219	0.2275	0.4208	0.089*
C17	0.2135 (7)	0.7529 (4)	0.4098 (3)	0.0422 (15)
C13	-0.0195 (8)	0.7860 (4)	0.3498 (3)	0.0434 (15)
C14	-0.0046 (9)	0.8868 (5)	0.3667 (3)	0.0515 (17)
H14	-0.0773	0.9320	0.3524	0.062*
C2	0.2241 (8)	0.2632 (5)	0.4706 (3)	0.0487 (16)
H2A	0.1421	0.2270	0.4917	0.058*
H2B	0.2784	0.3017	0.5021	0.058*
O9	0.3184 (6)	0.5422 (4)	0.2897 (2)	0.0581 (13)
N4	0.3530 (6)	0.4577 (5)	0.2744 (3)	0.0517 (16)
O8	0.3482 (6)	0.3918 (3)	0.3167 (3)	0.0573 (12)
C9	-0.2254 (8)	0.3989 (5)	0.2623 (3)	0.0447 (15)
H9	-0.2964	0.3741	0.2333	0.054*
O12	0.5289 (6)	0.5328 (4)	0.3902 (3)	0.0593 (13)
N5	0.5831 (8)	0.4621 (5)	0.4212 (3)	0.0605 (16)
C16	0.2267 (9)	0.8516 (5)	0.4248 (3)	0.0510 (17)
H16	0.3099	0.8738	0.4489	0.061*
C6	-0.2073 (8)	0.2359 (5)	0.3070 (3)	0.0506 (17)
H6	-0.2863	0.2162	0.2798	0.061*
C3	0.0358 (7)	0.2907 (5)	0.3862 (3)	0.0417 (14)
C11	-0.3100 (8)	0.6507 (5)	0.2514 (3)	0.0516 (17)
H11A	-0.3272	0.7036	0.2218	0.062*
H11B	-0.4022	0.6434	0.2774	0.062*
C10	-0.2754 (8)	0.5578 (5)	0.2171 (3)	0.0513 (16)
H10A	-0.3710	0.5285	0.2016	0.062*
H10B	-0.2073	0.5702	0.1815	0.062*

O11	0.4839 (6)	0.4045 (4)	0.4421 (3)	0.0728 (17)
C20	0.5726 (10)	0.7506 (6)	0.4231 (4)	0.071 (2)
H20A	0.6009	0.7031	0.3919	0.106*
H20B	0.6624	0.7672	0.4478	0.106*
H20C	0.5336	0.8081	0.4026	0.106*
C8	-0.0229 (7)	0.1970 (5)	0.3887 (4)	0.0484 (16)
H8	0.0192	0.1514	0.4164	0.058*
O10	0.3939 (7)	0.4348 (5)	0.2204 (2)	0.0808 (18)
O13	0.7234 (6)	0.4536 (7)	0.4307 (3)	0.099 (2)
C7	-0.1465 (9)	0.1717 (5)	0.3492 (3)	0.0563 (19)
H7	-0.1884	0.1092	0.3518	0.068*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C5	0.040 (3)	0.041 (3)	0.048 (4)	-0.006 (3)	-0.003 (3)	-0.002 (3)
Ho1	0.04247 (16)	0.03543 (16)	0.04639 (17)	-0.00007 (18)	-0.00389 (12)	0.00217 (13)
Ni1	0.0375 (4)	0.0340 (4)	0.0436 (4)	-0.0005 (3)	-0.0050 (4)	0.0047 (4)
O2	0.038 (2)	0.033 (2)	0.049 (2)	-0.0065 (18)	-0.0099 (18)	0.004 (2)
O1	0.045 (2)	0.026 (2)	0.052 (2)	0.0024 (19)	-0.014 (2)	0.004 (2)
N1	0.037 (2)	0.051 (3)	0.043 (3)	-0.005 (3)	-0.002 (2)	0.002 (3)
N2	0.040 (3)	0.038 (3)	0.049 (3)	-0.001 (2)	0.001 (2)	0.010 (3)
N3	0.085 (5)	0.037 (3)	0.054 (3)	0.003 (3)	0.007 (3)	-0.005 (3)
O6	0.066 (3)	0.076 (4)	0.052 (3)	0.003 (3)	-0.001 (3)	-0.001 (3)
C12	0.041 (3)	0.046 (4)	0.054 (4)	0.011 (3)	0.009 (3)	0.018 (4)
O3	0.045 (2)	0.027 (2)	0.054 (3)	-0.0026 (19)	-0.004 (2)	0.008 (2)
C18	0.039 (3)	0.034 (3)	0.043 (3)	-0.002 (3)	0.000 (3)	0.006 (3)
O4	0.049 (2)	0.037 (2)	0.056 (3)	-0.004 (2)	-0.016 (2)	0.000 (2)
O5	0.070 (3)	0.061 (3)	0.051 (3)	0.005 (3)	-0.008 (2)	0.002 (3)
C15	0.063 (4)	0.034 (4)	0.069 (5)	-0.007 (3)	0.008 (4)	0.003 (4)
C4	0.041 (3)	0.036 (3)	0.039 (3)	-0.004 (3)	0.001 (3)	-0.001 (3)
O7	0.120 (5)	0.077 (4)	0.055 (3)	0.004 (4)	0.013 (4)	0.003 (3)
C19	0.048 (4)	0.052 (4)	0.060 (4)	-0.002 (3)	-0.010 (4)	-0.006 (3)
C1	0.057 (4)	0.044 (4)	0.077 (5)	0.010 (3)	-0.016 (4)	-0.002 (4)
C17	0.053 (4)	0.031 (3)	0.043 (3)	0.004 (3)	0.010 (3)	0.003 (3)
C13	0.052 (4)	0.032 (3)	0.046 (3)	0.001 (3)	0.005 (3)	0.004 (3)
C14	0.061 (4)	0.032 (3)	0.062 (4)	0.011 (3)	0.001 (3)	0.008 (3)
C2	0.061 (4)	0.042 (4)	0.043 (3)	0.002 (3)	-0.010 (3)	0.008 (3)
O9	0.069 (3)	0.051 (3)	0.055 (3)	0.002 (3)	-0.004 (3)	0.012 (3)
N4	0.043 (3)	0.063 (4)	0.049 (3)	-0.018 (3)	-0.006 (3)	-0.003 (3)
O8	0.064 (3)	0.051 (3)	0.056 (3)	-0.007 (2)	0.001 (3)	-0.009 (3)
C9	0.047 (4)	0.041 (3)	0.046 (3)	-0.006 (3)	-0.003 (3)	0.000 (3)
O12	0.053 (3)	0.052 (3)	0.073 (3)	0.000 (2)	0.000 (3)	0.005 (3)
N5	0.055 (4)	0.068 (4)	0.059 (3)	-0.002 (3)	0.002 (3)	0.000 (3)
C16	0.054 (4)	0.047 (4)	0.053 (3)	-0.011 (3)	0.005 (3)	-0.001 (3)
C6	0.047 (4)	0.041 (4)	0.064 (4)	-0.006 (3)	0.002 (4)	-0.007 (4)
C3	0.042 (3)	0.034 (3)	0.049 (3)	-0.001 (3)	0.001 (3)	0.001 (3)
C11	0.046 (4)	0.056 (5)	0.053 (4)	0.005 (3)	-0.008 (3)	0.011 (4)

supplementary materials

C10	0.049 (4)	0.051 (4)	0.054 (3)	-0.007 (3)	-0.019 (3)	0.004 (3)
O11	0.054 (3)	0.081 (4)	0.083 (4)	0.005 (3)	-0.003 (3)	0.029 (3)
C20	0.066 (5)	0.058 (5)	0.087 (5)	-0.016 (5)	0.006 (5)	-0.012 (5)
C8	0.047 (4)	0.034 (4)	0.064 (4)	0.001 (3)	0.005 (3)	0.008 (3)
O10	0.080 (4)	0.115 (5)	0.048 (3)	-0.034 (4)	0.013 (3)	-0.018 (4)
O13	0.044 (3)	0.138 (6)	0.114 (5)	0.010 (4)	-0.012 (3)	0.022 (5)
C7	0.063 (4)	0.034 (4)	0.072 (5)	-0.013 (3)	0.000 (4)	-0.002 (4)

Geometric parameters (Å, °)

Ho1—O1	2.390 (4)	C15—H15	0.9300
Ho1—O2	2.343 (4)	C4—C3	1.398 (9)
Ho1—O3	2.540 (4)	C19—C20	1.484 (10)
Ho1—O4	2.594 (4)	C19—H19A	0.9700
Ho1—O5	2.430 (5)	C19—H19B	0.9700
Ho1—O6	2.468 (5)	C1—C2	1.529 (10)
Ho1—O8	2.460 (5)	C1—H1A	0.9600
Ho1—O9	2.492 (5)	C1—H1B	0.9600
Ho1—O11	2.531 (5)	C1—H1C	0.9600
Ho1—O12	2.425 (5)	C17—C16	1.394 (9)
Ni1—O1	1.846 (4)	C13—C14	1.433 (9)
Ni1—O2	1.847 (4)	C14—H14	0.9300
Ni1—N1	1.837 (5)	C2—H2A	0.9700
Ni1—N2	1.837 (5)	C2—H2B	0.9700
C5—C6	1.400 (9)	O9—N4	1.239 (8)
C5—C4	1.419 (8)	N4—O10	1.238 (7)
C5—C9	1.433 (9)	N4—O8	1.272 (8)
O2—C4	1.332 (7)	C9—H9	0.9300
O1—C18	1.327 (7)	O12—N5	1.259 (8)
N1—C9	1.313 (8)	N5—O13	1.227 (8)
N1—C10	1.453 (8)	N5—O11	1.242 (8)
N2—C12	1.280 (8)	C16—H16	0.9300
N2—C11	1.465 (9)	C6—C7	1.359 (10)
N3—O7	1.221 (7)	C6—H6	0.9300
N3—O6	1.255 (8)	C3—C8	1.380 (9)
N3—O5	1.279 (8)	C11—C10	1.496 (10)
C12—C13	1.445 (9)	C11—H11A	0.9700
C12—H12	0.9300	C11—H11B	0.9700
O3—C3	1.392 (7)	C10—H10A	0.9700
O3—C2	1.466 (7)	C10—H10B	0.9700
C18—C13	1.393 (8)	C20—H20A	0.9600
C18—C17	1.406 (9)	C20—H20B	0.9600
O4—C17	1.383 (7)	C20—H20C	0.9600
O4—C19	1.449 (8)	C8—C7	1.396 (10)
C15—C14	1.348 (10)	C8—H8	0.9300
C15—C16	1.382 (10)	C7—H7	0.9300
C6—C5—C4	119.3 (6)	C19—O4—Ho1	123.5 (4)
C6—C5—C9	118.2 (6)	N3—O5—Ho1	96.9 (4)
C4—C5—C9	122.4 (6)	C14—C15—C16	121.8 (6)

O2—Ho1—O1	62.44 (14)	C14—C15—H15	119.1
O2—Ho1—O12	142.78 (15)	C16—C15—H15	119.1
O1—Ho1—O12	116.07 (17)	O2—C4—C3	118.9 (5)
O2—Ho1—O5	125.12 (15)	O2—C4—C5	122.1 (6)
O1—Ho1—O5	114.48 (17)	C3—C4—C5	118.9 (5)
O12—Ho1—O5	90.49 (18)	O4—C19—C20	111.9 (6)
O2—Ho1—O8	73.78 (15)	O4—C19—H19A	109.2
O1—Ho1—O8	112.76 (15)	C20—C19—H19A	109.2
O12—Ho1—O8	73.62 (17)	O4—C19—H19B	109.2
O5—Ho1—O8	132.44 (19)	C20—C19—H19B	109.2
O2—Ho1—O6	80.15 (16)	H19A—C19—H19B	107.9
O1—Ho1—O6	71.83 (16)	C2—C1—H1A	109.5
O12—Ho1—O6	136.61 (17)	C2—C1—H1B	109.5
O5—Ho1—O6	51.82 (17)	H1A—C1—H1B	109.5
O8—Ho1—O6	146.11 (17)	C2—C1—H1C	109.5
O2—Ho1—O9	76.56 (16)	H1A—C1—H1C	109.5
O1—Ho1—O9	69.75 (16)	H1B—C1—H1C	109.5
O12—Ho1—O9	69.29 (18)	O4—C17—C16	125.6 (6)
O5—Ho1—O9	157.98 (18)	O4—C17—C18	113.8 (5)
O8—Ho1—O9	51.54 (17)	C16—C17—C18	120.6 (6)
O6—Ho1—O9	141.01 (17)	C18—C13—C14	119.1 (6)
O2—Ho1—O11	131.15 (18)	C18—C13—C12	122.4 (6)
O1—Ho1—O11	165.27 (17)	C14—C13—C12	118.5 (6)
O12—Ho1—O11	50.32 (17)	C15—C14—C13	119.9 (6)
O5—Ho1—O11	64.50 (19)	C15—C14—H14	120.0
O8—Ho1—O11	71.28 (19)	C13—C14—H14	120.0
O6—Ho1—O11	113.28 (17)	O3—C2—C1	111.0 (5)
O9—Ho1—O11	105.60 (18)	O3—C2—H2A	109.4
O2—Ho1—O3	63.79 (14)	C1—C2—H2A	109.4
O1—Ho1—O3	121.17 (14)	O3—C2—H2B	109.4
O12—Ho1—O3	120.80 (16)	C1—C2—H2B	109.4
O5—Ho1—O3	79.44 (17)	H2A—C2—H2B	108.0
O8—Ho1—O3	71.93 (16)	N4—O9—Ho1	94.8 (4)
O6—Ho1—O3	77.33 (17)	O10—N4—O9	123.1 (7)
O9—Ho1—O3	117.91 (17)	O10—N4—O8	118.8 (7)
O11—Ho1—O3	73.51 (16)	O9—N4—O8	118.1 (6)
O2—Ho1—O4	124.19 (15)	N4—O8—Ho1	95.5 (4)
O1—Ho1—O4	61.83 (14)	N1—C9—C5	124.2 (6)
O12—Ho1—O4	69.37 (16)	N1—C9—H9	117.9
O5—Ho1—O4	77.05 (16)	C5—C9—H9	117.9
O8—Ho1—O4	132.43 (16)	N5—O12—Ho1	99.7 (4)
O6—Ho1—O4	80.41 (17)	O13—N5—O11	123.6 (8)
O9—Ho1—O4	87.35 (17)	O13—N5—O12	121.4 (7)
O11—Ho1—O4	104.59 (17)	O11—N5—O12	114.9 (6)
O3—Ho1—O4	154.49 (18)	C15—C16—C17	119.3 (7)
N1—Ni1—N2	86.1 (3)	C15—C16—H16	120.3
N1—Ni1—O1	175.1 (2)	C17—C16—H16	120.3
N2—Ni1—O1	96.0 (2)	C7—C6—C5	120.0 (6)
N1—Ni1—O2	94.9 (2)	C7—C6—H6	120.0

supplementary materials


N2—Ni1—O2	175.7 (2)	C5—C6—H6	120.0
O1—Ni1—O2	83.27 (17)	C8—C3—O3	126.2 (6)
C4—O2—Ni1	126.1 (4)	C8—C3—C4	120.9 (6)
C4—O2—Ho1	125.1 (4)	O3—C3—C4	112.9 (5)
Ni1—O2—Ho1	107.79 (18)	N2—C11—C10	105.8 (5)
C18—O1—Ni1	126.2 (4)	N2—C11—H11A	110.6
C18—O1—Ho1	127.8 (4)	C10—C11—H11A	110.6
Ni1—O1—Ho1	105.95 (17)	N2—C11—H11B	110.6
C9—N1—C10	121.8 (5)	C10—C11—H11B	110.6
C9—N1—Ni1	126.3 (5)	H11A—C11—H11B	108.7
C10—N1—Ni1	111.9 (4)	N1—C10—C11	106.7 (5)
C12—N2—C11	119.2 (6)	N1—C10—H10A	110.4
C12—N2—Ni1	126.7 (5)	C11—C10—H10A	110.4
C11—N2—Ni1	113.9 (4)	N1—C10—H10B	110.4
O7—N3—O6	122.0 (7)	C11—C10—H10B	110.4
O7—N3—O5	122.8 (7)	H10A—C10—H10B	108.6
O6—N3—O5	115.3 (6)	N5—O11—Ho1	95.0 (4)
N3—O6—Ho1	95.8 (4)	C19—C20—H20A	109.5
N2—C12—C13	124.7 (6)	C19—C20—H20B	109.5
N2—C12—H12	117.7	H20A—C20—H20B	109.5
C13—C12—H12	117.7	C19—C20—H20C	109.5
C3—O3—C2	117.3 (5)	H20A—C20—H20C	109.5
C3—O3—Ho1	118.6 (3)	H20B—C20—H20C	109.5
C2—O3—Ho1	124.0 (4)	C3—C8—C7	119.0 (6)
O1—C18—C13	123.9 (6)	C3—C8—H8	120.5
O1—C18—C17	117.0 (5)	C7—C8—H8	120.5
C13—C18—C17	119.1 (6)	C6—C7—C8	121.7 (6)
C17—O4—C19	117.0 (5)	C6—C7—H7	119.1
C17—O4—Ho1	119.5 (4)	C8—C7—H7	119.1

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C20—H20A...O12	0.96	2.41	3.088 (9)	127
C12—H12...O10 ⁱ	0.93	2.37	3.286 (8)	169
C10—H10A...O10 ⁱⁱ	0.97	2.42	3.301 (9)	150
C2—H2B...O11	0.97	2.59	3.015 (9)	107
C1—H1C...O11	0.96	2.52	3.173 (10)	125


Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $x-1, y, z$.

Fig. 1



Article

Fig. 2



ArXiv