

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 3,4-Dihydroxybenzaldehyde thiosemi-carbazone

 Kong Wai Tan,<sup>a</sup> Yang Farina,<sup>b</sup> Chew Hee Ng,<sup>c\*</sup>  
 Mohd Jamil Maah<sup>a</sup> and Seik Weng Ng<sup>a</sup>

<sup>a</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, <sup>b</sup>School of Chemical Science and Food Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Malaysia, and <sup>c</sup>Faculty of Engineering and Science, Universiti Tunku Abdul Rahman, 53300 Kuala Lumpur, Malaysia  
 Correspondence e-mail: seikweng@um.edu.my


Received 6 May 2008; accepted 9 May 2008

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.061;  $wR$  factor = 0.190; data-to-parameter ratio = 15.9.

The asymmetric unit of the title compound,  $\text{C}_8\text{H}_9\text{N}_3\text{O}_2\text{S}$ , contains three independent molecules which are stacked approximately over each other. In the crystal structure, centrosymmetric pairs of molecules are formed through intermolecular hydroxy–hydroxy  $\text{O}-\text{H}\cdots\text{O}$  and hydroxy–sulfur  $\text{O}-\text{H}\cdots\text{S}$  hydrogen bonds which are, in turn, linked into a two-dimensional network by  $\text{N}-\text{H}\cdots\text{O}$  (hydroxy) hydrogen bonds.

### Related literature

For the structure of 3,4-dihydroxybenzaldehyde 4-phenylthiosemicarbazone, see: Swesi *et al.* (2006). For some metal complexes of the ligand, see: Zhu *et al.* (1991, 1997).



### Experimental

#### Crystal data

$\text{C}_8\text{H}_9\text{N}_3\text{O}_2\text{S}$   
 $M_r = 211.24$   
 Triclinic,  $P\bar{1}$   
 $a = 10.657$  (2) Å  
 $b = 11.794$  (2) Å  
 $c = 12.356$  (2) Å

$\alpha = 111.657$  (2)°  
 $\beta = 104.082$  (2)°  
 $\gamma = 90.929$  (2)°  
 $V = 1390.2$  (4) Å<sup>3</sup>  
 $Z = 6$   
 Mo  $K\alpha$  radiation

$\mu = 0.33$  mm<sup>-1</sup>  
 $T = 100$  (2) K

$0.20 \times 0.18 \times 0.04$  mm

#### Data collection

Bruker SMART APEX  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.938$ ,  $T_{\max} = 0.987$

8792 measured reflections  
 6298 independent reflections  
 3727 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.189$   
 $S = 1.01$   
 6298 reflections  
 397 parameters  
 6 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.51$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.40$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1o}\cdots\text{O6}^i$	0.84 (1)	2.07 (3)	2.784 (3)	143 (4)
$\text{O2}-\text{H2o}\cdots\text{S1}^{ii}$	0.84 (1)	2.47 (1)	3.300 (2)	171 (4)
$\text{N1}-\text{H1n2}\cdots\text{O5}^{iii}$	0.88	2.00	2.856 (4)	163
$\text{O3}-\text{H3o}\cdots\text{O4}^i$	0.84 (1)	2.11 (4)	2.732 (3)	130 (4)
$\text{O4}-\text{H4o}\cdots\text{S2}^{ii}$	0.84 (1)	2.38 (1)	3.219 (2)	174 (4)
$\text{N4}-\text{H4n2}\cdots\text{O3}^{iii}$	0.88	2.05	2.900 (4)	162
$\text{O5}-\text{H5o}\cdots\text{O2}^i$	0.84 (1)	2.16 (4)	2.742 (3)	127 (4)
$\text{O6}-\text{H6o}\cdots\text{S3}^{ii}$	0.84 (1)	2.40 (1)	3.244 (2)	177 (4)
$\text{N7}-\text{H7n2}\cdots\text{O1}^{iii}$	0.88	2.13	2.981 (4)	161

Symmetry codes: (i)  $-x+1, -y+1, -z+2$ ; (ii)  $x, y, z+1$ ; (iii)  $-x+1, -y+1, -z+1$ .

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

We thank the University of Malaya (P0265/2007A) for supporting this study. KWT thanks the Ministry of Higher Education for a SLAI scholarship in this research.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2626).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Swesi, A. T., Farina, Y., Kassim, M. & Ng, S. W. (2006). *Acta Cryst.* **E62**, o5457–o5458.  
 Westrip, S. P. (2008). *publCIF*. In preparation.  
 Zhu, X.-D., Wang, C.-G., Le, Z.-F., Mei, S.-Y., Yen, Z.-H. & Wu, Z.-S. (1991). *Synth. React. Inorg. Met. Org. Chem.* **21**, 1365–1373.  
 Zhu, X.-D., Wang, C.-G., Lu, Z.-P. & Dang, Y.-L. (1997). *Transition Met. Chem.* **22**, 9–13.

**supplementary materials**

*Acta Cryst.* (2008). E64, o1073 [ doi:10.1107/S160053680801386X ]

### 3,4-Dihydroxybenzaldehyde thiosemicarbazone

K. W. Tan, Y. Farina, C. H. Ng, M. J. Maah and S. W. Ng

#### Comment

A previous study of Schiff bases derived by condensing substituted benzaldehydes with 4-phenylthiosemicarbazides describes the 3,4-dihydroxybenzaldehyde derivative, which crystallizes as a hemihydrate. The compound features extensive hydrogen bonds (Swesi *et al.*, 2006). The condensation product of the reaction between thiosemicarbazide and 3,4-dihydroxybenzaldehyde has an amino  $-NH_2$  group in place of the phenyl group. In the crystal structure, a molecule is linked to an adjacent molecule by a hydrogen bond [ $O-H_{3\text{-hydroxy}}\cdots O_{4\text{-hydroxy}}$ ]; it is linked to another adjacent molecule by another hydrogen bond [ $O-H_{4\text{-hydroxy}}\cdots S$ ]. The structure is consolidated into a two-dimensional network motif by a  $N_{\text{terminal}}-H\cdots O_{4\text{-hydroxy}}$  hydrogen bond. The asymmetric unit features three molecules that are approximately stacked over each other (Fig. 1).

#### Experimental

Thiosemicarbazide (0.09 g, 1 mmol) and 2,4-dihydroxybenzaldehyde (0.14 g, 1 mmol) were heated in an ethanol/water mixture (20/5 ml) for 3 h. Slow evaporation of the solvent yielded yellow crystals.

#### Refinement

Carbon-bound H-atoms were placed in calculated positions ( $C-H$  0.95 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to 1.2  $U(C)$ . The amino H-atoms were similarly treated ( $N-H$  0.88 Å). The hydroxy H-atoms were located in a difference Fourier map, and were refined with a distance restraint of  $O-H$  0.85±0.01 Å; their temperature factors were tied by a factor of 1.5.

#### Figures




Fig. 1. Thermal ellipsoid (Barbour, 2001) plot of the three independent molecules of  $C_{14}H_{13}N_3O_2S$  at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radii.

### 3,4-Dihydroxybenzaldehyde thiosemicarbazone

#### Crystal data

$C_8H_9N_3O_2S$

$M_r = 211.24$

Triclinic,  $P\bar{1}$

$Z = 6$

$F_{000} = 660$

$D_x = 1.514 \text{ Mg m}^{-3}$

# supplementary materials

---

Hall symbol: -P 1

$a = 10.657(2) \text{ \AA}$

$b = 11.794(2) \text{ \AA}$

$c = 12.356(2) \text{ \AA}$

$\alpha = 111.657(2)^\circ$

$\beta = 104.082(2)^\circ$

$\gamma = 90.929(2)^\circ$

$V = 1390.2(4) \text{ \AA}^3$

Mo  $K\alpha$  radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1478 reflections

$\theta = 2.7\text{--}27.8^\circ$

$\mu = 0.33 \text{ mm}^{-1}$

$T = 100(2) \text{ K}$

Block, yellow

$0.20 \times 0.18 \times 0.04 \text{ mm}$

## Data collection

Bruker SMART APEX  
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100(2) \text{ K}$

$\omega$  scans

Absorption correction: multi-scan  
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.938$ ,  $T_{\max} = 0.987$

8792 measured reflections

6298 independent reflections

3727 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 1.8^\circ$

$h = -13 \rightarrow 13$

$k = -15 \rightarrow 9$

$l = -14 \rightarrow 16$

## Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.061$

$wR(F^2) = 0.189$

$S = 1.01$

6298 reflections

397 parameters

6 restraints

Primary atom site location: structure-invariant direct  
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring  
sites

H atoms treated by a mixture of  
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.1004P)^2]$$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.51 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.39 \text{ e \AA}^{-3}$

Extinction correction: none

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.84547 (10)	0.94314 (8)	0.21803 (8)	0.0401 (3)
S2	0.45653 (10)	0.85179 (8)	0.31556 (7)	0.0359 (2)
S3	0.10237 (10)	0.74686 (8)	0.41416 (8)	0.0390 (3)
O1	0.7414 (3)	0.5896 (2)	0.7057 (2)	0.0440 (7)
H1O	0.750 (4)	0.566 (4)	0.763 (3)	0.066*
O2	0.7519 (2)	0.7405 (2)	0.9368 (2)	0.0333 (6)

H2O	0.769 (4)	0.797 (3)	1.0057 (17)	0.050*
O3	0.4715 (4)	0.5066 (2)	0.8312 (2)	0.0615 (9)
H3O	0.476 (5)	0.496 (5)	0.895 (3)	0.092*
O4	0.4659 (3)	0.6536 (2)	1.05646 (19)	0.0359 (6)
H4O	0.464 (4)	0.710 (3)	1.122 (2)	0.054*
O5	0.2270 (3)	0.4329 (2)	0.9629 (2)	0.0441 (7)
H5O	0.232 (4)	0.426 (4)	1.029 (2)	0.066*
O6	0.1658 (3)	0.5600 (2)	1.1704 (2)	0.0370 (6)
H6O	0.152 (4)	0.609 (3)	1.235 (2)	0.056*
N1	0.7685 (3)	0.7537 (3)	0.2628 (2)	0.0407 (8)
H1N1	0.7486	0.7210	0.3109	0.049*
H1N2	0.7622	0.7072	0.1868	0.049*
N2	0.8145 (3)	0.9368 (2)	0.4209 (2)	0.0328 (7)
H2N	0.8332	1.0172	0.4520	0.039*
N3	0.7922 (3)	0.8755 (2)	0.4912 (2)	0.0316 (6)
N4	0.4570 (3)	0.6598 (3)	0.3786 (3)	0.0438 (8)
H4N1	0.4531	0.6259	0.4305	0.053*
H4N2	0.4637	0.6141	0.3061	0.053*
N5	0.4424 (3)	0.8421 (2)	0.5209 (2)	0.0294 (6)
H5N	0.4379	0.9218	0.5460	0.035*
N6	0.4390 (3)	0.7808 (2)	0.5960 (2)	0.0282 (6)
N7	0.1429 (3)	0.5636 (3)	0.4892 (3)	0.0470 (9)
H7N1	0.1457	0.5309	0.5431	0.056*
H7N2	0.1607	0.5210	0.4205	0.056*
N8	0.0866 (3)	0.7341 (3)	0.6182 (2)	0.0334 (7)
H8N	0.0617	0.8082	0.6370	0.040*
N9	0.1000 (3)	0.6760 (2)	0.6982 (2)	0.0299 (6)
C1	0.8075 (3)	0.8720 (3)	0.3043 (3)	0.0300 (7)
C2	0.7942 (3)	0.9415 (3)	0.5994 (3)	0.0306 (7)
H2	0.8072	1.0284	0.6258	0.037*
C3	0.7772 (3)	0.8876 (3)	0.6840 (3)	0.0279 (7)
C4	0.7636 (3)	0.7614 (3)	0.6542 (3)	0.0296 (7)
H4	0.7605	0.7070	0.5743	0.035*
C5	0.7545 (3)	0.7147 (3)	0.7389 (3)	0.0293 (7)
C6	0.7597 (3)	0.7934 (3)	0.8565 (3)	0.0257 (7)
C7	0.7714 (3)	0.9180 (3)	0.8871 (3)	0.0319 (8)
H7C	0.7737	0.9719	0.9670	0.038*
C8	0.7799 (3)	0.9652 (3)	0.8013 (3)	0.0327 (8)
H8	0.7875	1.0516	0.8228	0.039*
C9	0.4526 (3)	0.7787 (3)	0.4095 (3)	0.0273 (7)
C10	0.4204 (3)	0.8443 (3)	0.6983 (3)	0.0288 (7)
H10	0.4054	0.9277	0.7164	0.035*
C11	0.4218 (3)	0.7917 (3)	0.7876 (3)	0.0252 (7)
C12	0.4409 (3)	0.6698 (3)	0.7657 (3)	0.0298 (7)
H12	0.4459	0.6164	0.6879	0.036*
C13	0.4526 (3)	0.6261 (3)	0.8557 (3)	0.0321 (8)
C14	0.4480 (3)	0.7039 (3)	0.9711 (3)	0.0257 (7)
C15	0.4243 (3)	0.8234 (3)	0.9921 (3)	0.0316 (8)
H15	0.4172	0.8760	1.0694	0.038*

## supplementary materials

---

C16	0.4107 (3)	0.8671 (3)	0.9006 (3)	0.0304 (7)
H16	0.3936	0.9495	0.9153	0.036*
C17	0.1122 (3)	0.6761 (3)	0.5111 (3)	0.0306 (7)
C18	0.0760 (3)	0.7363 (3)	0.7984 (3)	0.0286 (7)
H18	0.0465	0.8147	0.8120	0.034*
C19	0.0928 (3)	0.6872 (3)	0.8926 (3)	0.0264 (7)
C20	0.1458 (3)	0.5772 (3)	0.8812 (3)	0.0288 (7)
H20	0.1662	0.5295	0.8081	0.035*
C21	0.1686 (3)	0.5376 (3)	0.9745 (3)	0.0291 (7)
C22	0.1375 (3)	0.6051 (3)	1.0814 (3)	0.0270 (7)
C23	0.0804 (3)	0.7116 (3)	1.0922 (3)	0.0295 (7)
H23	0.0556	0.7566	1.1637	0.035*
C24	0.0594 (3)	0.7524 (3)	0.9982 (3)	0.0300 (7)
H24	0.0214	0.8264	1.0064	0.036*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0718 (7)	0.0287 (5)	0.0288 (5)	0.0104 (4)	0.0234 (4)	0.0146 (4)
S2	0.0636 (6)	0.0242 (4)	0.0253 (4)	0.0079 (4)	0.0172 (4)	0.0123 (4)
S3	0.0631 (6)	0.0330 (5)	0.0286 (5)	0.0124 (4)	0.0187 (4)	0.0164 (4)
O1	0.083 (2)	0.0205 (12)	0.0354 (14)	0.0112 (13)	0.0242 (14)	0.0123 (11)
O2	0.0527 (15)	0.0269 (13)	0.0263 (12)	0.0067 (11)	0.0144 (11)	0.0144 (10)
O3	0.137 (3)	0.0278 (15)	0.0344 (15)	0.0361 (17)	0.0385 (18)	0.0173 (13)
O4	0.0637 (17)	0.0257 (13)	0.0238 (12)	0.0131 (12)	0.0166 (12)	0.0123 (10)
O5	0.083 (2)	0.0266 (13)	0.0343 (14)	0.0200 (13)	0.0278 (14)	0.0167 (12)
O6	0.0633 (17)	0.0276 (13)	0.0252 (12)	0.0146 (12)	0.0162 (12)	0.0127 (11)
N1	0.075 (2)	0.0255 (16)	0.0245 (15)	0.0053 (15)	0.0198 (15)	0.0091 (13)
N2	0.0565 (19)	0.0214 (14)	0.0249 (14)	0.0070 (13)	0.0161 (13)	0.0105 (12)
N3	0.0492 (18)	0.0255 (15)	0.0270 (14)	0.0069 (13)	0.0133 (13)	0.0156 (12)
N4	0.085 (3)	0.0232 (16)	0.0285 (16)	0.0122 (16)	0.0230 (16)	0.0106 (13)
N5	0.0500 (18)	0.0194 (13)	0.0235 (14)	0.0081 (12)	0.0140 (12)	0.0107 (11)
N6	0.0444 (17)	0.0221 (14)	0.0230 (13)	0.0032 (12)	0.0105 (12)	0.0131 (11)
N7	0.088 (3)	0.0285 (17)	0.0371 (17)	0.0156 (17)	0.0319 (17)	0.0165 (14)
N8	0.0529 (19)	0.0271 (15)	0.0302 (15)	0.0141 (13)	0.0172 (13)	0.0179 (13)
N9	0.0426 (17)	0.0239 (14)	0.0293 (14)	0.0056 (12)	0.0133 (12)	0.0147 (12)
C1	0.043 (2)	0.0262 (17)	0.0261 (17)	0.0123 (15)	0.0121 (15)	0.0143 (15)
C2	0.044 (2)	0.0221 (16)	0.0274 (17)	0.0042 (15)	0.0093 (15)	0.0119 (14)
C3	0.0387 (19)	0.0247 (17)	0.0249 (16)	0.0075 (14)	0.0106 (14)	0.0134 (14)
C4	0.042 (2)	0.0221 (16)	0.0261 (17)	0.0081 (14)	0.0122 (14)	0.0087 (14)
C5	0.0412 (19)	0.0190 (16)	0.0316 (17)	0.0082 (14)	0.0128 (15)	0.0119 (14)
C6	0.0326 (18)	0.0267 (17)	0.0235 (16)	0.0092 (14)	0.0116 (13)	0.0132 (14)
C7	0.050 (2)	0.0246 (17)	0.0223 (16)	0.0063 (15)	0.0118 (15)	0.0086 (14)
C8	0.049 (2)	0.0179 (16)	0.0316 (18)	0.0037 (15)	0.0121 (16)	0.0094 (14)
C9	0.0363 (18)	0.0234 (17)	0.0245 (16)	0.0080 (14)	0.0089 (14)	0.0111 (14)
C10	0.0357 (19)	0.0256 (17)	0.0293 (17)	0.0079 (14)	0.0114 (14)	0.0137 (14)
C11	0.0315 (17)	0.0240 (16)	0.0250 (16)	0.0059 (13)	0.0099 (13)	0.0133 (14)
C12	0.046 (2)	0.0250 (17)	0.0211 (15)	0.0092 (15)	0.0134 (14)	0.0092 (14)

C13	0.050 (2)	0.0209 (16)	0.0306 (17)	0.0120 (15)	0.0156 (15)	0.0126 (14)
C14	0.0371 (18)	0.0245 (16)	0.0211 (15)	0.0062 (14)	0.0111 (13)	0.0128 (13)
C15	0.047 (2)	0.0236 (17)	0.0259 (17)	0.0085 (15)	0.0163 (15)	0.0074 (14)
C16	0.048 (2)	0.0191 (16)	0.0296 (17)	0.0078 (14)	0.0133 (15)	0.0132 (14)
C17	0.0397 (19)	0.0259 (17)	0.0288 (17)	0.0040 (15)	0.0122 (15)	0.0116 (15)
C18	0.0333 (18)	0.0267 (17)	0.0289 (17)	0.0061 (14)	0.0096 (14)	0.0131 (14)
C19	0.0337 (18)	0.0221 (16)	0.0242 (16)	0.0019 (13)	0.0079 (13)	0.0098 (13)
C20	0.042 (2)	0.0212 (16)	0.0262 (16)	0.0048 (14)	0.0140 (14)	0.0094 (14)
C21	0.0408 (19)	0.0205 (16)	0.0300 (17)	0.0070 (14)	0.0141 (14)	0.0111 (14)
C22	0.0361 (18)	0.0240 (16)	0.0242 (16)	0.0036 (14)	0.0103 (14)	0.0115 (14)
C23	0.0405 (19)	0.0252 (17)	0.0245 (16)	0.0092 (15)	0.0131 (14)	0.0084 (14)
C24	0.0375 (19)	0.0248 (17)	0.0281 (17)	0.0067 (14)	0.0089 (14)	0.0103 (14)

*Geometric parameters (Å, °)*

S1—C1	1.693 (3)	N9—C18	1.270 (4)
S2—C9	1.689 (3)	C2—C3	1.453 (4)
S3—C17	1.680 (3)	C2—H2	0.9500
O1—C5	1.372 (4)	C3—C4	1.391 (4)
O1—H10	0.837 (10)	C3—C8	1.393 (4)
O2—C6	1.369 (4)	C4—C5	1.372 (4)
O2—H2O	0.840 (10)	C4—H4	0.9500
O3—C13	1.358 (4)	C5—C6	1.392 (4)
O3—H3O	0.836 (10)	C6—C7	1.371 (4)
O4—C14	1.367 (4)	C7—C8	1.387 (4)
O4—H4O	0.841 (10)	C7—H7C	0.9500
O5—C21	1.368 (4)	C8—H8	0.9500
O5—H5O	0.838 (10)	C10—C11	1.450 (4)
O6—C22	1.364 (4)	C10—H10	0.9500
O6—H6O	0.844 (10)	C11—C16	1.385 (4)
N1—C1	1.316 (4)	C11—C12	1.390 (4)
N1—H1N1	0.8800	C12—C13	1.369 (4)
N1—H1N2	0.8800	C12—H12	0.9500
N2—C1	1.341 (4)	C13—C14	1.396 (4)
N2—N3	1.375 (3)	C14—C15	1.375 (4)
N2—H2N	0.8800	C15—C16	1.385 (4)
N3—C2	1.269 (4)	C15—H15	0.9500
N4—C9	1.316 (4)	C16—H16	0.9500
N4—H4N1	0.8800	C18—C19	1.457 (4)
N4—H4N2	0.8800	C18—H18	0.9500
N5—C9	1.335 (4)	C19—C24	1.384 (4)
N5—N6	1.377 (3)	C19—C20	1.397 (4)
N5—H5N	0.8800	C20—C21	1.368 (4)
N6—C10	1.276 (4)	C20—H20	0.9500
N7—C17	1.316 (4)	C21—C22	1.390 (4)
N7—H7N1	0.8800	C22—C23	1.380 (4)
N7—H7N2	0.8800	C23—C24	1.386 (4)
N8—C17	1.344 (4)	C23—H23	0.9500
N8—N9	1.380 (3)	C24—H24	0.9500

## supplementary materials

---

N8—H8N	0.8800		
C5—O1—H1O	115 (3)	C3—C8—H8	119.6
C6—O2—H2O	107 (3)	N4—C9—N5	116.7 (3)
C13—O3—H3O	106 (4)	N4—C9—S2	123.5 (2)
C14—O4—H4O	107 (3)	N5—C9—S2	119.8 (2)
C21—O5—H5O	103 (3)	N6—C10—C11	121.2 (3)
C22—O6—H6O	112 (3)	N6—C10—H10	119.4
C1—N1—H1N1	120.0	C11—C10—H10	119.4
C1—N1—H1N2	120.0	C16—C11—C12	119.0 (3)
H1N1—N1—H1N2	120.0	C16—C11—C10	118.9 (3)
C1—N2—N3	118.9 (3)	C12—C11—C10	122.0 (3)
C1—N2—H2N	120.5	C13—C12—C11	120.3 (3)
N3—N2—H2N	120.5	C13—C12—H12	119.8
C2—N3—N2	116.4 (3)	C11—C12—H12	119.8
C9—N4—H4N1	120.0	O3—C13—C12	118.8 (3)
C9—N4—H4N2	120.0	O3—C13—C14	120.7 (3)
H4N1—N4—H4N2	120.0	C12—C13—C14	120.5 (3)
C9—N5—N6	118.9 (3)	O4—C14—C15	124.2 (3)
C9—N5—H5N	120.6	O4—C14—C13	116.4 (3)
N6—N5—H5N	120.6	C15—C14—C13	119.4 (3)
C10—N6—N5	116.1 (3)	C14—C15—C16	120.0 (3)
C17—N7—H7N1	120.0	C14—C15—H15	120.0
C17—N7—H7N2	120.0	C16—C15—H15	120.0
H7N1—N7—H7N2	120.0	C15—C16—C11	120.6 (3)
C17—N8—N9	119.0 (3)	C15—C16—H16	119.7
C17—N8—H8N	120.5	C11—C16—H16	119.7
N9—N8—H8N	120.5	N7—C17—N8	115.9 (3)
C18—N9—N8	115.9 (3)	N7—C17—S3	124.0 (3)
N1—C1—N2	116.6 (3)	N8—C17—S3	120.0 (3)
N1—C1—S1	123.3 (2)	N9—C18—C19	121.1 (3)
N2—C1—S1	120.1 (2)	N9—C18—H18	119.5
N3—C2—C3	121.7 (3)	C19—C18—H18	119.5
N3—C2—H2	119.2	C24—C19—C20	118.6 (3)
C3—C2—H2	119.2	C24—C19—C18	119.6 (3)
C4—C3—C8	118.5 (3)	C20—C19—C18	121.8 (3)
C4—C3—C2	122.7 (3)	C21—C20—C19	120.4 (3)
C8—C3—C2	118.8 (3)	C21—C20—H20	119.8
C5—C4—C3	120.7 (3)	C19—C20—H20	119.8
C5—C4—H4	119.7	O5—C21—C20	119.0 (3)
C3—C4—H4	119.7	O5—C21—C22	120.3 (3)
O1—C5—C4	118.7 (3)	C20—C21—C22	120.6 (3)
O1—C5—C6	121.0 (3)	O6—C22—C23	123.8 (3)
C4—C5—C6	120.3 (3)	O6—C22—C21	116.6 (3)
O2—C6—C7	123.0 (3)	C23—C22—C21	119.6 (3)
O2—C6—C5	117.1 (3)	C22—C23—C24	119.5 (3)
C7—C6—C5	119.9 (3)	C22—C23—H23	120.2
C6—C7—C8	119.8 (3)	C24—C23—H23	120.2
C6—C7—H7C	120.1	C19—C24—C23	121.2 (3)
C8—C7—H7C	120.1	C19—C24—H24	119.4

C7—C8—C3	120.8 (3)	C23—C24—H24	119.4
C7—C8—H8	119.6		
C1—N2—N3—C2	-176.9 (3)	C11—C12—C13—C14	-1.3 (5)
C9—N5—N6—C10	-175.5 (3)	O3—C13—C14—O4	1.2 (5)
C17—N8—N9—C18	179.2 (3)	C12—C13—C14—O4	-177.4 (3)
N3—N2—C1—N1	5.5 (5)	O3—C13—C14—C15	-177.8 (3)
N3—N2—C1—S1	-175.4 (2)	C12—C13—C14—C15	3.7 (5)
N2—N3—C2—C3	-177.5 (3)	O4—C14—C15—C16	178.4 (3)
N3—C2—C3—C4	2.9 (5)	C13—C14—C15—C16	-2.7 (5)
N3—C2—C3—C8	-179.7 (3)	C14—C15—C16—C11	-0.5 (5)
C8—C3—C4—C5	-0.8 (5)	C12—C11—C16—C15	2.9 (5)
C2—C3—C4—C5	176.6 (3)	C10—C11—C16—C15	-173.6 (3)
C3—C4—C5—O1	-179.9 (3)	N9—N8—C17—N7	3.0 (5)
C3—C4—C5—C6	-0.4 (5)	N9—N8—C17—S3	-177.6 (2)
O1—C5—C6—O2	0.4 (5)	N8—N9—C18—C19	-177.0 (3)
C4—C5—C6—O2	-179.1 (3)	N9—C18—C19—C24	-176.7 (3)
O1—C5—C6—C7	-179.3 (3)	N9—C18—C19—C20	5.3 (5)
C4—C5—C6—C7	1.2 (5)	C24—C19—C20—C21	-2.5 (5)
O2—C6—C7—C8	179.5 (3)	C18—C19—C20—C21	175.5 (3)
C5—C6—C7—C8	-0.9 (5)	C19—C20—C21—O5	-176.6 (3)
C6—C7—C8—C3	-0.3 (5)	C19—C20—C21—C22	1.1 (5)
C4—C3—C8—C7	1.1 (5)	O5—C21—C22—O6	-1.2 (5)
C2—C3—C8—C7	-176.4 (3)	C20—C21—C22—O6	-178.8 (3)
N6—N5—C9—N4	0.9 (5)	O5—C21—C22—C23	179.0 (3)
N6—N5—C9—S2	179.8 (2)	C20—C21—C22—C23	1.4 (5)
N5—N6—C10—C11	-176.7 (3)	O6—C22—C23—C24	177.8 (3)
N6—C10—C11—C16	175.3 (3)	C21—C22—C23—C24	-2.5 (5)
N6—C10—C11—C12	-1.1 (5)	C20—C19—C24—C23	1.4 (5)
C16—C11—C12—C13	-2.0 (5)	C18—C19—C24—C23	-176.6 (3)
C10—C11—C12—C13	174.4 (3)	C22—C23—C24—C19	1.0 (5)
C11—C12—C13—O3	-179.9 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O1—H1o...O6 <sup>i</sup>	0.84 (1)	2.07 (3)	2.784 (3)	143 (4)
O2—H2o...S1 <sup>ii</sup>	0.84 (1)	2.47 (1)	3.300 (2)	171 (4)
N1—H1n2...O5 <sup>iii</sup>	0.88	2.00	2.856 (4)	163
O3—H3o...O4 <sup>i</sup>	0.84 (1)	2.11 (4)	2.732 (3)	130 (4)
O4—H4o...S2 <sup>ii</sup>	0.84 (1)	2.38 (1)	3.219 (2)	174 (4)
N4—H4n2...O3 <sup>iii</sup>	0.88	2.05	2.900 (4)	162
O5—H5o...O2 <sup>i</sup>	0.84 (1)	2.16 (4)	2.742 (3)	127 (4)
O6—H6o...S3 <sup>ii</sup>	0.84 (1)	2.40 (1)	3.244 (2)	177 (4)
N7—H7n2...O1 <sup>iii</sup>	0.88	2.13	2.981 (4)	161

Symmetry codes: (i)  $-x+1, -y+1, -z+2$ ; (ii)  $x, y, z+1$ ; (iii)  $-x+1, -y+1, -z+1$ .

Fig. 1

