

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-Methyl-1*H*-benzimidazole-2(3*H*)-thione

 Hizbullah Khan,^a Amin Badshah,^{a*} Farkhanda Shaheen,^a Christine Gieck^b and Rizwana Aleem Qureshi^c
^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad, Pakistan, ^bDISTA, Università del Piemonte Orientale, Alessandria I-15100, Italy, and ^cDepartment of Plant Sciences, Faculty of Biological Sciences, Quaid-i-Azam University, Islamabad 45320, Pakistan

Correspondence e-mail: aminbadshah@yahoo.com


Received 21 April 2008; accepted 18 May 2008

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.029; wR factor = 0.078; data-to-parameter ratio = 9.5.

The title compound, $\text{C}_8\text{H}_8\text{N}_2\text{S}$, was prepared by the condensation of *N*-methyl-1,2-phenylenediamine and carbon disulfide. The crystal structure is stabilized by a $\text{C}-\text{H}\cdots\pi$ interaction between a benzene H atom and the benzene ring of a neighbouring molecule, and by intermolecular $\text{N}-\text{H}\cdots\text{S}$ interactions.

Related literature

For related literature, see: Baily *et al.* (1996); Koch (2001); Namgun *et al.* (2001); Schuster *et al.* (1990); Patel & Chedekel (1984).



Experimental

Crystal data

 $\text{C}_8\text{H}_8\text{N}_2\text{S}$
 $M_r = 164.22$

 Monoclinic, $P2_1/n$
 $a = 9.997$ (4) Å

 $b = 5.8140$ (7) Å
 $c = 13.703$ (4) Å
 $\beta = 94.05$ (3)°
 $V = 794.5$ (4) Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 0.34$ mm⁻¹
 $T = 293$ (2) K
 $0.20 \times 0.10 \times 0.02$ mm

Data collection

 Oxford Diffraction Xcalibur2 CCD diffractometer
 Absorption correction: analytical (*CrysAlis RED*; Oxford Diffraction; 2004; Clark & Reid, 1995)
 $T_{\min} = 0.929$, $T_{\max} = 0.967$

 7237 measured reflections
 962 independent reflections
 855 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\max} = 23.1^\circ$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.077$
 $S = 1.09$
 962 reflections

 101 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2}\cdots\text{S}^i$	0.86	2.57	3.408 (2)	166
$\text{C3}-\text{H3}\cdots\text{Cg}^{ii}$	0.93	2.74	3.464 (3)	136

 Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$. Cg is the centroid of the C2–C7 ring.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2004); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2004); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *PLATON*.

The authors acknowledge the Higher Education Commission, Pakistan, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2055).

References

- Baily, N., Dean, A. W., Judd, D. B., Middlemiss, D., Storer, R. & Watson, S. P. (1996). *Bioorg. Med. Chem. Lett.* **6**, 1409–1413.
 Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
 Koch, K. R. (2001). *Coord. Chem. Rev.* **216**, 473–482.
 Namgun, L., Mi-Hyun, C. & Tack, H. K. (2001). *J. Korean Chem. Soc.* **45**, 96–99.
 Oxford Diffraction (2004). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
 Patel, D. G. & Chedekel, M. R. (1984). *J. Org. Chem.* **49**, 997–1000.
 Schuster, M., Kugler, B. & König, K. H. (1990). *J. Anal. Chem.* **338**, 717–720.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o1141 [doi:10.1107/S1600536808015043]

1-Methyl-1*H*-benzimidazole-2(3*H*)-thione

H. Khan, A. Badshah, F. Shaheen, C. Gieck and R. A. Qureshi

Comment

N,N'-disubstituted and *N*-substituted thiourea derivatives are the major building blocks of organic macromolecular compounds. Thiourea derivatives such as benzothiazoles have been isolated by bromination of arylthioureas (Patil & Chedekel, 1984) and by condensation of 2-aminothiazole (Baily *et al.*, 1996), by cyclization of *N*-(2-hydroxyethyl)-*N*-methylthioureas and 2-methyl-aminothiazole (Namgun *et al.*, 2001). Aliphatic and acylthioureas have a wide range of application due to their coordination behavior towards transition metals (Schuster *et al.*, 1990). *N,N*-dialkyl-*N*-arylthioureas have been used for the extraction of metals such as nickel, palladium and platinum (Koch, 2001). Here we report the crystal structure of the title compound, 1-methyl-2*H*-benzimidazole-2-thione (Fig. 1).

The benzimidazole unit is essentially planar, with a mean deviation of 0.023 Å from the least-squares plane defined by the nine constituent atoms. The molecular packing (Fig. 2) is stabilized by a C—H \cdots π interaction between a benzene H atom and the benzene ring of neighbouring molecules, with a C3—H3 \cdots Cgⁱ separation of 2.735 (3) Å (Fig. 2 and Table 1; Cg is the C2-C7 benzene ring, symmetry code as in Fig. 2). Additionally, intermolecular N—H \cdots S interactions in the structure were observed (Fig. 2 and Table 1; symmetry code as in Fig. 2).

Experimental

Compound (I) was synthesized by the addition of carbondisulfide (0.79 ml, 13.02 mmol) to *N*-methyl-1,2-phenylenediamine (0.744 ml, 6.55 mmol) in methanol (20 ml). The resulting mixture was stirred for 24 h, at 0°C temperature, giving a clear light yellow solution. The solution was evaporated under reduced pressure to give a light yellow solid, which was recrystallized in methanol/petroleum ether (9:1) to afford compound (I) (yield : 76%).

Refinement

All H atoms were placed in idealized positions (C—H = 0.96 Å ° (methyl); C—H = 0.93 Å ° (aromatic); N—H = 0.86 Å °) and refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C, N})$ or $1.5U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The structure of (I), with displacement ellipsoids drawn at the 50% probability level.




Fig. 2. C—H... π and N—H...S interactions (dotted lines) in the title compound. Cg denotes the ring centroid. [Symmetry code: (i) $-x+1/2, y-1/2, -z+3/2$; (ii) $-x+3/2, y+1/2, -z+3/2$; (iii) $-x+3/2, y-1/2, -z+3/2$; (iv) $-x+1/2, y+1/2, -z+3/2$.]

1-Methyl-1H-benzimidazole-2(3H)-thione

Crystal data

$C_8H_8N_2S$	$F_{000} = 344$
$M_r = 164.22$	$D_x = 1.373 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 402 K
Hall symbol: $-P_2yn$	Mo $K\alpha$ radiation
$a = 9.997(4) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 5.8140(7) \text{ \AA}$	Cell parameters from 5701 reflections
$c = 13.703(4) \text{ \AA}$	$\theta = 3.7\text{--}23.1^\circ$
$\beta = 94.05(3)^\circ$	$\mu = 0.34 \text{ mm}^{-1}$
$V = 794.5(4) \text{ \AA}^3$	$T = 293(2) \text{ K}$
$Z = 4$	Block, colourless
	$0.20 \times 0.10 \times 0.02 \text{ mm}$

Data collection

Oxford Diffraction Xcalibur2 CCD diffractometer	962 independent reflections
Radiation source: Enhance (Mo) X-ray Source	855 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
Detector resolution: $10.0 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 23.1^\circ$
$T = 293(2) \text{ K}$	$\theta_{\text{min}} = 3.8^\circ$
ω scans	$h = -10 \rightarrow 11$
Absorption correction: analytical (CrysAlis RED; Oxford Diffraction; 2004; Clark & Reid, 1995)	$k = -6 \rightarrow 6$
$T_{\text{min}} = 0.929, T_{\text{max}} = 0.967$	$l = -14 \rightarrow 14$
7237 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.029$	H-atom parameters constrained
$wR(F^2) = 0.078$	$w = 1/[\sigma^2(F_o^2) + (0.0353P)^2 + 0.4983P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
962 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$

101 parameters

$$\Delta\rho_{\min} = -0.19 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S	0.73449 (6)	0.62833 (10)	0.62836 (4)	0.0475 (3)
N1	0.52959 (17)	0.3736 (3)	0.69273 (12)	0.0345 (5)
N2	0.57646 (17)	0.6789 (3)	0.78071 (12)	0.0380 (5)
H2	0.6119	0.8071	0.8004	0.046*
C1	0.6119 (2)	0.5599 (4)	0.70145 (15)	0.0359 (6)
C2	0.4462 (2)	0.3690 (3)	0.77058 (15)	0.0323 (5)
C3	0.3528 (2)	0.2095 (4)	0.79703 (16)	0.0399 (6)
H3	0.3342	0.0773	0.7603	0.048*
C4	0.2882 (2)	0.2566 (4)	0.88107 (17)	0.0461 (6)
H4	0.2255	0.1524	0.9017	0.055*
C5	0.3145 (2)	0.4548 (4)	0.93499 (17)	0.0466 (6)
H5	0.2684	0.4812	0.9906	0.056*
C6	0.4079 (2)	0.6150 (4)	0.90824 (16)	0.0412 (6)
H6	0.4246	0.7492	0.9440	0.049*
C7	0.4752 (2)	0.5656 (4)	0.82572 (15)	0.0330 (5)
C8	0.5331 (3)	0.1997 (4)	0.61682 (17)	0.0518 (7)
H8A	0.5508	0.2717	0.5560	0.078*
H8B	0.4482	0.1221	0.6098	0.078*
H8C	0.6026	0.0904	0.6345	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S	0.0446 (4)	0.0454 (4)	0.0547 (4)	0.0026 (3)	0.0184 (3)	0.0102 (3)
N1	0.0345 (10)	0.0313 (10)	0.0384 (10)	0.0019 (8)	0.0071 (8)	-0.0027 (8)
N2	0.0379 (11)	0.0323 (10)	0.0443 (11)	-0.0042 (8)	0.0064 (9)	-0.0040 (9)
C1	0.0350 (12)	0.0332 (12)	0.0396 (13)	0.0068 (11)	0.0026 (10)	0.0050 (10)
C2	0.0283 (11)	0.0339 (13)	0.0348 (12)	0.0062 (10)	0.0024 (9)	0.0023 (10)
C3	0.0365 (12)	0.0356 (13)	0.0478 (14)	-0.0024 (11)	0.0043 (11)	-0.0016 (11)

supplementary materials

C4	0.0380 (13)	0.0483 (16)	0.0529 (15)	-0.0053 (11)	0.0094 (12)	0.0080 (13)
C5	0.0425 (13)	0.0601 (16)	0.0383 (13)	0.0015 (13)	0.0104 (11)	0.0021 (12)
C6	0.0423 (13)	0.0431 (14)	0.0380 (13)	0.0040 (11)	0.0017 (10)	-0.0060 (11)
C7	0.0302 (12)	0.0330 (12)	0.0358 (12)	0.0021 (10)	0.0019 (10)	0.0020 (10)
C8	0.0536 (15)	0.0493 (15)	0.0542 (15)	-0.0015 (13)	0.0150 (12)	-0.0150 (13)

Geometric parameters (Å, °)

S—C1	1.684 (2)	C3—H3	0.9300
N1—C1	1.361 (3)	C4—C5	1.384 (3)
N1—C2	1.400 (3)	C4—H4	0.9300
N1—C8	1.453 (3)	C5—C6	1.386 (3)
N2—C1	1.356 (3)	C5—H5	0.9300
N2—C7	1.389 (3)	C6—C7	1.386 (3)
N2—H2	0.8600	C6—H6	0.9300
C2—C3	1.383 (3)	C8—H8A	0.9600
C2—C7	1.389 (3)	C8—H8B	0.9600
C3—C4	1.387 (3)	C8—H8C	0.9600
C1—N1—C2	109.71 (17)	C3—C4—H4	119.2
C1—N1—C8	124.88 (18)	C4—C5—C6	121.6 (2)
C2—N1—C8	125.34 (18)	C4—C5—H5	119.2
C1—N2—C7	110.71 (18)	C6—C5—H5	119.2
C1—N2—H2	124.6	C7—C6—C5	116.8 (2)
C7—N2—H2	124.6	C7—C6—H6	121.6
N2—C1—N1	106.62 (18)	C5—C6—H6	121.6
N2—C1—S	126.72 (18)	C6—C7—N2	132.5 (2)
N1—C1—S	126.65 (17)	C6—C7—C2	121.3 (2)
C3—C2—C7	121.86 (19)	N2—C7—C2	106.21 (17)
C3—C2—N1	131.46 (19)	N1—C8—H8A	109.5
C7—C2—N1	106.66 (17)	N1—C8—H8B	109.5
C2—C3—C4	116.6 (2)	H8A—C8—H8B	109.5
C2—C3—H3	121.7	N1—C8—H8C	109.5
C4—C3—H3	121.7	H8A—C8—H8C	109.5
C5—C4—C3	121.7 (2)	H8B—C8—H8C	109.5
C5—C4—H4	119.2		
C7—N2—C1—N1	2.4 (2)	C2—C3—C4—C5	0.9 (3)
C7—N2—C1—S	-177.28 (16)	C3—C4—C5—C6	-0.7 (4)
C2—N1—C1—N2	-3.2 (2)	C4—C5—C6—C7	-1.0 (3)
C8—N1—C1—N2	179.79 (19)	C5—C6—C7—N2	-177.1 (2)
C2—N1—C1—S	176.50 (15)	C5—C6—C7—C2	2.4 (3)
C8—N1—C1—S	-0.5 (3)	C1—N2—C7—C6	178.9 (2)
C1—N1—C2—C3	-175.6 (2)	C1—N2—C7—C2	-0.7 (2)
C8—N1—C2—C3	1.4 (3)	C3—C2—C7—C6	-2.3 (3)
C1—N1—C2—C7	2.8 (2)	N1—C2—C7—C6	179.11 (19)
C8—N1—C2—C7	179.79 (19)	C3—C2—C7—N2	177.32 (19)
C7—C2—C3—C4	0.6 (3)	N1—C2—C7—N2	-1.2 (2)
N1—C2—C3—C4	178.8 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2···S ⁱ	0.86	2.57	3.408 (2)	166
C3—H3···Cg ⁱⁱ	0.93	2.74	3.464 (3)	136

Symmetry codes: (i) $-x+3/2, y+1/2, -z+3/2$; (ii) $-x+1/2, y-1/2, -z+3/2$.

Fig. 1




Fig. 2

