

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

trans-5-(4-Chlorophenyl)-*N*-cyclohexyl-4-methyl-2-oxo-1,3-thiazolidine-3-carboxamide

Ying-Hui Yu, Ji-Long Liu, Li-Zhong Niu, Guang-Feng Hou and Jin-Sheng Gao*


 College of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China
 Correspondence e-mail: hg1000@163.com

Received 24 April 2008; accepted 30 April 2008

 Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.109; data-to-parameter ratio = 19.6.

The title pesticide, $\text{C}_{17}\text{H}_{21}\text{ClN}_2\text{O}_2\text{S}$, has a *trans* arrangement of the 4-chlorophenyl and 4-methyl substituents of the thiazolidine ring; the structure features an intramolecular amide-ring carbonyl $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond. The thiazolidine ring is almost planar, the largest deviation being 0.199 (1) Å for the methyl-substituted C atom, and the cyclohexane ring has a chair conformation.

Related literature

 For the synthesis of the pesticide Hexythiazox, see: Iwataki *et al.* (1981); Yamada *et al.* (1983).


Experimental

Crystal data

 $\text{C}_{17}\text{H}_{21}\text{ClN}_2\text{O}_2\text{S}$
 $M_r = 352.88$
 Monoclinic, $P2_1/c$
 $a = 10.284$ (4) Å

 $b = 11.799$ (5) Å
 $c = 15.902$ (5) Å
 $\beta = 111.830$ (14)°
 $V = 1791.2$ (12) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.34$ mm⁻¹
 $T = 291$ (2) K
 $0.27 \times 0.26 \times 0.25$ mm

Data collection

 Rigaku R-Axis RAPID diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.913$, $T_{\max} = 0.918$

 17194 measured reflections
 4088 independent reflections
 3066 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.109$
 $S = 1.06$
 4088 reflections

 209 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.27$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H10}\cdots\text{O1}$	0.84	2.03	2.706 (2)	137
$\text{C2}-\text{H1}\cdots\text{O2}^i$	0.93	2.47	3.386 (2)	170
$\text{C5}-\text{H3}\cdots\text{S1}$	0.93	2.79	3.168 (2)	105
$\text{C12}-\text{H11}\cdots\text{O2}$	0.98	2.44	2.831 (2)	103

 Symmetry code: (i) $-x + 1, -y + 2, -z + 2$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank Heilongjiang University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2450).

References

- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 Iwataki, I., Kaeriyama, M., Matsui, N. & Yamada, T. (1981). German Patent 3 037 105.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MS (2002). *CrystalStructure*. Rigaku/MS Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Yamada, T., Kaeriyama, M., Matsui, N. & Yoneda, H. (1983). Japan Patent 58 110 577.

supplementary materials

Acta Cryst. (2008). E64, o1019 [doi:10.1107/S1600536808012634]

***trans*-5-(4-Chlorophenyl)-*N*-cyclohexyl-4-methyl-2-oxo-1,3-thiazolidine-3-carboxamide**

Y.-H. Yu, J.-L. Liu, L.-Z. Niu, G.-F. Hou and J.-S. Gao

Comment

Hexythiazox, chemically named *trans*-5-(4-chlorophenyl)- *N*-cyclohexyl-4-methyl-2-oxo-3-thiazolidinecarboxamide, is known as a high efficiency of pesticide. In this paper, we first report the crystal structure of hexythiazox (I).

The title compound (I), consists of a planar phenyl ring (A), a S-contained five-numbers ring (B) and a cyclohexane ring (C). The S-contained five-numbers ring is almost coplanar, with the largest deviation being 0.199 (1) Å for atom C8, and the cyclohexane ring is chair forms. The three rings make the following dihedral angles: A/B 82.20 (0.06)°, A/C 54.22 (0.07)° and B/C 81.70 (0.06)°.

In the crystal structure, an extensive network of intramolecular N—H···O and intermolecular C—H···O hydrogen bonds stabilizes the packing (Table 1; Fig. 2).

Experimental

Hexythiazox was synthesized by the reaction of 5-(4-chlorophenyl)-4-methylthiazolidin-2-one and isocyanatocyclohexane in toluene solution in the patent literature. Crystals suitable for X-ray experiments were obtained by slow evaporation of an ethanol solution.

Refinement

H atoms bound to C atoms were placed in calculated positions and treated as riding on their parent atoms, with C—H = 0.93 Å (aromatic), C—H = 0.98 Å (methine), C—H = 0.97 Å (methylene), C—H = 0.96 Å (methyl) and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. N-H atoms were initially located in a difference Fourier map but they were treated as riding on their parent atoms with N—H = 0.85 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of (I), showing displacement ellipsoids at the 30% probability level for non-H atoms.




Fig. 2. A partial packing view, showing the three-dimensional hydrogen-bonding network. Dashed lines indicate the hydrogen-bonding interactions. H atoms not involved in hydrogen bonds have been omitted for clarity.

***trans*-5-(4-Chlorophenyl)-*N*-cyclohexyl-4-methyl-2-oxo-1,3-thiazolidine-3-carboxamide**

Crystal data

$C_{17}H_{21}ClN_2O_2S$

$M_r = 352.88$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.284$ (4) Å

$b = 11.799$ (5) Å

$c = 15.902$ (5) Å

$\beta = 111.830$ (14)°

$V = 1791.2$ (12) Å³

$Z = 4$

$F_{000} = 744$

$D_x = 1.309$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 12810 reflections

$\theta = 3.3$ – 27.5 °

$\mu = 0.34$ mm⁻¹

$T = 291$ (2) K

Block, colorless

$0.27 \times 0.26 \times 0.25$ mm

Data collection

Rigaku R-Axis RAPID
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 291$ (2) K

ω scans

Absorption correction: multi-scan
(ABSCOR; Higashi, 1995)

$T_{\min} = 0.913$, $T_{\max} = 0.918$

17194 measured reflections

4088 independent reflections

3066 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$

$\theta_{\max} = 27.5$ °

$\theta_{\min} = 3.3$ °

$h = -13 \rightarrow 11$

$k = -15 \rightarrow 15$

$l = -19 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.109$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0513P)^2 + 0.331P]$

$S = 1.06$

4088 reflections

209 parameters

Primary atom site location: structure-invariant direct methods

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5475 (2)	1.00752 (16)	0.78248 (11)	0.0503 (4)
C2	0.5027 (2)	1.07700 (16)	0.83577 (12)	0.0564 (5)
H1	0.5544	1.1405	0.8637	0.068*
C3	0.3794 (2)	1.05051 (15)	0.84691 (12)	0.0540 (5)
H2	0.3480	1.0976	0.8823	0.065*
C4	0.30034 (18)	0.95543 (13)	0.80670 (10)	0.0426 (4)
C5	0.3492 (2)	0.88779 (15)	0.75323 (11)	0.0509 (4)
H3	0.2985	0.8239	0.7253	0.061*
C6	0.4717 (2)	0.91368 (16)	0.74085 (12)	0.0547 (5)
H4	0.5027	0.8679	0.7045	0.066*
C7	0.17090 (18)	0.93163 (14)	0.82722 (12)	0.0460 (4)
H5	0.1281	1.0046	0.8313	0.055*
C8	0.20030 (17)	0.86779 (12)	0.91720 (11)	0.0392 (4)
H6	0.2982	0.8802	0.9567	0.047*
C9	0.1069 (2)	0.90648 (17)	0.96633 (16)	0.0638 (5)
H7	0.0106	0.9016	0.9262	0.096*
H8	0.1289	0.9835	0.9859	0.096*
H9	0.1220	0.8588	1.0181	0.096*
C10	0.08671 (16)	0.72176 (14)	0.80822 (11)	0.0418 (4)
C11	0.24351 (16)	0.66650 (13)	0.96508 (10)	0.0382 (3)
C12	0.27713 (17)	0.46582 (13)	1.00756 (11)	0.0426 (4)
H11	0.3500	0.4975	1.0615	0.051*
C13	0.34475 (19)	0.37870 (16)	0.96687 (12)	0.0506 (4)
H12	0.4216	0.4134	0.9548	0.061*
H13	0.2767	0.3519	0.9098	0.061*
C14	0.3994 (2)	0.27909 (16)	1.03110 (15)	0.0640 (6)

supplementary materials

H14	0.4387	0.2227	1.0029	0.077*
H15	0.4734	0.3049	1.0861	0.077*
C15	0.2844 (3)	0.22642 (17)	1.05453 (16)	0.0745 (7)
H16	0.3218	0.1637	1.0959	0.089*
H17	0.2127	0.1969	1.0000	0.089*
C16	0.2203 (2)	0.3122 (2)	1.09784 (15)	0.0721 (6)
H18	0.2902	0.3371	1.1549	0.087*
H19	0.1441	0.2772	1.1105	0.087*
C17	0.1650 (2)	0.41410 (18)	1.03619 (13)	0.0561 (5)
H20	0.0863	0.3908	0.9828	0.067*
H21	0.1318	0.4707	1.0677	0.067*
C11	0.70303 (6)	1.03996 (6)	0.76763 (4)	0.07627 (19)
N1	0.18010 (13)	0.74624 (10)	0.89396 (8)	0.0369 (3)
N2	0.21922 (17)	0.55814 (12)	0.94286 (10)	0.0558 (4)
H10	0.1648	0.5422	0.8896	0.067*
O1	0.04055 (13)	0.62916 (10)	0.77874 (8)	0.0545 (3)
O2	0.31689 (14)	0.70170 (10)	1.03933 (7)	0.0520 (3)
S1	0.03912 (5)	0.84657 (4)	0.74263 (3)	0.06226 (18)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0621 (11)	0.0475 (10)	0.0381 (8)	-0.0070 (8)	0.0149 (8)	0.0067 (7)
C2	0.0721 (12)	0.0442 (10)	0.0500 (10)	-0.0206 (9)	0.0192 (9)	-0.0081 (8)
C3	0.0723 (12)	0.0393 (9)	0.0511 (10)	-0.0112 (8)	0.0238 (9)	-0.0112 (8)
C4	0.0533 (9)	0.0299 (8)	0.0347 (7)	-0.0024 (7)	0.0048 (7)	0.0050 (6)
C5	0.0650 (11)	0.0354 (9)	0.0431 (9)	-0.0083 (8)	0.0092 (8)	-0.0059 (7)
C6	0.0706 (12)	0.0479 (11)	0.0434 (9)	-0.0012 (9)	0.0187 (9)	-0.0042 (8)
C7	0.0457 (9)	0.0291 (8)	0.0518 (9)	0.0032 (7)	0.0047 (7)	0.0046 (7)
C8	0.0427 (8)	0.0263 (7)	0.0462 (8)	-0.0002 (6)	0.0137 (7)	-0.0025 (6)
C9	0.0720 (13)	0.0448 (11)	0.0865 (15)	0.0026 (9)	0.0431 (12)	-0.0134 (10)
C10	0.0359 (8)	0.0384 (9)	0.0428 (8)	0.0002 (6)	0.0049 (7)	0.0003 (7)
C11	0.0430 (8)	0.0312 (8)	0.0389 (8)	-0.0008 (6)	0.0137 (7)	-0.0006 (6)
C12	0.0500 (9)	0.0285 (8)	0.0398 (8)	0.0008 (6)	0.0058 (7)	0.0011 (6)
C13	0.0475 (9)	0.0485 (10)	0.0537 (10)	0.0018 (8)	0.0164 (8)	-0.0029 (8)
C14	0.0599 (12)	0.0448 (11)	0.0698 (12)	0.0186 (9)	0.0039 (10)	-0.0057 (9)
C15	0.0853 (15)	0.0351 (10)	0.0751 (13)	-0.0044 (10)	-0.0026 (12)	0.0133 (9)
C16	0.0732 (13)	0.0748 (15)	0.0641 (12)	-0.0126 (12)	0.0206 (11)	0.0223 (11)
C17	0.0517 (10)	0.0593 (12)	0.0569 (10)	0.0086 (9)	0.0197 (9)	0.0045 (9)
C11	0.0787 (4)	0.0850 (4)	0.0734 (3)	-0.0163 (3)	0.0380 (3)	0.0019 (3)
N1	0.0414 (7)	0.0269 (6)	0.0374 (6)	0.0017 (5)	0.0087 (5)	-0.0016 (5)
N2	0.0764 (10)	0.0290 (7)	0.0418 (7)	-0.0011 (7)	-0.0013 (7)	0.0002 (6)
O1	0.0542 (7)	0.0403 (7)	0.0520 (7)	-0.0077 (5)	0.0001 (6)	-0.0066 (5)
O2	0.0719 (8)	0.0352 (6)	0.0370 (6)	-0.0045 (6)	0.0066 (6)	-0.0013 (5)
S1	0.0508 (3)	0.0494 (3)	0.0583 (3)	-0.0045 (2)	-0.0125 (2)	0.0140 (2)

Geometric parameters (\AA , $^\circ$)

C1—C6	1.374 (3)	C10—S1	1.7655 (18)
-------	-----------	--------	-------------

C1—C2	1.376 (3)	C11—O2	1.2139 (19)
C1—C11	1.744 (2)	C11—N2	1.325 (2)
C2—C3	1.380 (3)	C11—N1	1.4286 (19)
C2—H1	0.9300	C12—N2	1.464 (2)
C3—C4	1.393 (2)	C12—C13	1.514 (2)
C3—H2	0.9300	C12—C17	1.517 (3)
C4—C5	1.389 (3)	C12—H11	0.9800
C4—C7	1.510 (3)	C13—C14	1.521 (3)
C5—C6	1.379 (3)	C13—H12	0.9700
C5—H3	0.9300	C13—H13	0.9700
C6—H4	0.9300	C14—C15	1.501 (3)
C7—C8	1.544 (2)	C14—H14	0.9700
C7—S1	1.8159 (17)	C14—H15	0.9700
C7—H5	0.9800	C15—C16	1.507 (4)
C8—N1	1.4763 (19)	C15—H16	0.9700
C8—C9	1.517 (3)	C15—H17	0.9700
C8—H6	0.9800	C16—C17	1.522 (3)
C9—H7	0.9600	C16—H18	0.9700
C9—H8	0.9600	C16—H19	0.9700
C9—H9	0.9600	C17—H20	0.9700
C10—O1	1.214 (2)	C17—H21	0.9700
C10—N1	1.375 (2)	N2—H10	0.8445
C6—C1—C2	121.10 (19)	N2—C12—C17	110.75 (15)
C6—C1—C11	119.77 (16)	C13—C12—C17	112.11 (15)
C2—C1—C11	119.14 (15)	N2—C12—H11	108.1
C1—C2—C3	118.59 (17)	C13—C12—H11	108.1
C1—C2—H1	120.7	C17—C12—H11	108.1
C3—C2—H1	120.7	C12—C13—C14	110.57 (16)
C2—C3—C4	122.04 (18)	C12—C13—H12	109.5
C2—C3—H2	119.0	C14—C13—H12	109.5
C4—C3—H2	119.0	C12—C13—H13	109.5
C5—C4—C3	117.47 (18)	C14—C13—H13	109.5
C5—C4—C7	125.03 (15)	H12—C13—H13	108.1
C3—C4—C7	117.46 (16)	C15—C14—C13	110.99 (16)
C6—C5—C4	121.16 (16)	C15—C14—H14	109.4
C6—C5—H3	119.4	C13—C14—H14	109.4
C4—C5—H3	119.4	C15—C14—H15	109.4
C1—C6—C5	119.64 (18)	C13—C14—H15	109.4
C1—C6—H4	120.2	H14—C14—H15	108.0
C5—C6—H4	120.2	C14—C15—C16	110.77 (17)
C4—C7—C8	113.97 (13)	C14—C15—H16	109.5
C4—C7—S1	114.69 (13)	C16—C15—H16	109.5
C8—C7—S1	104.59 (11)	C14—C15—H17	109.5
C4—C7—H5	107.8	C16—C15—H17	109.5
C8—C7—H5	107.8	H16—C15—H17	108.1
S1—C7—H5	107.8	C15—C16—C17	111.00 (18)
N1—C8—C9	111.37 (14)	C15—C16—H18	109.4
N1—C8—C7	106.33 (13)	C17—C16—H18	109.4
C9—C8—C7	112.78 (15)	C15—C16—H19	109.4

supplementary materials

N1—C8—H6	108.8	C17—C16—H19	109.4
C9—C8—H6	108.8	H18—C16—H19	108.0
C7—C8—H6	108.8	C12—C17—C16	111.58 (16)
C8—C9—H7	109.5	C12—C17—H20	109.3
C8—C9—H8	109.5	C16—C17—H20	109.3
H7—C9—H8	109.5	C12—C17—H21	109.3
C8—C9—H9	109.5	C16—C17—H21	109.3
H7—C9—H9	109.5	H20—C17—H21	108.0
H8—C9—H9	109.5	C10—N1—C11	126.26 (13)
O1—C10—N1	126.92 (15)	C10—N1—C8	115.70 (12)
O1—C10—S1	122.68 (12)	C11—N1—C8	117.47 (12)
N1—C10—S1	110.40 (11)	C11—N2—C12	122.90 (14)
O2—C11—N2	125.23 (15)	C11—N2—H10	118.1
O2—C11—N1	118.71 (14)	C12—N2—H10	118.9
N2—C11—N1	116.04 (13)	C10—S1—C7	93.26 (8)
N2—C12—C13	109.64 (15)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H10 \cdots O1	0.84	2.03	2.706 (2)	137
C2—H1 \cdots O2 ⁱ	0.93	2.47	3.386 (2)	170
C5—H3 \cdots S1	0.93	2.79	3.168 (2)	105
C12—H11 \cdots O2	0.98	2.44	2.831 (2)	103

Symmetry codes: (i) $-x+1, -y+2, -z+2$.

Fig. 1




Fig. 2

