

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(4-Iodophenyl)-5-methyl-3-methylsulfinyl-1-benzofuran

 Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}

^aDepartment of Chemistry, Donggeui University, San 24 Kaya-dong, Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea

Correspondence e-mail: uklee@pknu.ac.kr


Received 6 May 2008; accepted 8 May 2008

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.027; wR factor = 0.066; data-to-parameter ratio = 17.6.

The title compound, $\text{C}_{16}\text{H}_{13}\text{IO}_2\text{S}$, was prepared by the oxidation of 2-(4-iodophenyl)-5-methyl-3-methylsulfonyl-1-benzofuran with 3-chloroperoxybenzoic acid. The 4-iodophenyl ring makes a dihedral angle of 37.97 (9)° with the plane of the benzofuran fragment, and the O atom and the methyl group of the methylsulfinyl substituent lie on opposite sides of this plane. The molecular packing is stabilized by $\text{C}-\text{H}\cdots\pi$ interactions between H atoms on the 4-iodophenyl ring and the benzofuran rings, and by an $\text{I}\cdots\text{O}$ halogen bond of 3.252 (2) Å with a nearly linear $\text{C}-\text{I}\cdots\text{O}$ angle of 163.06 (8)°. In addition, the stacked molecules exhibit inversion-related $\text{S}\cdots\text{O}$ contacts [3.209 (2) Å] involving the sulfinyl groups.

Related literature

For the crystal structures of similar 2-aryl-5-methyl-3-methylsulfinyl-1-benzofuran compounds, see: Choi *et al.* (2007*a,b*). For a review of halogen bonding, see: Politzer *et al.* (2007). For details of sulfinyl–sulfinyl interactions, see: Choi *et al.* (2007*c*). For a review of carbonyl–carbonyl interactions, see: Allen *et al.* (1998).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{13}\text{IO}_2\text{S}$
 $M_r = 396.22$
 Monoclinic, $P2_1/n$
 $a = 9.258$ (2) Å
 $b = 15.939$ (3) Å
 $c = 10.299$ (2) Å
 $\beta = 103.471$ (3)°
 $V = 1477.9$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 2.31$ mm⁻¹
 $T = 173$ (2) K
 $0.40 \times 0.30 \times 0.30$ mm

Data collection

Bruker SMART CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2000)
 $T_{\min} = 0.443$, $T_{\max} = 0.508$
 8743 measured reflections
 3227 independent reflections
 2934 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.065$
 $S = 1.15$
 3227 reflections
 183 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.54$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.94$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---|-------|-------------|-------------|---------------|
| $\text{C10}-\text{H10}\cdots\text{Cg1}^1$ | 0.95 | 3.01 | 3.617 (4) | 125 |
| $\text{C11}-\text{H11}\cdots\text{Cg2}^1$ | 0.95 | 2.77 | 3.643 (4) | 148 |

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$. Cg1 and Cg2 are the centroids of the C2–C7 benzene ring and the O1/C8/C1/C2/C7 furan ring, respectively.

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 1998); software used to prepare material for publication: SHELXL97.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2496).

References

- Allen, F. H., Baalham, C. A., Lommerse, J. P. M. & Raithby, P. R. (1998). *Acta Cryst.* **B54**, 320–329.
 Brandenburg, K. (1998). *DIAMOND*, Crystal Impact GbR, Bonn, Germany.
 Bruker (2001). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2007*a*). *Acta Cryst.* **E63**, o3295.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2007*b*). *Acta Cryst.* **E63**, o4282.
 Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2007*c*). *Acta Cryst.* **E63**, o4811.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Politzer, P., Lane, P., Concha, M. C., Ma, Y. & Murray, J. S. (2007). *J. Mol. Model.* **13**, 305–311.
 Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o1061 [doi:10.1107/S1600536808013706]

2-(4-Iodophenyl)-5-methyl-3-methylsulfinyl-1-benzofuran

H. D. Choi, P. J. Seo, B. W. Son and U. Lee

Comment

This work is related to our preceding communications on the synthesis and structure of 2-aryl-5-methyl-3-methylsulfinyl-1-benzofuran derivatives, *viz.* 2-(4-bromophenyl)-5-methyl-3-methylsulfinyl-1-benzofuran (Choi *et al.*, 2007*a*) and 2-(4-bromophenyl)-5,7-dimethyl-3-methylsulfinyl-1-benzofuran (Choi *et al.*, 2007*b*). Here we report the crystal structure of the title compound, 2-(4-iodophenyl)-5-methyl-3-methylsulfinyl-1-benzofuran (Fig. 1).

The benzofuran unit is essentially planar, with a mean deviation of 0.008 Å from the least-squares plane defined by the nine constituent atoms. The 4-iodophenyl ring (C9—C14) makes a dihedral angle of 37.97 (9)° with the plane of the benzofuran fragment. The molecular packing (Fig. 2) is stabilized by two different C—H \cdots π interactions within each stack of molecules; one between a 4-iodophenyl H atom and the benzene ring (Cg1ⁱ), with a C10—H10 \cdots Cg1ⁱ separation of 3.617 (4) Å, and a second between a 4-iodophenyl H atom and the furan ring (Cg2ⁱ), with a C11—H11 \cdots Cg2ⁱ separation of 3.643 (4) Å, (Fig. 2 and Table 1; Cg1 and Cg2 are the centroids of the C2—C7 benzene ring and the O1/C8/C1/C2/C7 furan ring, respectively, symmetry code as in Fig. 2). The molecular packing is further stabilized by an I \cdots O halogen bond (Politzer *et al.*, 2007) between the iodine atom and the oxygen of a neighbouring S=O unit, with a I \cdots O2ⁱⁱ distance of 3.252 (2) Å (symmetry code as Fig. 2). In addition, the crystal packing exhibits a sulfinyl-sulfinyl interaction (Choi *et al.*, 2007*c*) interpreted as similar to a type-II carbonyl-carbonyl interaction (Allen *et al.*, 1998), with S \cdots O2ⁱⁱⁱ and O2 \cdots Sⁱⁱⁱ distance of 3.209 (2) Å (symmetry code as in Fig. 2)

Experimental

3-Chloroperoxybenzoic acid (77%, 247 mg, 1.1 mmol) was added in small portions to a stirred solution of 2-(4-iodophenyl)-5-methyl-3-methylsulanyl-1-benzofuran (380 mg, 1.0 mmol) in dichloromethane (30 ml) at 273 K. After stirring at room temperature for 2 h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated in vacuum. The residue was purified by column chromatography (ethyl acetate) to afford the title compound as a colorless solid [yield 84%, m.p. 472–473 K; R_f = 0.61 (ethyl acetate)]. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of title compound in tetrahydrofuran at room temperature. Spectroscopic analysis: ¹H NMR (CDCl₃, 400 MHz) δ 2.48 (s, 3H), 3.11 (s, 3H), 7.22 (d, J = 8.04 Hz, 1H), 7.42 (d, J = 8.44 Hz, 1H), 7.75 (d, J = 6.96 Hz, 2H), 7.85 (d, J = 6.96 Hz, 2H), 7.99 (s, 1H); EI—MS 396 [M^+].

Refinement

All H atoms were geometrically located in ideal positions and refined using a riding model, with C—H = 0.95 Å for aromatic H atoms and 0.98 Å for methyl H atoms, and with $U_{iso}(H) = 1.2U_{eq}(C)$ for aromatic H atoms, and 1.5 $U_{eq}(C)$ for methyl H atoms.

Figures




Fig. 1. The molecular structure of the title compound, showing displacement ellipsoids drawn at the 50% probability level.




Fig. 2. C—H... π , I...O halogen bond and S...O interactions (dotted lines) in the title compound. Cg denotes the ring centroids. [Symmetry code: (i) $x - 1/2, -y + 3/2, z - 1/2$; (ii) $x + 1/2, y, z - 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x + 1, y, z + 1$; (v) $x + 1/2, -y + 3/2, z + 1/2$.]

2-(4-Iodophenyl)-5-methyl-3-methylsulfinyl-1-benzofuran

Crystal data

$C_{16}H_{13}IO_2S$

$M_r = 396.22$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 9.258 (2) \text{ \AA}$

$b = 15.939 (3) \text{ \AA}$

$c = 10.299 (2) \text{ \AA}$

$\beta = 103.471 (3)^\circ$

$V = 1477.9 (5) \text{ \AA}^3$

$Z = 4$

$F_{000} = 776$

$D_x = 1.781 \text{ Mg m}^{-3}$

Melting point = 472–473 K

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6653 reflections

$\theta = 2.4\text{--}28.2^\circ$

$\mu = 2.31 \text{ mm}^{-1}$

$T = 173 (2) \text{ K}$

Block, colorless

$0.40 \times 0.30 \times 0.30 \text{ mm}$

Data collection

Bruker SMART CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 10.0 pixels mm^{-1}

$T = 173(2) \text{ K}$

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 2000)

$T_{\min} = 0.443, T_{\max} = 0.508$

8743 measured reflections

3227 independent reflections

2934 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 27.0^\circ$

$\theta_{\min} = 2.4^\circ$

$h = -11 \rightarrow 6$

$k = -20 \rightarrow 20$

$l = -11 \rightarrow 13$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.026$$

$$wR(F^2) = 0.065$$

$$S = 1.15$$

3227 reflections

183 parameters

Primary atom site location: structure-invariant direct methods

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0226P)^2 + 1.1637P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.54 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.94 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|----------------|---------------|----------------|----------------------------------|
| I | -0.134684 (19) | 0.613119 (11) | -0.281817 (17) | 0.03372 (7) |
| S | 0.51165 (7) | 0.54138 (4) | 0.32703 (6) | 0.02482 (13) |
| O1 | 0.61196 (19) | 0.66982 (11) | 0.02820 (17) | 0.0251 (4) |
| O2 | 0.5977 (2) | 0.57384 (14) | 0.45895 (18) | 0.0389 (5) |
| C1 | 0.5913 (3) | 0.59181 (15) | 0.2070 (2) | 0.0220 (5) |
| C2 | 0.7456 (3) | 0.61452 (15) | 0.2246 (2) | 0.0231 (5) |
| C3 | 0.8760 (3) | 0.60093 (15) | 0.3218 (3) | 0.0262 (5) |
| H3 | 0.8746 | 0.5687 | 0.3991 | 0.031* |
| C4 | 1.0087 (3) | 0.63528 (17) | 0.3040 (3) | 0.0309 (6) |
| C5 | 1.0091 (3) | 0.68158 (17) | 0.1884 (3) | 0.0329 (6) |
| H5 | 1.1006 | 0.7038 | 0.1771 | 0.039* |
| C6 | 0.8810 (3) | 0.69632 (16) | 0.0895 (3) | 0.0308 (6) |
| H6 | 0.8824 | 0.7277 | 0.0114 | 0.037* |
| C7 | 0.7515 (3) | 0.66243 (15) | 0.1120 (2) | 0.0249 (5) |
| C8 | 0.5155 (3) | 0.62626 (14) | 0.0892 (2) | 0.0228 (5) |
| C9 | 0.3610 (3) | 0.62508 (15) | 0.0136 (2) | 0.0217 (5) |
| C10 | 0.3001 (3) | 0.69536 (16) | -0.0607 (3) | 0.0270 (5) |
| H10 | 0.3572 | 0.7453 | -0.0564 | 0.032* |
| C11 | 0.1575 (3) | 0.69273 (16) | -0.1403 (3) | 0.0284 (5) |
| H11 | 0.1162 | 0.7410 | -0.1893 | 0.034* |
| C12 | 0.0746 (3) | 0.61946 (15) | -0.1483 (2) | 0.0235 (5) |
| C13 | 0.1323 (3) | 0.54923 (15) | -0.0743 (2) | 0.0230 (5) |

supplementary materials

| | | | | |
|------|------------|--------------|------------|------------|
| H13 | 0.0746 | 0.4995 | -0.0797 | 0.028* |
| C14 | 0.2744 (3) | 0.55201 (15) | 0.0075 (2) | 0.0218 (5) |
| H14 | 0.3134 | 0.5044 | 0.0595 | 0.026* |
| C15 | 1.1498 (3) | 0.62338 (19) | 0.4100 (3) | 0.0417 (7) |
| H15A | 1.1345 | 0.6425 | 0.4962 | 0.063* |
| H15B | 1.2298 | 0.6561 | 0.3867 | 0.063* |
| H15C | 1.1769 | 0.5638 | 0.4160 | 0.063* |
| C16 | 0.5769 (3) | 0.43689 (17) | 0.3073 (3) | 0.0317 (6) |
| H16A | 0.6833 | 0.4386 | 0.3096 | 0.048* |
| H16B | 0.5227 | 0.4139 | 0.2216 | 0.048* |
| H16C | 0.5604 | 0.4013 | 0.3801 | 0.048* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|--------------|--------------|--------------|--------------|--------------|--------------|
| I | 0.02443 (11) | 0.04004 (12) | 0.03256 (11) | -0.00188 (7) | -0.00175 (7) | 0.00229 (7) |
| S | 0.0250 (3) | 0.0303 (3) | 0.0206 (3) | -0.0021 (2) | 0.0083 (2) | -0.0031 (2) |
| O1 | 0.0205 (8) | 0.0272 (9) | 0.0277 (9) | -0.0025 (7) | 0.0057 (7) | 0.0030 (7) |
| O2 | 0.0418 (12) | 0.0538 (13) | 0.0214 (9) | -0.0143 (10) | 0.0082 (8) | -0.0077 (9) |
| C1 | 0.0224 (12) | 0.0229 (11) | 0.0215 (11) | -0.0001 (9) | 0.0066 (9) | -0.0038 (9) |
| C2 | 0.0239 (13) | 0.0225 (11) | 0.0233 (11) | 0.0009 (9) | 0.0062 (10) | -0.0069 (9) |
| C3 | 0.0251 (13) | 0.0251 (12) | 0.0274 (12) | 0.0031 (9) | 0.0039 (10) | -0.0077 (9) |
| C4 | 0.0231 (13) | 0.0290 (13) | 0.0382 (14) | 0.0037 (10) | 0.0024 (11) | -0.0142 (11) |
| C5 | 0.0209 (13) | 0.0293 (13) | 0.0497 (16) | -0.0028 (10) | 0.0108 (11) | -0.0095 (12) |
| C6 | 0.0257 (13) | 0.0281 (13) | 0.0413 (15) | -0.0005 (10) | 0.0133 (11) | -0.0007 (11) |
| C7 | 0.0212 (12) | 0.0227 (12) | 0.0313 (12) | -0.0002 (9) | 0.0067 (10) | -0.0042 (10) |
| C8 | 0.0217 (12) | 0.0216 (11) | 0.0259 (12) | -0.0013 (9) | 0.0076 (10) | -0.0025 (9) |
| C9 | 0.0208 (12) | 0.0245 (12) | 0.0203 (11) | -0.0006 (9) | 0.0055 (9) | -0.0019 (9) |
| C10 | 0.0252 (13) | 0.0236 (12) | 0.0319 (13) | -0.0040 (10) | 0.0062 (10) | 0.0030 (10) |
| C11 | 0.0286 (13) | 0.0250 (12) | 0.0308 (13) | 0.0012 (10) | 0.0051 (11) | 0.0068 (10) |
| C12 | 0.0190 (11) | 0.0295 (13) | 0.0221 (11) | 0.0002 (9) | 0.0046 (9) | -0.0014 (9) |
| C13 | 0.0252 (12) | 0.0211 (11) | 0.0245 (11) | -0.0034 (9) | 0.0095 (10) | -0.0024 (9) |
| C14 | 0.0247 (12) | 0.0209 (11) | 0.0202 (11) | 0.0007 (9) | 0.0062 (9) | 0.0014 (9) |
| C15 | 0.0240 (14) | 0.0445 (17) | 0.0514 (18) | 0.0020 (12) | -0.0017 (13) | -0.0162 (14) |
| C16 | 0.0351 (15) | 0.0325 (14) | 0.0296 (13) | 0.0038 (11) | 0.0116 (11) | 0.0046 (11) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|--------------------|-------------|---------|-----------|
| I—O2 ⁱ | 3.252 (2) | C6—H6 | 0.9500 |
| I—C12 | 2.101 (3) | C8—C9 | 1.461 (3) |
| S—O2 | 1.4981 (19) | C9—C10 | 1.399 (3) |
| S—O2 ⁱⁱ | 3.209 (2) | C9—C14 | 1.407 (3) |
| S—C1 | 1.773 (2) | C10—C11 | 1.383 (4) |
| S—C16 | 1.799 (3) | C10—H10 | 0.9500 |
| O1—C7 | 1.381 (3) | C11—C12 | 1.389 (3) |
| O1—C8 | 1.391 (3) | C11—H11 | 0.9500 |
| C1—C8 | 1.367 (3) | C12—C13 | 1.389 (3) |
| C1—C2 | 1.444 (3) | C13—C14 | 1.388 (3) |

| | | | |
|-----------------------|--------------|---------------|--------------|
| C2—C3 | 1.394 (4) | C13—H13 | 0.9500 |
| C2—C7 | 1.400 (3) | C14—H14 | 0.9500 |
| C3—C4 | 1.395 (4) | C15—H15A | 0.9800 |
| C3—H3 | 0.9500 | C15—H15B | 0.9800 |
| C4—C5 | 1.402 (4) | C15—H15C | 0.9800 |
| C4—C15 | 1.507 (4) | C16—H16A | 0.9800 |
| C5—C6 | 1.391 (4) | C16—H16B | 0.9800 |
| C5—H5 | 0.9500 | C16—H16C | 0.9800 |
| C6—C7 | 1.383 (4) | | |
| C12—I—O2 ⁱ | 163.06 (8) | C10—C9—C8 | 120.1 (2) |
| O2—S—C1 | 104.84 (12) | C14—C9—C8 | 120.7 (2) |
| O2—S—C16 | 107.47 (13) | C11—C10—C9 | 120.5 (2) |
| C1—S—C16 | 97.80 (12) | C11—C10—H10 | 119.8 |
| C7—O1—C8 | 106.30 (18) | C9—C10—H10 | 119.8 |
| C8—C1—C2 | 107.4 (2) | C10—C11—C12 | 119.9 (2) |
| C8—C1—S | 126.1 (2) | C10—C11—H11 | 120.1 |
| C2—C1—S | 125.89 (18) | C12—C11—H11 | 120.1 |
| C3—C2—C7 | 119.1 (2) | C13—C12—C11 | 120.6 (2) |
| C3—C2—C1 | 135.8 (2) | C13—C12—I | 119.88 (18) |
| C7—C2—C1 | 105.1 (2) | C11—C12—I | 119.44 (18) |
| C2—C3—C4 | 119.1 (2) | C14—C13—C12 | 119.7 (2) |
| C2—C3—H3 | 120.5 | C14—C13—H13 | 120.1 |
| C4—C3—H3 | 120.5 | C12—C13—H13 | 120.1 |
| C3—C4—C5 | 119.7 (2) | C13—C14—C9 | 120.2 (2) |
| C3—C4—C15 | 119.8 (3) | C13—C14—H14 | 119.9 |
| C5—C4—C15 | 120.6 (3) | C9—C14—H14 | 119.9 |
| C6—C5—C4 | 122.7 (2) | C4—C15—H15A | 109.5 |
| C6—C5—H5 | 118.7 | C4—C15—H15B | 109.5 |
| C4—C5—H5 | 118.7 | H15A—C15—H15B | 109.5 |
| C7—C6—C5 | 115.9 (3) | C4—C15—H15C | 109.5 |
| C7—C6—H6 | 122.1 | H15A—C15—H15C | 109.5 |
| C5—C6—H6 | 122.1 | H15B—C15—H15C | 109.5 |
| O1—C7—C6 | 125.7 (2) | S—C16—H16A | 109.5 |
| O1—C7—C2 | 110.7 (2) | S—C16—H16B | 109.5 |
| C6—C7—C2 | 123.6 (2) | H16A—C16—H16B | 109.5 |
| C1—C8—O1 | 110.5 (2) | S—C16—H16C | 109.5 |
| C1—C8—C9 | 134.8 (2) | H16A—C16—H16C | 109.5 |
| O1—C8—C9 | 114.6 (2) | H16B—C16—H16C | 109.5 |
| C10—C9—C14 | 119.1 (2) | | |
| O2—S—C1—C8 | -136.9 (2) | C1—C2—C7—C6 | -179.6 (2) |
| C16—S—C1—C8 | 112.7 (2) | C2—C1—C8—O1 | 0.4 (3) |
| O2—S—C1—C2 | 33.6 (2) | S—C1—C8—O1 | 172.33 (17) |
| C16—S—C1—C2 | -76.9 (2) | C2—C1—C8—C9 | 178.0 (3) |
| C8—C1—C2—C3 | 178.7 (3) | S—C1—C8—C9 | -10.1 (4) |
| S—C1—C2—C3 | 6.8 (4) | C7—O1—C8—C1 | -0.5 (3) |
| C8—C1—C2—C7 | -0.2 (3) | C7—O1—C8—C9 | -178.63 (19) |
| S—C1—C2—C7 | -172.09 (18) | C1—C8—C9—C10 | 146.5 (3) |
| C7—C2—C3—C4 | -0.1 (3) | O1—C8—C9—C10 | -36.0 (3) |

supplementary materials

| | | | |
|--------------|------------|-----------------|--------------|
| C1—C2—C3—C4 | -178.9 (3) | C1—C8—C9—C14 | -37.5 (4) |
| C2—C3—C4—C5 | -0.9 (4) | O1—C8—C9—C14 | 140.0 (2) |
| C2—C3—C4—C15 | 178.2 (2) | C14—C9—C10—C11 | -0.8 (4) |
| C3—C4—C5—C6 | 1.0 (4) | C8—C9—C10—C11 | 175.3 (2) |
| C15—C4—C5—C6 | -178.1 (2) | C9—C10—C11—C12 | -1.0 (4) |
| C4—C5—C6—C7 | 0.1 (4) | C10—C11—C12—C13 | 1.7 (4) |
| C8—O1—C7—C6 | 179.8 (2) | C10—C11—C12—I | -174.85 (19) |
| C8—O1—C7—C2 | 0.4 (2) | C11—C12—C13—C14 | -0.6 (4) |
| C5—C6—C7—O1 | 179.4 (2) | I—C12—C13—C14 | 175.94 (17) |
| C5—C6—C7—C2 | -1.2 (4) | C12—C13—C14—C9 | -1.2 (3) |
| C3—C2—C7—O1 | -179.3 (2) | C10—C9—C14—C13 | 1.9 (3) |
| C1—C2—C7—O1 | -0.2 (3) | C8—C9—C14—C13 | -174.2 (2) |
| C3—C2—C7—C6 | 1.3 (4) | | |

Symmetry codes: (i) $x-1, y, z-1$; (ii) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-------------------------------------|-------|-------------|-------------|---------------|
| C10—H10 \cdots Cg1 ⁱⁱⁱ | 0.95 | 3.01 | 3.617 (4) | 125 |
| C11—H11 \cdots Cg2 ⁱⁱⁱ | 0.95 | 2.77 | 3.643 (4) | 148 |

Symmetry codes: (iii) $x-1/2, -y+3/2, z-1/2$.

Fig. 1




Fig. 2

