

Acta Crystallographica Section E

Structure Reports
Online

ISSN 1600-5368

Bis[hexaaquacobalt(II)] 25,26,27,28-tetrahydroxy-2,8,14,19-tetrathiacalix[4]-arene-5,11,17,23-tetrasulfonate monohydrate

Haruo Akashi* and Misato Ichikawa

Research Institute of Natural Sciences, Okayama University of Science, Ridai-cho, Okayama 700-0005, Japan

Correspondence e-mail: akashi@high.ous.ac.jp


Received 15 April 2008; accepted 24 April 2008

 Key indicators: single-crystal X-ray study; $T = 93$ K; mean $\sigma(C-C) = 0.004$ Å; H-atom completeness 95%; R factor = 0.045; wR factor = 0.144; data-to-parameter ratio = 13.4.

In the crystal structure of the title compound, $[Co(H_2O)_6]_2 \cdot (C_{24}H_{12}O_{16}S_8) \cdot H_2O$, the thiacalix[4]arenetetrasulfonate (= TCAS⁴⁻) anions adopt a cone-type conformation with an additional water molecule as a guest molecule in the hydrophobic cavity. The TCAS⁴⁻ anions are arranged in layers in an up-down fashion. These anionic layers alternate with cationic layers consisting of rather regular octahedral cations (symmetry m). Several medium O—H...O hydrogen-bond interactions exist between the aqua ligands of the $[Co(H_2O)_6]^{2+}$ cations and the O atoms of the sulfonate groups. In addition to the two crystallographically different Co atoms, two S and four O atoms are situated on mirror planes.

Related literature

For the structure of sodium thiacalix[4]arene tetrasulfonate monohydrate, see: Akashi & Yamauchi (2003), and for the Cd salt of the same anion, see: Zhao *et al.* (2005). Assemblies of thiacalix[4]arene tetrasulfonates with several transition metal ions were described by Guo *et al.* (2004).


Experimental
Crystal data
 $[Co(H_2O)_6]_2(C_{24}H_{12}O_{16}S_8) \cdot H_2O$
 $M_r = 1164.89$

 Monoclinic, $P2_1/m$
 $a = 11.9955$ (5) Å

 $b = 14.0628$ (10) Å

 $c = 12.8907$ (11) Å

 $\beta = 95.638$ (3)°
 $V = 2164.0$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation

 $\mu = 1.25$ mm⁻¹
 $T = 93$ K
 $0.40 \times 0.40 \times 0.10$ mm

Data collection

 Rigaku R-Axis-IV diffractometer
 Absorption correction: multi-scan
 (*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.767$, $T_{\max} = 0.883$

 15194 measured reflections
 4319 independent reflections
 3640 reflections with $F^2 > 2\sigma(F^2)$
 $R_{\text{int}} = 0.047$
Refinement
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.144$
 $S = 1.00$
 4296 reflections

 320 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 1.65$ e Å⁻³
 $\Delta\rho_{\min} = -0.81$ e Å⁻³
Table 1

Selected bond lengths (Å).

Co1—O9	2.079 (3)	Co2—O13	2.076 (3)
Co1—O10	2.064 (3)	Co2—O14	2.090 (3)
Co1—O11	2.085 (2)	Co2—O15	2.078 (2)
Co1—O12	2.088 (2)	Co2—O16	2.085 (2)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1—H5...O1 ⁱ	0.78	2.17	2.877 (3)	151
O2—H6...O1	0.88	2.26	3.082 (3)	156
O2—H6...O12 ⁱⁱ	0.88	2.71	3.154 (3)	113
O9—H7...O4 ⁱⁱⁱ	0.82	2.06	2.878 (3)	173
O9—H7...O4 ⁱⁱⁱ	0.82	2.06	2.878 (3)	173
O10—H8...O8 ^{iv}	0.93	1.83	2.729 (2)	163
O10—H8...O8 ^{iv}	0.93	1.83	2.729 (2)	163
O11—H9...O3 ^v	0.81	1.95	2.741 (3)	165
O11—H10...O7 ^v	0.77	2.04	2.795 (3)	169
O12—H11...O5	0.91	1.86	2.764 (3)	173
O12—H12...O6 ^{iv}	0.82	2.00	2.812 (3)	174
O13—H13...O7 ^{vi}	0.81	1.95	2.739 (2)	165
O13—H13...O7 ^{vi}	0.81	1.95	2.739 (2)	165
O14—H14...O5	0.78	2.05	2.818 (2)	172
O14—H14...O5	0.78	2.05	2.818 (2)	172
O15—H15...O6 ^{vii}	0.84	1.91	2.739 (3)	170
O15—H16...O4 ^v	0.87	1.89	2.743 (3)	164
O16—H17...O8 ^{viii}	0.77	2.01	2.764 (3)	166
O16—H18...O3 ^{viii}	0.84	1.92	2.719 (3)	159

Symmetry codes: (i) $x, -y + \frac{1}{2}, z$; (ii) $-x + 1, -y + 1, -z + 1$; (iii) $-x, y - \frac{1}{2}, -z + 1$; (iv) $x, y, z + 1$; (v) $-x, -y + 1, -z + 1$; (vi) $-x, y - \frac{1}{2}, -z$; (vii) $-x, -y + 1, -z$; (viii) $x, -y + \frac{1}{2}, z$.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalStructure* (Rigaku/MS, 2006); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *CrystalStructure*; molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure*.

This research was partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research, 18550063, 2007.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WM2177).

References

- Akashi, H. & Yamauchi, T. (2003). *Acta Cryst.* **E59**, m336–m338.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Guo, Q.-L., Zhu, W.-X., Ma, S.-L., Dong, S.-J. & Xu, M.-Q. (2004). *Polyhedron*, **23**, 1461–1466.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSO (2006). *CrystalStructure*. Rigaku/MSO, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Zhao, B.-T., Hu, P.-Z., Wang, J.-G., Ma, L.-F., Wang, L.-Y. & Ng, S. W. (2005). *Acta Cryst.* **E61**, m1957–m1959.

supplementary materials

Acta Cryst. (2008). E64, m797-m798 [doi:10.1107/S1600536808011781]

Bis[hexaaquacobalt(II)] 25,26,27,28-tetrahydroxy-2,8,14,19-tetrathiacalix[4]arene-5,11,17,23-tetrasulfonate monohydrate

H. Akashi and M. Ichikawa

Comment

The water soluble thiocalix[4]arene tetrasulfonate anion (= TCAS⁴⁻) is well-known as a second-sphere ligand of various transition metal coordination compounds containing complex metal aqua cations (Zhao, 2005). In the title compound, [Co(H₂O)₆]₂(S₈C₂₄O₁₆H₁₂)·H₂O, (I), the second-sphere TCAS⁴⁻ anionic ligands are linked to the aqua ligands of [Co(H₂O)₆] via several medium O-H...H hydrogen bonding interactions (see Table 2).

In the crystal structure of (I) the TCAS⁴⁻ molecules adopt the cone-type conformation. The Co²⁺ ions are not directly bonded to the TCAS⁴⁻ moieties, but exist as two octahedral aqua complexes, [Co(H₂O)₆]²⁺, both with *m* symmetry (Figure 1, Table 1). Layers of TCAS⁴⁻ anions are arranged in an up-down fashion and alternate with cationic layers that contain the [Co(H₂O)₆]²⁺ cations (Figure 2).

The TCAS⁴⁻ molecule include a water molecule as a guest-molecule in the hydrophobic cavity. Interactions between the H atoms of the aqua ligands of [Co(H₂O)₆]²⁺ and of O atoms of the TCAS⁴⁻ moieties constitute a three-dimensional hydrogen bond network, and from there the second-sphere coordination is established.

Experimental

A solution of Na₄(S₈C₂₄O₁₆H₁₂)·14H₂O (=Na₄TCAS·14H₂O) in 2 M HCl and a 2M HCl solution containing CoCl₂·6H₂O were mixed in the molar ratio (Co/Na₄TCAS = 5), poured into a vial and stirred for 1 h. This solution was then kept at room temperature from which pale pink crystals of (I) were obtained within a few days. Elemental analysis of (I) is in agreement with the refined structure model. Found: C 24.85, H 2.89%; calculated for (I): C 24.74, H 3.29%.

Refinement

Refinement was performed using 4296 reflections ($5 < 2\theta < 51^\circ$) out of 4319 independent reflections. The positions of all the H atoms were initially located from a difference map. All H atoms were refined by using the riding model approximation with C—H distances in the range 0.77–0.93 Å. The isotropic displacement parameters for all H atoms were fixed at 1.2 times the value of the equivalent isotropic displacement parameter of their carrier atom. The H atoms attached to the water molecule in the hydrophobic cavity of thiocalix[4]arenetetrasulfonate could not be found from a difference map. Therefore the positions of the H atoms were not refined. The maximum of the remaining electron density is located 2.66 Å away from atom O15.

Figures




Fig. 1. The molecular entities of (I) with the atom-labelling scheme and displacement ellipsoids drawn at 50% probability level. Labelling of H atoms has been omitted for clarity. [Symmetry code: (i) $x, 1/2 - y, z$.]




Fig. 2. The projection of the structure of (I) along the b axis.

Bis[hexaaquacobalt(II)] 25,26,27,28-tetrahydroxy-2,8,14,19-tetrathiacalix[4]arene-5,11,17,23-tetrasulfonate monohydrate

Crystal data

$[\text{Co}(\text{H}_2\text{O})_6]_2(\text{C}_{24}\text{H}_{12}\text{O}_{16}\text{S}_8) \cdot \text{H}_2\text{O}$

$M_r = 1164.89$

Monoclinic, $P2_1/m$

Hall symbol: $-P\ 2yb$

$a = 11.9955\ (5)\ \text{\AA}$

$b = 14.0628\ (10)\ \text{\AA}$

$c = 12.8907\ (11)\ \text{\AA}$

$\beta = 95.638\ (3)^\circ$

$V = 2164.0\ (3)\ \text{\AA}^3$

$Z = 2$

$F_{000} = 1192.00$

$D_x = 1.788\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71069\ \text{\AA}$

Cell parameters from 14562 reflections

$\theta = 2.2\text{--}27.5^\circ$

$\mu = 1.25\ \text{mm}^{-1}$

$T = 93\ \text{K}$

Block, pale pink

$0.40 \times 0.40 \times 0.10\ \text{mm}$

Data collection

Rigaku R-AXIS-IV
diffractometer

Detector resolution: $10.00\ \text{pixels mm}^{-1}$

ω scans

Absorption correction: multi-scan
(ABSCOR; Higashi, 1995)

$T_{\min} = 0.767$, $T_{\max} = 0.883$

15194 measured reflections

4319 independent reflections

3640 reflections with $F^2 > 2\sigma(F^2)$

$R_{\text{int}} = 0.047$

$\theta_{\text{max}} = 25.5^\circ$

$h = -15 \rightarrow 15$

$k = -17 \rightarrow 17$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[0.0031F_o^2 + \sigma(F_o^2)]/(4F_o^2)$
$wR(F^2) = 0.144$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.00$	$\Delta\rho_{\max} = 1.65 \text{ e } \text{\AA}^{-3}$
4296 reflections	$\Delta\rho_{\min} = -0.81 \text{ e } \text{\AA}^{-3}$
320 parameters	Extinction correction: none

Special details

Refinement. Refinement was performed using 4296 reflections ($5 < 2\theta < 51^\circ$). The weighted R -factor (wR) and goodness of fit (S) are based on F^2 . R factor (gt) are based on F . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.04693 (6)	0.2500	0.75294 (5)	0.01428 (17)
Co2	0.00632 (6)	0.2500	0.24647 (5)	0.01639 (17)
S1	0.50722 (9)	0.7500	0.56024 (9)	0.0120 (2)
S2	0.53799 (7)	0.47539 (6)	0.25884 (6)	0.01238 (19)
S3	0.52961 (10)	0.7500	-0.04805 (9)	0.0129 (2)
S4	0.15582 (6)	0.50788 (5)	0.45489 (6)	0.00843 (18)
S5	0.17521 (6)	0.50752 (5)	-0.03055 (6)	0.00836 (18)
O1	0.60482 (18)	0.64771 (18)	0.38463 (19)	0.0174 (5)
O2	0.61612 (19)	0.64728 (17)	0.14682 (18)	0.0162 (5)
O3	0.10300 (19)	0.50174 (16)	0.34755 (17)	0.0122 (5)
O4	0.10609 (19)	0.58262 (17)	0.51376 (17)	0.0130 (5)
O5	0.15942 (19)	0.41514 (16)	0.50631 (17)	0.0135 (5)
O6	0.1697 (2)	0.50641 (16)	-0.14431 (18)	0.0132 (5)
O7	0.10106 (19)	0.57942 (16)	0.00710 (17)	0.0121 (5)
O8	0.15982 (19)	0.41379 (16)	0.01357 (17)	0.0117 (5)
O9	-0.0351 (3)	0.250000 (10)	0.6031 (2)	0.0206 (9)
O10	0.1310 (4)	0.250000 (10)	0.9006 (2)	0.0393 (12)
O11	-0.0636 (2)	0.35446 (18)	0.79486 (19)	0.0257 (6)
O12	0.1547 (2)	0.35660 (17)	0.71050 (18)	0.0184 (6)
O13	-0.0660 (4)	0.250000 (10)	0.0934 (2)	0.0380 (11)
O14	0.0769 (3)	0.250000 (10)	0.4013 (2)	0.0178 (8)
O15	-0.1064 (2)	0.35324 (17)	0.28526 (18)	0.0201 (6)
O16	0.1180 (2)	0.14395 (18)	0.21098 (19)	0.0240 (6)
O17	0.3199 (9)	0.7500	0.2349 (9)	0.154 (4)
C1	0.5066 (2)	0.6116 (2)	0.4082 (2)	0.0123 (7)
C2	0.4618 (2)	0.5311 (2)	0.3545 (2)	0.0107 (7)
C3	0.3575 (2)	0.4978 (2)	0.3728 (2)	0.0113 (7)
C4	0.2960 (2)	0.5438 (2)	0.4440 (2)	0.0103 (7)
C5	0.3419 (2)	0.6201 (2)	0.5015 (2)	0.0107 (7)

supplementary materials

C6	0.4478 (2)	0.6531 (2)	0.4843 (2)	0.0102 (7)
C7	0.5189 (2)	0.6115 (2)	0.1034 (2)	0.0114 (7)
C8	0.4647 (2)	0.6537 (2)	0.0126 (2)	0.0098 (7)
C9	0.3610 (2)	0.6200 (2)	-0.0297 (2)	0.0093 (7)
C10	0.3118 (2)	0.5430 (2)	0.0163 (2)	0.0106 (7)
C11	0.3666 (2)	0.4982 (2)	0.1027 (2)	0.0108 (7)
C12	0.4701 (2)	0.5325 (2)	0.1461 (2)	0.0105 (7)
H1	0.3306	0.4470	0.3377	0.013*
H2	0.2996	0.6492	0.5510	0.013*
H3	0.3269	0.6542	-0.0940	0.011*
H4	0.3380	0.4414	0.1315	0.013*
H5	0.6164	0.6994	0.4048	0.021*
H6	0.6272	0.6330	0.2133	0.020*
H7	-0.0525	0.2040	0.5657	0.025*
H8	0.1529	0.2997	0.9450	0.046*
H9	-0.0875	0.3956	0.7543	0.032*
H10	-0.0648	0.3742	0.8501	0.031*
H11	0.1560	0.3806	0.6450	0.022*
H12	0.1556	0.4022	0.7502	0.022*
H13	-0.0783	0.2055	0.0540	0.045*
H14	0.1057	0.2940	0.4286	0.021*
H15	-0.1274	0.4002	0.2485	0.024*
H16	-0.0996	0.3831	0.3450	0.024*
H17	0.1182	0.1257	0.1545	0.030*
H18	0.1217	0.0916	0.2420	0.030*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0313 (4)	0.0051 (3)	0.0070 (3)	0.0000	0.0045 (2)	0.0000
Co2	0.0372 (4)	0.0050 (3)	0.0075 (3)	0.0000	0.0052 (2)	0.0000
S1	0.0131 (5)	0.0110 (5)	0.0113 (5)	0.0000	-0.0020 (4)	0.0000
S2	0.0133 (3)	0.0130 (4)	0.0113 (3)	0.0074 (2)	0.0034 (3)	0.0013 (2)
S3	0.0145 (5)	0.0112 (5)	0.0143 (5)	0.0000	0.0088 (4)	0.0000
S4	0.0107 (3)	0.0076 (3)	0.0075 (3)	0.0005 (2)	0.0029 (3)	-0.0007 (2)
S5	0.0114 (3)	0.0058 (3)	0.0080 (3)	0.0007 (2)	0.0011 (3)	-0.0010 (2)
O1	0.0095 (10)	0.0191 (12)	0.0240 (12)	-0.0005 (9)	0.0047 (10)	0.0014 (10)
O2	0.0136 (10)	0.0168 (12)	0.0181 (11)	-0.0003 (9)	0.0001 (9)	0.0011 (9)
O3	0.0180 (11)	0.0100 (12)	0.0089 (10)	-0.0007 (8)	0.0021 (9)	-0.0004 (8)
O4	0.0146 (10)	0.0111 (11)	0.0142 (10)	0.0021 (8)	0.0063 (9)	-0.0032 (8)
O5	0.0169 (11)	0.0101 (11)	0.0135 (10)	-0.0003 (9)	0.0009 (9)	0.0023 (9)
O6	0.0186 (11)	0.0106 (11)	0.0100 (10)	-0.0004 (9)	-0.0007 (9)	-0.0006 (8)
O7	0.0126 (10)	0.0092 (11)	0.0152 (10)	0.0031 (8)	0.0045 (9)	-0.0004 (8)
O8	0.0167 (11)	0.0077 (11)	0.0104 (10)	-0.0015 (8)	0.0001 (9)	0.0003 (8)
O9	0.042 (2)	0.0102 (17)	0.0094 (15)	0.0000	0.0008 (15)	0.0000
O10	0.095 (3)	0.0078 (18)	0.0115 (16)	0.0000	-0.015 (2)	0.0000
O11	0.0573 (18)	0.0121 (12)	0.0094 (10)	0.0091 (12)	0.0126 (12)	0.0010 (9)
O12	0.0365 (14)	0.0086 (11)	0.0108 (10)	-0.0018 (10)	0.0051 (10)	-0.0022 (9)

O13	0.094 (3)	0.0091 (18)	0.0086 (15)	0.0000	-0.009 (2)	0.0000
O14	0.034 (2)	0.0075 (16)	0.0118 (15)	0.0000	0.0012 (15)	0.0000
O15	0.0406 (15)	0.0089 (11)	0.0106 (10)	0.0045 (10)	0.0011 (11)	-0.0006 (9)
O16	0.0523 (17)	0.0089 (12)	0.0132 (11)	0.0059 (11)	0.0150 (12)	0.0015 (9)
O17	0.141 (9)	0.149 (9)	0.180 (10)	0.0000	0.054 (8)	0.0000
C1	0.0094 (13)	0.0123 (16)	0.0148 (14)	0.0035 (12)	-0.0011 (12)	0.0048 (12)
C2	0.0150 (15)	0.0087 (15)	0.0084 (13)	0.0059 (12)	0.0009 (12)	0.0008 (11)
C3	0.0148 (14)	0.0095 (16)	0.0098 (14)	0.0032 (12)	0.0016 (12)	0.0002 (11)
C4	0.0133 (14)	0.0071 (15)	0.0110 (13)	-0.0002 (11)	0.0031 (12)	0.0023 (11)
C5	0.0133 (14)	0.0090 (15)	0.0103 (14)	0.0042 (11)	0.0040 (12)	0.0007 (11)
C6	0.0132 (14)	0.0076 (14)	0.0092 (13)	0.0026 (11)	-0.0019 (12)	-0.0006 (11)
C7	0.0091 (13)	0.0132 (16)	0.0124 (14)	0.0043 (12)	0.0038 (12)	-0.0030 (12)
C8	0.0114 (14)	0.0091 (15)	0.0103 (13)	0.0006 (11)	0.0075 (12)	0.0006 (11)
C9	0.0101 (13)	0.0091 (15)	0.0088 (13)	0.0005 (11)	0.0008 (12)	-0.0016 (11)
C10	0.0128 (14)	0.0070 (15)	0.0123 (13)	0.0026 (11)	0.0031 (12)	-0.0004 (11)
C11	0.0175 (15)	0.0078 (15)	0.0077 (13)	0.0030 (12)	0.0040 (13)	-0.0013 (11)
C12	0.0129 (14)	0.0092 (15)	0.0097 (13)	0.0070 (11)	0.0027 (12)	-0.0008 (11)

Geometric parameters (Å, °)

Co1—O9	2.079 (3)	C3—C4	1.392 (4)
Co1—O10	2.064 (3)	C4—C5	1.386 (4)
Co1—O11	2.085 (2)	C5—C6	1.390 (4)
Co1—O11 ⁱ	2.085 (2)	C7—C8	1.413 (4)
Co1—O12	2.088 (2)	C7—C12	1.393 (4)
Co1—O12 ⁱ	2.088 (2)	C8—C9	1.391 (4)
Co2—O13	2.076 (3)	C9—C10	1.394 (4)
Co2—O14	2.090 (3)	C10—C11	1.387 (4)
Co2—O15	2.078 (2)	C11—C12	1.397 (4)
Co2—O15 ⁱ	2.078 (2)	O1—H5	0.779
Co2—O16	2.085 (2)	O2—H6	0.878
Co2—O16 ⁱ	2.085 (2)	O9—H7	0.822
S1—C6	1.786 (3)	O9—H7 ⁱ	0.822
S1—C6 ⁱⁱ	1.786 (3)	O10—H8	0.925
S2—C2	1.786 (3)	O10—H8 ⁱ	0.925
S2—C12	1.785 (3)	O11—H9	0.813
S3—C8	1.781 (3)	O11—H10	0.766
S3—C8 ⁱⁱ	1.781 (3)	O12—H11	0.911
S4—O3	1.467 (2)	O12—H12	0.820
S4—O4	1.458 (2)	O13—H13	0.811
S4—O5	1.462 (2)	O13—H13 ⁱ	0.811
S4—C4	1.775 (3)	O14—H14	0.776
S5—O6	1.462 (2)	O14—H14 ⁱ	0.776
S5—O7	1.461 (2)	O15—H15	0.837
S5—O8	1.455 (2)	O15—H16	0.874
S5—C10	1.762 (3)	O16—H17	0.772
O1—C1	1.345 (3)	O16—H18	0.837

supplementary materials

O2—C7	1.341 (3)	C3—H1	0.888
C1—C2	1.406 (4)	C5—H2	0.947
C1—C6	1.392 (4)	C9—H3	1.009
C2—C3	1.378 (4)	C11—H4	0.958
O9—Co1—O10	179.01 (18)	C3—C4—C5	120.3 (2)
O9—Co1—O11	89.19 (10)	C4—C5—C6	119.6 (2)
O9—Co1—O11 ⁱ	89.19 (10)	S1—C6—C1	120.2 (2)
O9—Co1—O12	90.31 (10)	S1—C6—C5	119.3 (2)
O9—Co1—O12 ⁱ	90.31 (10)	C1—C6—C5	120.4 (2)
O10—Co1—O11	91.51 (12)	O2—C7—C8	119.9 (2)
O10—Co1—O11 ⁱ	91.51 (12)	O2—C7—C12	121.1 (2)
O10—Co1—O12	89.00 (12)	C8—C7—C12	119.0 (2)
O10—Co1—O12 ⁱ	89.00 (12)	S3—C8—C7	119.9 (2)
O11—Co1—O11 ⁱ	89.58 (11)	S3—C8—C9	119.9 (2)
O11—Co1—O12	89.31 (10)	C7—C8—C9	120.2 (2)
O11—Co1—O12 ⁱ	178.79 (10)	C8—C9—C10	119.8 (2)
O11 ⁱ —Co1—O12	178.79 (10)	S5—C10—C9	119.5 (2)
O11 ⁱ —Co1—O12 ⁱ	89.31 (10)	S5—C10—C11	119.8 (2)
O12—Co1—O12 ⁱ	91.80 (10)	C9—C10—C11	120.5 (2)
O13—Co2—O14	179.19 (18)	C10—C11—C12	119.8 (2)
O13—Co2—O15	90.44 (12)	S2—C12—C7	120.3 (2)
O13—Co2—O15 ⁱ	90.44 (12)	S2—C12—C11	119.1 (2)
O13—Co2—O16	90.52 (12)	C7—C12—C11	120.5 (2)
O13—Co2—O16 ⁱ	90.52 (12)	C1—O1—H5	114.1
O14—Co2—O15	88.99 (9)	C7—O2—H6	111.3
O14—Co2—O15 ⁱ	88.99 (9)	Co1—O9—H7	128.0
O14—Co2—O16	90.04 (10)	Co1—O9—H7 ⁱ	128.0
O14—Co2—O16 ⁱ	90.04 (10)	H7—O9—H7 ⁱ	103.8
O15—Co2—O15 ⁱ	88.64 (10)	Co1—O10—H8	130.8
O15—Co2—O16	178.36 (10)	Co1—O10—H8 ⁱ	130.8
O15—Co2—O16 ⁱ	90.02 (10)	H8—O10—H8 ⁱ	98.2
O15 ⁱ —Co2—O16	90.02 (10)	Co1—O11—H9	121.8
O15 ⁱ —Co2—O16 ⁱ	178.36 (10)	Co1—O11—H10	124.5
O16—Co2—O16 ⁱ	91.31 (10)	H9—O11—H10	107.6
C6—S1—C6 ⁱⁱ	99.56 (15)	Co1—O12—H11	125.2
C2—S2—C12	98.15 (14)	Co1—O12—H12	111.6
C8—S3—C8 ⁱⁱ	99.04 (15)	H11—O12—H12	106.7
O3—S4—O4	111.87 (13)	Co2—O13—H13	129.0
O3—S4—O5	111.47 (12)	Co2—O13—H13 ⁱ	129.0
O3—S4—C4	105.50 (14)	H13—O13—H13 ⁱ	101.1
O4—S4—O5	113.74 (13)	Co2—O14—H14	123.7
O4—S4—C4	105.98 (14)	Co2—O14—H14 ⁱ	123.7
O5—S4—C4	107.68 (13)	H14—O14—H14 ⁱ	105.7

O6—S5—O7	111.83 (13)	Co2—O15—H15	125.6
O6—S5—O8	112.81 (13)	Co2—O15—H16	122.8
O6—S5—C10	107.21 (14)	H15—O15—H16	96.7
O7—S5—O8	113.11 (13)	Co2—O16—H17	120.4
O7—S5—C10	105.42 (14)	Co2—O16—H18	121.9
O8—S5—C10	105.79 (14)	H17—O16—H18	98.9
O1—C1—C2	119.6 (2)	C2—C3—H1	118.5
O1—C1—C6	121.1 (2)	C4—C3—H1	121.3
C2—C1—C6	119.2 (2)	C4—C5—H2	118.9
S2—C2—C1	119.7 (2)	C6—C5—H2	121.4
S2—C2—C3	120.2 (2)	C8—C9—H3	115.8
C1—C2—C3	120.0 (2)	C10—C9—H3	124.4
C2—C3—C4	120.2 (2)	C10—C11—H4	122.0
S4—C4—C3	119.1 (2)	C12—C11—H4	118.1
S4—C4—C5	120.5 (2)		

Symmetry codes: (i) $x, -y+1/2, z$; (ii) $x, -y+3/2, z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H5 \cdots O1 ⁱⁱ	0.78	2.17	2.877 (3)	151
O2—H6 \cdots O1	0.88	2.26	3.082 (3)	156
O2—H6 \cdots O12 ⁱⁱⁱ	0.88	2.71	3.154 (3)	113
O9—H7 \cdots O4 ^{iv}	0.82	2.06	2.878 (3)	173
O9—H7 \cdots O4 ^{iv}	0.82	2.06	2.878 (3)	173
O10—H8 \cdots O8 ^v	0.93	1.83	2.729 (2)	163
O10—H8 \cdots O8 ^v	0.93	1.83	2.729 (2)	163
O11—H9 \cdots O3 ^{vi}	0.81	1.95	2.741 (3)	165
O11—H10 \cdots O7 ^{vi}	0.77	2.04	2.795 (3)	169
O12—H11 \cdots O5	0.91	1.86	2.764 (3)	173
O12—H12 \cdots O6 ^v	0.82	2.00	2.812 (3)	174
O13—H13 \cdots O7 ^{vii}	0.81	1.95	2.739 (2)	165
O13—H13 \cdots O7 ^{vii}	0.81	1.95	2.739 (2)	165
O14—H14 \cdots O5	0.78	2.05	2.818 (2)	172
O14—H14 \cdots O5	0.78	2.05	2.818 (2)	172
O15—H15 \cdots O6 ^{viii}	0.84	1.91	2.739 (3)	170
O15—H16 \cdots O4 ^{vi}	0.87	1.89	2.743 (3)	164
O16—H17 \cdots O8 ^{ix}	0.77	2.01	2.764 (3)	166
O16—H18 \cdots O3 ^{ix}	0.84	1.92	2.719 (3)	159

Symmetry codes: (ii) $x, -y+3/2, z$; (iii) $-x+1, -y+1, -z+1$; (iv) $-x, y-1/2, -z+1$; (v) $x, y, z+1$; (vi) $-x, -y+1, -z+1$; (vii) $-x, y-1/2, -z$; (viii) $-x, -y+1, -z$; (ix) $x, -y+1/2, z$.

Fig. 1




Fig. 2

