

Hexaaquanickel(II) bis[4-(2-hydroxybenzylideneamino)benzenesulfonate]Xi-Shi Tai,^{a*} Jun Xu,^b Yi-Min Feng^a and Zu-Pei Liang^a

^aDepartment of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China, and ^bWeifang Institute of Supervision and Inspection, on Product Quality, Weifang 261031, People's Republic of China
Correspondence e-mail: taixishi@lzu.edu.cn


Received 16 May 2008; accepted 21 May 2008

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.011\text{ \AA}$; R factor = 0.067; wR factor = 0.138; data-to-parameter ratio = 13.0.

In the title compound, $[\text{Ni}(\text{H}_2\text{O})_6](\text{C}_{13}\text{H}_{10}\text{NO}_4\text{S})_2$, the nickel(II) atom, lying on a center of symmetry, is six-coordinated by six aqua O-atom donors. The dihedral angle between the two benzene rings is $33.1(3)^\circ$. The crystal structure is stabilized by aqua-anion O-H \cdots O hydrogen bonds. Intramolecular O-H \cdots N and C-H \cdots O hydrogen-bonding interactions occur in the anion.

Related literature

For related literature, see: Tai & Feng (2008); Tai *et al.* (2003, 2008); Tai, Yin & Feng (2007); Tai, Yin & Hao (2007); Tai, Yin, Feng & Kong (2007); Wang *et al.* (2007).

**Experimental***Crystal data* $M_r = 719.37$ Monoclinic, $P2_1/c$ $a = 6.3047(6)\text{ \AA}$ $b = 35.193(3)\text{ \AA}$ $c = 9.3536(10)\text{ \AA}$ $\beta = 131.822(2)^\circ$ $V = 1546.6(3)\text{ \AA}^3$ $Z = 2$ Mo $K\alpha$ radiation $\mu = 0.83\text{ mm}^{-1}$ $T = 298(2)\text{ K}$ $0.43 \times 0.38 \times 0.25\text{ mm}$ **Data collection**

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.716$, $T_{\max} = 0.819$

7465 measured reflections
2661 independent reflections
2428 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.067$
 $wR(F^2) = 0.137$
 $S = 1.28$
2661 reflections

205 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.32\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.00\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4-H4 \cdots N1	0.82	1.86	2.595 (9)	148
O5-H5A \cdots O1 ⁱ	0.85	1.96	2.737 (6)	151
O5-H5B \cdots O2 ⁱⁱ	0.85	1.98	2.751 (6)	150
O6-H6A \cdots O3 ⁱ	0.85	1.95	2.764 (7)	161
O6-H6B \cdots O1 ⁱⁱⁱ	0.85	1.97	2.768 (8)	156
O7-H7A \cdots O2	0.85	2.00	2.757 (8)	148
O7-H7B \cdots O3 ⁱⁱ	0.85	2.00	2.769 (7)	150
C2-H2 \cdots O3	0.93	2.56	2.917 (7)	104

Symmetry codes: (i) $x, y, z + 1$; (ii) $x - 1, y, z$; (iii) $x + 1, y, z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (20671073), the Natural Science Foundation of Shandong (Y2007B60), the Science and Technology Foundation of Weifang and Weifang University for research grants.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2568).

References

- Bruker (2000). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Tai, X.-S. & Feng, Y.-M. (2008). *Acta Cryst. E* **64**, o707.
- Tai, X.-S., Feng, Y.-M. & Zhang, H.-X. (2008). *Acta Cryst. E* **64**, m502.
- Tai, X. S., Yin, J. & Feng, Y. M. (2007). *Z. Kristallogr. New Cryst. Struct.* **222**, 398–400.
- Tai, X. S., Yin, J., Feng, Y. M. & Kong, F. Y. (2007). *Chin. J. Inorg. Chem.* **23**, 1812–1814.
- Tai, X.-S., Yin, J. & Hao, M.-Y. (2007). *Acta Cryst. E* **63**, m1061–m1062.
- Tai, X.-S., Yin, X.-H., Tan, M.-Y. & Li, Y.-Z. (2003). *Acta Cryst. E* **59**, o681–o682.
- Wang, L.-H., Yin, J. & Tai, X.-S. (2007). *Acta Cryst. E* **63**, m1664.

supporting information

Acta Cryst. (2008). E64, m893 [doi:10.1107/S1600536808015262]

Hexaaquanickel(II) bis[4-(2-hydroxybenzylideneamino)benzenesulfonate]

Xi-Shi Tai, Jun Xu, Yi-Min Feng and Zu-Pei Liang

S1. Comment

As part of our ongoing studies of the coordination chemistry of Schiffbase ligands (Tai *et al.*, 2003; (Tai, Yin & Feng, 2007; Tai, Yin, Feng & Kong, 2007; Tai, Yin & Hao, 2007; Wang *et al.*, 2007; Tai *et al.*, 2008; Tai *et al.*, 2008), we now report the synthesis and structure of the title compound, (I), (Fig. 1).

In the molecule of (I), the nickel (II) center is six-coordinate with six O donor of H₂O. The C7—N1 [1.288 (8) Å] is close to double-bond. Otherwise, the geometrical parameters for (I) are normal. The dihedral angle between the two benzene rings is 33.1°. The packing is stabilized by O_{water}—H···O_{anion} hydrogen bonds. The intramolecular O—H···N and C—H···O hydrogen bonding interactions occur in the anion. (Table 1).

S2. Experimental

1 mmol of Ni(CH₃COO)₂·4H₂O was added to a solution of salicylaldehyde-4-aminobenzene sulfonic acid (1 mmol) in 10 ml of 95% ethanol. The mixture was stirred for 2 h at refluxing temperature. Evaporating some ethanol, clear blocks of (I) were obtained after one weeks.

S3. Refinement

The H atoms were placed geometrically (C—H = 0.93 Å and O—H = 0.82–0.85 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{carrier})$.




Figure 1

The molecular structure of (I) showing 30% displacement ellipsoids.

Hexaaquanickel(II) bis[4-(2-hydroxybenzylideneamino)benzenesulfonate]

Crystal data

[Ni(H₂O)₆](C₁₃H₁₀NO₄S)₂

$M_r = 719.37$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 6.3047 (6)$ Å

$b = 35.193 (3)$ Å

$c = 9.3536 (10)$ Å

$\beta = 131.822 (2)^\circ$

$V = 1546.6 (3) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 748$
 $D_x = 1.545 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 4537 reflections

$\theta = 2.9\text{--}27.7^\circ$
 $\mu = 0.83 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
Block, colourless
 $0.43 \times 0.38 \times 0.25 \text{ mm}$

Data collection

Bruker SMART CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2000)
 $T_{\min} = 0.716$, $T_{\max} = 0.819$

7465 measured reflections
2661 independent reflections
2428 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.3^\circ$
 $h = -7 \rightarrow 5$
 $k = -41 \rightarrow 40$
 $l = -9 \rightarrow 11$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.067$
 $wR(F^2) = 0.137$
 $S = 1.28$
2661 reflections
205 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0161P)^2 + 4.8144P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.32 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -1.00 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.5000	1.0000	1.0000	0.0310 (2)
N1	0.4029 (11)	0.76969 (14)	0.4874 (8)	0.0565 (13)
O1	0.3354 (7)	0.95021 (11)	0.3248 (5)	0.0486 (10)
O2	0.6843 (8)	0.94957 (11)	0.6726 (5)	0.0462 (9)
O3	0.8302 (7)	0.93929 (10)	0.4933 (5)	0.0392 (8)
O4	0.0669 (10)	0.71618 (13)	0.4256 (8)	0.0767 (14)
H4	0.1257	0.7379	0.4400	0.115*
O5	0.2072 (7)	0.97532 (11)	0.9978 (5)	0.0474 (9)
H5A	0.2819	0.9621	1.0975	0.071*
H5B	0.0852	0.9623	0.8972	0.071*
O6	0.7945 (8)	0.96016 (12)	1.1929 (6)	0.0549 (11)

H6A	0.8124	0.9592	1.2913	0.082*
H6B	0.9559	0.9642	1.2273	0.082*
O7	0.3791 (8)	0.96590 (13)	0.7796 (6)	0.0600 (12)
H7A	0.4709	0.9703	0.7444	0.090*
H7B	0.2009	0.9665	0.6838	0.090*
S1	0.6046 (2)	0.93471 (4)	0.49681 (17)	0.0319 (3)
C1	0.5566 (9)	0.88532 (14)	0.4986 (7)	0.0313 (10)
C2	0.6556 (12)	0.86005 (16)	0.4406 (8)	0.0451 (13)
H2	0.7524	0.8689	0.4039	0.054*
C3	0.6082 (14)	0.82145 (16)	0.4382 (9)	0.0529 (15)
H3	0.6758	0.8042	0.4012	0.063*
C4	0.4623 (12)	0.80853 (16)	0.4899 (9)	0.0483 (14)
C5	0.3614 (13)	0.83417 (16)	0.5454 (9)	0.0509 (15)
H5	0.2614	0.8253	0.5795	0.061*
C6	0.4079 (12)	0.87246 (15)	0.5503 (8)	0.0438 (13)
H6	0.3404	0.8896	0.5878	0.053*
C7	0.5721 (13)	0.74368 (17)	0.5168 (9)	0.0525 (15)
H7	0.7384	0.7507	0.5446	0.063*
C8	0.5084 (15)	0.70327 (16)	0.5074 (9)	0.0527 (14)
C9	0.2544 (14)	0.69149 (18)	0.4569 (9)	0.0589 (17)
C10	0.1982 (16)	0.65268 (18)	0.4407 (9)	0.0643 (18)
H10	0.0295	0.6445	0.4068	0.077*
C11	0.3845 (17)	0.6265 (2)	0.4733 (9)	0.068 (2)
H11	0.3396	0.6008	0.4588	0.082*
C12	0.6424 (19)	0.6376 (2)	0.5281 (11)	0.077 (2)
H12	0.7726	0.6196	0.5549	0.092*
C13	0.7008 (15)	0.67614 (18)	0.5417 (9)	0.0610 (17)
H13	0.8689	0.6840	0.5739	0.073*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0214 (4)	0.0436 (5)	0.0287 (4)	-0.0007 (4)	0.0170 (4)	-0.0012 (4)
N1	0.060 (3)	0.044 (3)	0.061 (3)	-0.010 (2)	0.038 (3)	-0.004 (2)
O1	0.0284 (19)	0.064 (2)	0.048 (2)	0.0099 (18)	0.0234 (19)	0.0211 (19)
O2	0.042 (2)	0.056 (2)	0.051 (2)	-0.0096 (18)	0.035 (2)	-0.0105 (18)
O3	0.0269 (17)	0.055 (2)	0.041 (2)	-0.0029 (16)	0.0250 (16)	0.0003 (17)
O4	0.056 (3)	0.059 (3)	0.094 (4)	-0.008 (2)	0.042 (3)	0.002 (3)
O5	0.0309 (19)	0.074 (3)	0.035 (2)	-0.0104 (19)	0.0206 (17)	-0.0005 (19)
O6	0.038 (2)	0.082 (3)	0.055 (3)	0.018 (2)	0.035 (2)	0.025 (2)
O7	0.036 (2)	0.100 (3)	0.053 (3)	-0.023 (2)	0.033 (2)	-0.033 (2)
S1	0.0229 (6)	0.0420 (7)	0.0311 (6)	-0.0014 (5)	0.0181 (5)	0.0003 (5)
C1	0.021 (2)	0.041 (3)	0.027 (2)	-0.004 (2)	0.014 (2)	-0.002 (2)
C2	0.053 (3)	0.049 (3)	0.047 (3)	-0.003 (3)	0.039 (3)	-0.004 (3)
C3	0.066 (4)	0.046 (3)	0.054 (4)	-0.006 (3)	0.043 (3)	-0.012 (3)
C4	0.041 (3)	0.046 (3)	0.049 (3)	-0.007 (3)	0.026 (3)	0.001 (3)
C5	0.054 (4)	0.046 (3)	0.068 (4)	-0.004 (3)	0.047 (4)	0.001 (3)
C6	0.053 (3)	0.044 (3)	0.053 (3)	-0.004 (3)	0.043 (3)	0.001 (3)

C7	0.050 (4)	0.056 (4)	0.048 (4)	-0.014 (3)	0.031 (3)	-0.009 (3)
C8	0.059 (4)	0.045 (3)	0.047 (3)	-0.008 (3)	0.032 (3)	-0.008 (3)
C9	0.050 (4)	0.049 (4)	0.048 (4)	-0.003 (3)	0.021 (3)	0.002 (3)
C10	0.065 (4)	0.049 (4)	0.055 (4)	-0.014 (3)	0.030 (4)	-0.002 (3)
C11	0.082 (5)	0.050 (4)	0.051 (4)	-0.013 (4)	0.035 (4)	0.000 (3)
C12	0.098 (6)	0.049 (4)	0.079 (5)	0.000 (4)	0.057 (5)	0.002 (4)
C13	0.068 (4)	0.052 (4)	0.063 (4)	-0.007 (3)	0.044 (4)	-0.007 (3)

Geometric parameters (\AA , $^{\circ}$)

Ni1—O5 ⁱ	2.028 (3)	C1—C2	1.388 (7)
Ni1—O5	2.028 (3)	C2—C3	1.388 (8)
Ni1—O7 ⁱ	2.043 (4)	C2—H2	0.9300
Ni1—O7	2.043 (4)	C3—C4	1.371 (8)
Ni1—O6	2.049 (4)	C3—H3	0.9300
Ni1—O6 ⁱ	2.049 (4)	C4—C5	1.389 (8)
N1—C7	1.288 (8)	C5—C6	1.373 (8)
N1—C4	1.414 (7)	C5—H5	0.9300
O1—S1	1.458 (4)	C6—H6	0.9300
O2—S1	1.459 (4)	C7—C8	1.465 (8)
O3—S1	1.454 (3)	C7—H7	0.9300
O4—C9	1.333 (8)	C8—C9	1.397 (9)
O4—H4	0.8209	C8—C13	1.402 (9)
O5—H5A	0.8494	C9—C10	1.394 (8)
O5—H5B	0.8492	C10—C11	1.357 (10)
O6—H6A	0.8500	C10—H10	0.9300
O6—H6B	0.8498	C11—C12	1.396 (10)
O7—H7A	0.8497	C11—H11	0.9300
O7—H7B	0.8504	C12—C13	1.387 (9)
S1—C1	1.767 (5)	C12—H12	0.9300
C1—C6	1.387 (7)	C13—H13	0.9300
O5 ⁱ —Ni1—O5	180.0 (2)	C3—C2—H2	120.5
O5 ⁱ —Ni1—O7 ⁱ	90.78 (15)	C1—C2—H2	120.5
O5—Ni1—O7 ⁱ	89.22 (15)	C4—C3—C2	120.4 (5)
O5 ⁱ —Ni1—O7	89.22 (15)	C4—C3—H3	119.8
O5—Ni1—O7	90.78 (15)	C2—C3—H3	119.8
O7 ⁱ —Ni1—O7	180.000 (2)	C3—C4—C5	119.9 (5)
O5 ⁱ —Ni1—O6	89.99 (15)	C3—C4—N1	123.2 (6)
O5—Ni1—O6	90.01 (15)	C5—C4—N1	116.9 (5)
O7 ⁱ —Ni1—O6	90.23 (18)	C6—C5—C4	120.6 (5)
O7—Ni1—O6	89.77 (18)	C6—C5—H5	119.7
O5 ⁱ —Ni1—O6 ⁱ	90.01 (15)	C4—C5—H5	119.7
O5—Ni1—O6 ⁱ	89.99 (15)	C5—C6—C1	119.1 (5)
O7 ⁱ —Ni1—O6 ⁱ	89.77 (18)	C5—C6—H6	120.4
O7—Ni1—O6 ⁱ	90.23 (18)	C1—C6—H6	120.4
O6—Ni1—O6 ⁱ	180.000 (3)	N1—C7—C8	121.5 (6)
C7—N1—C4	120.7 (5)	N1—C7—H7	119.3

C9—O4—H4	109.5	C8—C7—H7	119.3
Ni1—O5—H5A	112.9	C9—C8—C13	119.8 (6)
Ni1—O5—H5B	112.8	C9—C8—C7	121.0 (6)
H5A—O5—H5B	110.6	C13—C8—C7	119.1 (6)
Ni1—O6—H6A	111.7	O4—C9—C8	122.0 (6)
Ni1—O6—H6B	111.9	O4—C9—C10	119.4 (6)
H6A—O6—H6B	109.6	C8—C9—C10	118.6 (7)
Ni1—O7—H7A	113.0	C11—C10—C9	121.4 (7)
Ni1—O7—H7B	112.9	C11—C10—H10	119.3
H7A—O7—H7B	110.5	C9—C10—H10	119.3
O3—S1—O1	111.7 (2)	C10—C11—C12	120.9 (7)
O3—S1—O2	112.3 (2)	C10—C11—H11	119.5
O1—S1—O2	112.5 (2)	C12—C11—H11	119.5
O3—S1—C1	106.6 (2)	C13—C12—C11	118.7 (8)
O1—S1—C1	107.3 (2)	C13—C12—H12	120.6
O2—S1—C1	105.9 (2)	C11—C12—H12	120.6
C6—C1—C2	120.8 (5)	C12—C13—C8	120.5 (7)
C6—C1—S1	119.0 (4)	C12—C13—H13	119.7
C2—C1—S1	120.2 (4)	C8—C13—H13	119.7
C3—C2—C1	119.0 (5)		
O3—S1—C1—C6	163.8 (4)	C2—C1—C6—C5	0.6 (8)
O1—S1—C1—C6	-76.3 (5)	S1—C1—C6—C5	177.7 (4)
O2—S1—C1—C6	44.1 (5)	C4—N1—C7—C8	-177.6 (5)
O3—S1—C1—C2	-19.0 (5)	N1—C7—C8—C9	3.2 (10)
O1—S1—C1—C2	100.8 (4)	N1—C7—C8—C13	-179.6 (6)
O2—S1—C1—C2	-138.8 (4)	C13—C8—C9—O4	179.1 (6)
C6—C1—C2—C3	-1.1 (8)	C7—C8—C9—O4	-3.8 (10)
S1—C1—C2—C3	-178.2 (4)	C13—C8—C9—C10	-0.4 (10)
C1—C2—C3—C4	0.8 (9)	C7—C8—C9—C10	176.8 (6)
C2—C3—C4—C5	-0.1 (10)	O4—C9—C10—C11	-179.3 (6)
C2—C3—C4—N1	178.9 (6)	C8—C9—C10—C11	0.2 (10)
C7—N1—C4—C3	29.9 (9)	C9—C10—C11—C12	1.3 (11)
C7—N1—C4—C5	-151.1 (6)	C10—C11—C12—C13	-2.4 (11)
C3—C4—C5—C6	-0.4 (10)	C11—C12—C13—C8	2.2 (10)
N1—C4—C5—C6	-179.5 (6)	C9—C8—C13—C12	-0.8 (10)
C4—C5—C6—C1	0.2 (9)	C7—C8—C13—C12	-178.0 (6)

Symmetry code: (i) $-x+1, -y+2, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O4—H4 \cdots N1	0.82	1.86	2.595 (9)	148
O5—H5A \cdots O1 ⁱⁱ	0.85	1.96	2.737 (6)	151
O5—H5B \cdots O2 ⁱⁱⁱ	0.85	1.98	2.751 (6)	150
O6—H6A \cdots O3 ⁱⁱ	0.85	1.95	2.764 (7)	161
O6—H6B \cdots O1 ^{iv}	0.85	1.97	2.768 (8)	156

O7—H7A···O2	0.85	2.00	2.757 (8)	148
O7—H7B···O3 ⁱⁱⁱ	0.85	2.00	2.769 (7)	150
C2—H2···O3	0.93	2.56	2.917 (7)	104

Symmetry codes: (ii) $x, y, z+1$; (iii) $x-1, y, z$; (iv) $x+1, y, z+1$.