

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1-Benzyl-2-(4-chlorophenyl)-4,5-diphenyl-1*H*-imidazole

 Mahmood Ghoranneviss,<sup>a</sup> Ghodsi Mohammadi Ziarani,<sup>b</sup> Alireza Abbasi,<sup>c\*</sup> Mohammad Reza Hantehzadeh<sup>a</sup> and Zahra Farahani<sup>b</sup>
<sup>a</sup>Plasma Physics Research Center, Science & Reseach Campus, Islamic Azad University, Tehran, Iran, <sup>b</sup>Department of Chemistry, University of Alzahra, Tehran, Iran, and <sup>c</sup>School of Chemistry, University College of Science, University of Tehran, Tehran, Iran

Correspondence e-mail: aabbasi@khayam.ut.ac.ir


Received 20 May 2008; accepted 2 June 2008

 Key indicators: single-crystal X-ray study;  $T = 290$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.042;  $wR$  factor = 0.091; data-to-parameter ratio = 14.7.

The molecular conformation of the title compound,  $\text{C}_{28}\text{H}_{21}\text{ClN}_2$ , is stabilized by an intramolecular  $\text{C}-\text{H}\cdots\text{N}$  hydrogen bond. It has many pharmacological properties, such as being an inhibitor of P38 MAP Kinase, and can play an important role in biochemical processes.

## Related literature

For related structures and properties, see: Balalaie *et al.* (2003); Nagarapu *et al.* (2007); Kidwai *et al.* (2007).



## Experimental

### Crystal data

$\text{C}_{28}\text{H}_{21}\text{ClN}_2$	$\gamma = 87.842$ (13) $^\circ$
$M_r = 420.92$	$V = 1082.6$ (3) Å <sup>3</sup>
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.4880$ (11) Å	Mo $K\alpha$ radiation
$b = 9.2711$ (16) Å	$\mu = 0.19$ mm <sup>-1</sup>
$c = 16.049$ (3) Å	$T = 290$ (2) K
$\alpha = 87.169$ (13) $^\circ$	$0.3 \times 0.2 \times 0.2$ mm
$\beta = 76.704$ (12) $^\circ$	

### Data collection

STOE IPDS-II diffractometer	4246 independent reflections
Absorption correction: none	2814 reflections with $I > 2\sigma(I)$
8769 measured reflections	$R_{\text{int}} = 0.034$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.090$	$\Delta\rho_{\text{max}} = 0.13$ e Å <sup>-3</sup>
$S = 0.94$	$\Delta\rho_{\text{min}} = -0.20$ e Å <sup>-3</sup>
4246 reflections	
288 parameters	

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C3}-\text{H3}\cdots\text{N1}$	0.93	2.56	2.874 (2)	100

Data collection: *X-Area* (Stoe & Cie, 1997); cell refinement: *X-Area*; data reduction: *X-Area*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2001); software used to prepare material for publication: *SHELXL97*.

This work was supported by a grant from the University of Tehran and the University of Alzahra.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2715).

## References

- Balalaie, S., Hashemi, M. M. & Akhbari, M. (2003). *Tetrahedron Lett.* **44**, 1709–1711.
- Brandenburg, K. (2001). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Kidwai, M., Mothsra, P., Bansal, V., Somvanshi, R. K., Ethayathulla, A. S., Dey, S. & Singh, T. P. (2007). *J. Mol. Catal. A Chem.* **265**, 177–182.
- Nagarapu, L., Apuri, S. & Kantevari, S. (2007). *J. Mol. Catal. A Chem.* **266**, 104–108.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (1997). *X-Area*. Stoe & Cie GmbH, Darmstadt, Germany.

**supplementary materials**

*Acta Cryst.* (2008). E64, o1233 [ doi:10.1107/S1600536808016802 ]

## 1-Benzyl-2-(4-chlorophenyl)-4,5-diphenyl-1*H*-imidazole

M. Ghoranneviss, G. Mohammadi Ziarani, A. Abbasi, M. R. Hantehzadeh and Z. Farahani

### Comment

The synthesis, reactions and biological properties of substituted imidazole constitutes a significant part of modern heterocyclic chemistry. Compounds containing an imidazole ring system have many pharmacological properties and play important roles in biochemical processes. Many of substituted diaryl imidazoles are known as inhibitors of P38 MAP Kinase (Balalaie, *et al.*, 2003; Nagarapu, *et al.*, 2007; Kidwai, *et al.*, 2007).

The molecular structure of the title compound and the atom-numbering scheme are shown in Fig. 1. The phenyl rings attached at C2, N2, C9 and C8 enclose dihedral angles of 40.74 (8)°, 85.60 (7)°, 77.10 (6)°, 16.55 (11)°, respectively, with the imidazole ring. An intramolecular hydrogen bond stabilises the molecular conformation. Dipole-dipole and van der Waals interactions are effective in the molecular packing.

### Experimental

A mixture of benzil (2.5 mmol), 4-chlorobenzaldehyde (2.5 mmol), benzylamine (2.5 mmol), ammonium acetate (5 mmol) and activated SBA-sulfonic acid (0.02 g) was heated at 140°C for 6 minutes. The progress of reaction was monitored by TLC. After cooling to room temperature, the mixture was dissolved in hot ethylacetate and the catalyst was removed by filtration. The filtrate was left for crystallization.

### Refinement

Aromatic H atoms were placed in calculated positions (C—H = 0.93 Å) and constrained to ride on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . Methylene H atoms were located in difference density maps and their coordinates and isotropic displacement parameters were refined freely.

### Figures




Fig. 1. Molecular structure of (I), with 50% probability displacement ellipsoids. H atoms are shown as circles of arbitrary radii.

## (I)

### Crystal data

$C_{28}H_{21}ClN_2$	$Z = 2$
$M_r = 420.92$	$F_{000} = 440$
Triclinic, $P\bar{1}$	$D_x = 1.291 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation
$a = 7.4880 (11) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 9.2711 (16) \text{ \AA}$	Cell parameters from 6987 reflections
$c = 16.049 (3) \text{ \AA}$	$\theta = 1.2\text{--}29.8^\circ$
$\alpha = 87.169 (13)^\circ$	$\mu = 0.19 \text{ mm}^{-1}$
$\beta = 76.704 (12)^\circ$	$T = 290 (2) \text{ K}$
$\gamma = 87.842 (13)^\circ$	Block, colorless
$V = 1082.6 (3) \text{ \AA}^3$	$0.3 \times 0.2 \times 0.2 \text{ mm}$

### Data collection

STOE IPDS-II diffractometer	2814 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.034$
Monochromator: graphite	$\theta_{\text{max}} = 26.0^\circ$
$T = 290(2) \text{ K}$	$\theta_{\text{min}} = 2.2^\circ$
Area detector – phi oscillation scans	$h = -9 \rightarrow 9$
Absorption correction: none	$k = -11 \rightarrow 10$
8769 measured reflections	$l = -19 \rightarrow 19$
4246 independent reflections	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.041$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.090$	$w = 1/[\sigma^2(F_o^2) + (0.0457P)^2]$
$S = 0.94$	where $P = (F_o^2 + 2F_c^2)/3$
4246 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
288 parameters	$\Delta\rho_{\text{max}} = 0.13 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.20 \text{ e \AA}^{-3}$
	Extinction correction: none

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.17061 (7)	0.52946 (7)	0.32233 (4)	0.0741 (2)
N1	0.60096 (17)	0.82368 (15)	0.59681 (9)	0.0412 (3)
N2	0.36337 (17)	0.74456 (15)	0.69583 (8)	0.0403 (3)
C1	0.2919 (2)	0.56492 (19)	0.54889 (12)	0.0456 (4)
H1	0.2804	0.5096	0.5997	0.055*
C2	0.4446 (2)	0.75675 (17)	0.61039 (10)	0.0393 (4)
C3	0.9378 (2)	0.9430 (2)	0.61528 (12)	0.0476 (4)
H3	0.9370	0.8959	0.5656	0.057*
C4	0.7878 (2)	0.93196 (18)	0.68467 (11)	0.0413 (4)
C5	0.4409 (2)	0.80553 (19)	0.83352 (11)	0.0419 (4)
C6	0.2487 (2)	0.5939 (2)	0.40748 (12)	0.0492 (5)
C7	0.2308 (2)	0.5117 (2)	0.48201 (12)	0.0491 (4)
H7	0.1784	0.4213	0.4874	0.059*
C8	0.6244 (2)	0.85566 (17)	0.67670 (10)	0.0391 (4)
C9	0.4790 (2)	0.80756 (18)	0.73880 (11)	0.0398 (4)
C10	0.1795 (2)	0.55784 (19)	0.79077 (11)	0.0435 (4)
C11	0.3703 (2)	0.69955 (18)	0.54181 (11)	0.0399 (4)
C12	0.3880 (2)	0.77975 (19)	0.46497 (11)	0.0475 (4)
H12	0.4418	0.8696	0.4587	0.057*
C13	0.5348 (3)	0.7078 (2)	0.87682 (12)	0.0567 (5)
H13	0.6178	0.6416	0.8463	0.068*
C14	0.3263 (3)	0.7273 (2)	0.39788 (12)	0.0530 (5)
H14	0.3373	0.7817	0.3468	0.064*
C15	1.0919 (3)	1.0925 (2)	0.69172 (14)	0.0607 (5)
H15	1.1921	1.1472	0.6939	0.073*
C16	0.1802 (2)	0.6943 (2)	0.73629 (12)	0.0440 (4)
H16A	0.114 (2)	0.6806 (18)	0.6900 (11)	0.046 (5)*
H16B	0.114 (2)	0.7692 (19)	0.7710 (11)	0.046 (5)*
C17	1.0877 (2)	1.0226 (2)	0.61875 (13)	0.0550 (5)
H17	1.1863	1.0290	0.5715	0.066*
C18	0.0320 (3)	0.5323 (2)	0.85851 (13)	0.0625 (5)
H18	-0.0639	0.6003	0.8704	0.075*
C19	0.1645 (4)	0.3071 (3)	0.89254 (17)	0.0854 (8)

## supplementary materials

---

H19	0.1596	0.2226	0.9267	0.102*
C20	0.7963 (2)	1.0024 (2)	0.75824 (12)	0.0505 (4)
H20	0.6988	0.9963	0.8060	0.061*
C21	0.3165 (3)	0.9005 (2)	0.88092 (12)	0.0577 (5)
H21	0.2503	0.9662	0.8533	0.069*
C22	0.2891 (3)	0.8988 (3)	0.96904 (13)	0.0699 (6)
H22	0.2046	0.9633	1.0003	0.084*
C23	0.9463 (3)	1.0807 (2)	0.76139 (13)	0.0604 (5)
H23	0.9495	1.1263	0.8113	0.072*
C24	0.3199 (3)	0.4554 (2)	0.77506 (13)	0.0557 (5)
H24	0.4208	0.4706	0.7298	0.067*
C25	0.3117 (4)	0.3305 (3)	0.82608 (17)	0.0747 (6)
H25	0.4072	0.2621	0.8150	0.090*
C26	0.3851 (3)	0.8033 (3)	1.01068 (13)	0.0680 (6)
H26	0.3676	0.8037	1.0699	0.082*
C27	0.5065 (3)	0.7074 (3)	0.96484 (14)	0.0688 (6)
H27	0.5708	0.6412	0.9931	0.083*
C28	0.0251 (4)	0.4068 (3)	0.90890 (15)	0.0820 (7)
H28	-0.0755	0.3905	0.9542	0.098*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0667 (3)	0.1008 (5)	0.0631 (3)	-0.0114 (3)	-0.0247 (3)	-0.0313 (3)
N1	0.0440 (7)	0.0441 (8)	0.0370 (8)	-0.0063 (6)	-0.0110 (6)	-0.0032 (6)
N2	0.0392 (7)	0.0453 (8)	0.0374 (8)	-0.0061 (6)	-0.0101 (6)	-0.0014 (6)
C1	0.0421 (9)	0.0474 (11)	0.0493 (10)	-0.0038 (8)	-0.0142 (8)	-0.0024 (8)
C2	0.0399 (8)	0.0402 (10)	0.0396 (9)	-0.0009 (7)	-0.0126 (7)	-0.0028 (7)
C3	0.0473 (9)	0.0531 (11)	0.0429 (10)	-0.0051 (8)	-0.0102 (8)	-0.0057 (8)
C4	0.0445 (9)	0.0407 (10)	0.0407 (9)	-0.0038 (7)	-0.0136 (7)	-0.0021 (7)
C5	0.0408 (8)	0.0473 (10)	0.0383 (9)	-0.0095 (7)	-0.0096 (7)	0.0000 (8)
C6	0.0400 (9)	0.0621 (13)	0.0497 (11)	0.0020 (8)	-0.0156 (8)	-0.0205 (9)
C7	0.0400 (9)	0.0491 (11)	0.0598 (12)	-0.0063 (8)	-0.0124 (8)	-0.0105 (9)
C8	0.0427 (8)	0.0372 (9)	0.0384 (9)	-0.0002 (7)	-0.0109 (7)	-0.0040 (7)
C9	0.0401 (8)	0.0419 (9)	0.0394 (9)	-0.0012 (7)	-0.0128 (7)	-0.0033 (7)
C10	0.0417 (9)	0.0516 (11)	0.0402 (10)	-0.0117 (8)	-0.0136 (8)	-0.0035 (8)
C11	0.0366 (8)	0.0437 (10)	0.0404 (9)	0.0010 (7)	-0.0100 (7)	-0.0064 (8)
C12	0.0558 (10)	0.0434 (11)	0.0443 (10)	-0.0058 (8)	-0.0122 (8)	-0.0055 (8)
C13	0.0580 (11)	0.0655 (13)	0.0497 (12)	0.0048 (9)	-0.0196 (9)	-0.0046 (10)
C14	0.0636 (11)	0.0563 (12)	0.0419 (10)	0.0011 (9)	-0.0174 (9)	-0.0059 (9)
C15	0.0520 (10)	0.0654 (13)	0.0692 (14)	-0.0171 (9)	-0.0200 (10)	-0.0062 (11)
C16	0.0337 (8)	0.0529 (11)	0.0454 (10)	-0.0023 (8)	-0.0083 (8)	-0.0041 (9)
C17	0.0450 (10)	0.0636 (13)	0.0551 (12)	-0.0105 (9)	-0.0080 (9)	-0.0002 (10)
C18	0.0561 (11)	0.0763 (15)	0.0532 (12)	-0.0186 (10)	-0.0061 (9)	-0.0019 (11)
C19	0.140 (2)	0.0633 (16)	0.0644 (16)	-0.0376 (17)	-0.0452 (17)	0.0139 (13)
C20	0.0515 (10)	0.0560 (12)	0.0443 (10)	-0.0104 (8)	-0.0088 (8)	-0.0082 (9)
C21	0.0628 (11)	0.0599 (13)	0.0483 (11)	0.0059 (10)	-0.0092 (9)	-0.0034 (9)
C22	0.0746 (14)	0.0808 (16)	0.0480 (13)	0.0019 (12)	0.0005 (11)	-0.0138 (11)

C23	0.0634 (12)	0.0680 (13)	0.0554 (12)	-0.0141 (10)	-0.0206 (10)	-0.0166 (10)
C24	0.0565 (11)	0.0577 (13)	0.0543 (12)	-0.0039 (9)	-0.0161 (9)	0.0028 (10)
C25	0.0997 (17)	0.0578 (14)	0.0771 (16)	-0.0012 (12)	-0.0425 (14)	0.0001 (12)
C26	0.0800 (14)	0.0880 (17)	0.0359 (11)	-0.0195 (12)	-0.0106 (10)	0.0004 (11)
C27	0.0821 (14)	0.0792 (16)	0.0513 (13)	-0.0035 (12)	-0.0298 (11)	0.0076 (11)
C28	0.0988 (18)	0.092 (2)	0.0522 (14)	-0.0498 (16)	-0.0062 (12)	0.0070 (13)

*Geometric parameters (Å, °)*

C11—C6	1.7436 (17)	C13—C27	1.379 (3)
N1—C2	1.314 (2)	C13—H13	0.9300
N1—C8	1.381 (2)	C14—H14	0.9300
N2—C2	1.367 (2)	C15—C17	1.373 (3)
N2—C9	1.385 (2)	C15—C23	1.374 (3)
N2—C16	1.459 (2)	C15—H15	0.9300
C1—C7	1.378 (2)	C16—H16A	1.000 (18)
C1—C11	1.388 (2)	C16—H16B	0.960 (17)
C1—H1	0.9300	C17—H17	0.9300
C2—C11	1.471 (2)	C18—C28	1.379 (3)
C3—C17	1.380 (2)	C18—H18	0.9300
C3—C4	1.391 (2)	C19—C28	1.356 (4)
C3—H3	0.9300	C19—C25	1.362 (4)
C4—C20	1.392 (2)	C19—H19	0.9300
C4—C8	1.470 (2)	C20—C23	1.372 (3)
C5—C21	1.378 (2)	C20—H20	0.9300
C5—C13	1.383 (3)	C21—C22	1.381 (3)
C5—C9	1.480 (2)	C21—H21	0.9300
C6—C7	1.369 (3)	C22—C26	1.364 (3)
C6—C14	1.373 (3)	C22—H22	0.9300
C7—H7	0.9300	C23—H23	0.9300
C8—C9	1.368 (2)	C24—C25	1.380 (3)
C10—C24	1.377 (2)	C24—H24	0.9300
C10—C18	1.378 (3)	C25—H25	0.9300
C10—C16	1.501 (3)	C26—C27	1.362 (3)
C11—C12	1.390 (2)	C26—H26	0.9300
C12—C14	1.381 (2)	C27—H27	0.9300
C12—H12	0.9300	C28—H28	0.9300
C2—N1—C8	105.97 (13)	C17—C15—C23	119.26 (17)
C2—N2—C9	106.82 (13)	C17—C15—H15	120.4
C2—N2—C16	128.26 (14)	C23—C15—H15	120.4
C9—N2—C16	124.44 (14)	N2—C16—C10	114.03 (14)
C7—C1—C11	121.22 (18)	N2—C16—H16A	107.8 (10)
C7—C1—H1	119.4	C10—C16—H16A	110.3 (10)
C11—C1—H1	119.4	N2—C16—H16B	108.6 (10)
N1—C2—N2	111.47 (14)	C10—C16—H16B	109.0 (10)
N1—C2—C11	123.73 (15)	H16A—C16—H16B	106.7 (13)
N2—C2—C11	124.79 (14)	C15—C17—C3	120.22 (17)
C17—C3—C4	121.31 (16)	C15—C17—H17	119.9
C17—C3—H3	119.3	C3—C17—H17	119.9

## supplementary materials

---

C4—C3—H3	119.3	C10—C18—C28	120.7 (2)
C3—C4—C20	117.33 (16)	C10—C18—H18	119.7
C3—C4—C8	119.95 (15)	C28—C18—H18	119.7
C20—C4—C8	122.61 (15)	C28—C19—C25	120.0 (2)
C21—C5—C13	118.12 (17)	C28—C19—H19	120.0
C21—C5—C9	122.19 (16)	C25—C19—H19	120.0
C13—C5—C9	119.68 (15)	C23—C20—C4	121.03 (17)
C7—C6—C14	121.57 (17)	C23—C20—H20	119.5
C7—C6—C11	119.75 (15)	C4—C20—H20	119.5
C14—C6—C11	118.69 (16)	C5—C21—C22	120.6 (2)
C6—C7—C1	118.92 (17)	C5—C21—H21	119.7
C6—C7—H7	120.5	C22—C21—H21	119.7
C1—C7—H7	120.5	C26—C22—C21	120.6 (2)
C9—C8—N1	109.98 (14)	C26—C22—H22	119.7
C9—C8—C4	129.88 (15)	C21—C22—H22	119.7
N1—C8—C4	120.14 (14)	C20—C23—C15	120.83 (18)
C8—C9—N2	105.76 (14)	C20—C23—H23	119.6
C8—C9—C5	132.19 (15)	C15—C23—H23	119.6
N2—C9—C5	121.91 (14)	C10—C24—C25	120.4 (2)
C24—C10—C18	118.34 (19)	C10—C24—H24	119.8
C24—C10—C16	122.53 (16)	C25—C24—H24	119.8
C18—C10—C16	119.12 (17)	C19—C25—C24	120.3 (2)
C1—C11—C12	118.42 (16)	C19—C25—H25	119.8
C1—C11—C2	122.62 (16)	C24—C25—H25	119.8
C12—C11—C2	118.87 (15)	C27—C26—C22	119.51 (19)
C14—C12—C11	120.65 (17)	C27—C26—H26	120.2
C14—C12—H12	119.7	C22—C26—H26	120.2
C11—C12—H12	119.7	C26—C27—C13	120.5 (2)
C27—C13—C5	120.73 (19)	C26—C27—H27	119.8
C27—C13—H13	119.6	C13—C27—H27	119.8
C5—C13—H13	119.6	C19—C28—C18	120.2 (2)
C6—C14—C12	119.22 (18)	C19—C28—H28	119.9
C6—C14—H14	120.4	C18—C28—H28	119.9
C12—C14—H14	120.4		
C8—N1—C2—N2	0.58 (18)	N1—C2—C11—C12	-39.5 (2)
C8—N1—C2—C11	-178.04 (15)	N2—C2—C11—C12	142.03 (17)
C9—N2—C2—N1	-0.62 (19)	C1—C11—C12—C14	0.9 (2)
C16—N2—C2—N1	171.59 (15)	C2—C11—C12—C14	177.51 (15)
C9—N2—C2—C11	177.97 (15)	C21—C5—C13—C27	-1.1 (3)
C16—N2—C2—C11	-9.8 (3)	C9—C5—C13—C27	177.59 (18)
C17—C3—C4—C20	-1.0 (3)	C7—C6—C14—C12	0.0 (3)
C17—C3—C4—C8	175.34 (17)	C11—C6—C14—C12	179.73 (13)
C14—C6—C7—C1	0.3 (3)	C11—C12—C14—C6	-0.6 (3)
C11—C6—C7—C1	-179.45 (12)	C2—N2—C16—C10	115.85 (19)
C11—C1—C7—C6	0.1 (2)	C9—N2—C16—C10	-73.2 (2)
C2—N1—C8—C9	-0.31 (18)	C24—C10—C16—N2	-28.4 (2)
C2—N1—C8—C4	-179.85 (15)	C18—C10—C16—N2	152.54 (16)
C3—C4—C8—C9	165.88 (18)	C23—C15—C17—C3	0.8 (3)
C20—C4—C8—C9	-17.9 (3)	C4—C3—C17—C15	0.3 (3)

C3—C4—C8—N1	-14.7 (2)	C24—C10—C18—C28	-0.3 (3)
C20—C4—C8—N1	161.51 (16)	C16—C10—C18—C28	178.75 (18)
N1—C8—C9—N2	-0.06 (18)	C3—C4—C20—C23	0.7 (3)
C4—C8—C9—N2	179.42 (16)	C8—C4—C20—C23	-175.60 (17)
N1—C8—C9—C5	175.61 (17)	C13—C5—C21—C22	0.9 (3)
C4—C8—C9—C5	-4.9 (3)	C9—C5—C21—C22	-177.70 (18)
C2—N2—C9—C8	0.40 (18)	C5—C21—C22—C26	0.1 (3)
C16—N2—C9—C8	-172.20 (15)	C4—C20—C23—C15	0.4 (3)
C2—N2—C9—C5	-175.82 (15)	C17—C15—C23—C20	-1.2 (3)
C16—N2—C9—C5	11.6 (2)	C18—C10—C24—C25	0.1 (3)
C21—C5—C9—C8	105.1 (2)	C16—C10—C24—C25	-178.92 (18)
C13—C5—C9—C8	-73.6 (3)	C28—C19—C25—C24	-0.1 (3)
C21—C5—C9—N2	-79.8 (2)	C10—C24—C25—C19	0.1 (3)
C13—C5—C9—N2	101.5 (2)	C21—C22—C26—C27	-1.1 (3)
C7—C1—C11—C12	-0.7 (2)	C22—C26—C27—C13	0.9 (3)
C7—C1—C11—C2	-177.10 (14)	C5—C13—C27—C26	0.2 (3)
N1—C2—C11—C1	136.88 (17)	C25—C19—C28—C18	-0.1 (4)
N2—C2—C11—C1	-41.5 (2)	C10—C18—C28—C19	0.3 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C3—H3...N1	0.93	2.56	2.874 (2)	100

Fig. 1

