

## 3,3-Dimethyl-1,2,3,4,6,11-hexahydrobenzo[*d*]naphtho[2,3-*b*]furan-1,6,11-trione

Huayou Hu

Department of Chemistry and Chemical Engineering, Southeast University, Nanjing, People's Republic of China  
 Correspondence e-mail: njuhhy@hotmail.com


Received 11 July 2008; accepted 11 July 2008

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.048;  $wR$  factor = 0.137; data-to-parameter ratio = 13.4.

In the title compound,  $\text{C}_{18}\text{H}_{14}\text{O}_4$ , the cyclohexene ring adopts a sofa conformation. In the crystalline state, the molecules are linked into a chain by weak intermolecular  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

### Related literature

For related literature, see: Correa & Romo (1966); Greve & Friedrichsen (2000); Hirai *et al.* (1999); Hu *et al.* (2005); Ito *et al.* (2000). For related structures, see: Goldstein *et al.* (1975); Park *et al.* (1992).



### Experimental

#### Crystal data

$\text{C}_{18}\text{H}_{14}\text{O}_4$	$\gamma = 78.18(3)^\circ$
$M_r = 294.29$	$V = 695.2(2)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 5.8080(12)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 6.7510(14)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$c = 18.332(4)\text{ \AA}$	$T = 293(2)\text{ K}$
$\alpha = 89.82(3)^\circ$	$0.25 \times 0.18 \times 0.16\text{ mm}$
$\beta = 81.32(3)^\circ$	

### Data collection

Bruker APEX CCD diffractometer	2958 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 2000)	2676 independent reflections
$T_{\min} = 0.976$ , $T_{\max} = 0.984$	1845 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.018$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$	199 parameters
$wR(F^2) = 0.137$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\max} = 0.16\text{ e \AA}^{-3}$
2676 reflections	$\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

**Table 1**  
 Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C16—H16A $\cdots$ O4 <sup>i</sup>	0.93	2.54	3.177 (3)	126

Symmetry code: (i)  $x - 1, y + 1, z$ .

Data collection: *SMART* (Bruker, 2000); cell refinement: *SMART*; data reduction: *SAINT* (Bruker, 2000); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The author thanks the Program for Young Excellent Talents in Southeast University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2746).

### References

- Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Correa, J. & Romo, J. (1966). *Tetrahedron*, **22**, 685–691.
- Goldstein, P. (1975). *Acta Cryst. B* **31**, 2086–2097.
- Greve, S. & Friedrichsen, W. (2000). *Prog. Heterocycl. Chem.* **12**, 134–160.
- Hirai, K. I., Koyama, J., Pan, J., Simamura, E., Shimada, H., Yamori, T., Sato, S., Tagahara, K. & Tsuruo, T. (1999). *Cancer Detect. Prev.* **23**, 539–550.
- Hu, H.-Y., Ye, Z., Wang, L. & Xu, J.-H. (2005). *Synthesis*, pp. 1605–1610.
- Ito, C., Katsuno, S., Kondo, Y., Tan, H. T.-W. & Furukawa, H. (2000). *Chem. Pharm. Bull.* **48**, 339–343.
- Park, I. Y., Kim, B. K. & Kim, Y. B. (1992). *Arch. Pharm. Res.* **15**, 52–57.
- Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2008). E64, o1534 [doi:10.1107/S1600536808021600]

## 3,3-Dimethyl-1,2,3,4,6,11-hexahydrobenzo[*d*]naphtho[2,3-*b*]furan-1,6,11-trione

Huayou Hu

### S1. Comment

Furan derivatives and annulated furan derivatives widely occur in nature. Many of these naturally occurring furan and annulated furan derivatives and their unnatural analogs have a wide range of biological activity and are important precursors for the synthesis of natural products (Greve & Friedrichsen, 2000). Especially, naphtho[2,3-*b*]furan-4,9-dione derivatives as represented by avicequinones (Ito *et al.*, 2000) and maturinones (Correa & Romo, 1966) have shown a diversity of biological activities of medical importance, such as anticancer, antibacterial and anti-inflammatory activity (Hirai *et al.*, 1999). Recently, we had reported an one-pot synthesis method for naphtho[2,3-*b*]furan-4,9-dione derivatives by reacting 2,3-dichloro-1,4-naphthoquinone with 1,3-dicarbonyl compounds (Hu *et al.*, 2005).


The title compound, C<sub>18</sub>H<sub>14</sub>O<sub>4</sub>, is a naphtho[2,3-*b*]furan-4,9-dione derivative. It is formed as one product from refluxing 2,3-dichloro-1,4-naphthoquinone, K<sub>2</sub>CO<sub>3</sub> and 5,5-dimethylcyclohexane-1,3-dione in MeCN for 6 h. In the crystalline state, the molecules are linked to a one-dimensional chain by intermolecular weak C—H···O hydrogen bonds (Table 1).

### S2. Experimental


A mixture of 2,3-dichloro-1,4-naphthoquinone (0.227 g, 1.0 mmol), K<sub>2</sub>CO<sub>3</sub> (0.345 g, 2.5 mmol) and 5,5-dimethylcyclohexane-1,3-dione (0.154 g, 1.1 mmol) in MeCN (15 ml) was stirred at reflux temperature for 6 h. The reaction mixture was poured into H<sub>2</sub>O (150 ml) and filtered. The collected solid product was separated by silica gel column chromatography [petroleum ether - EtOAc (10:1)] yield the title compound (0.118 g, 40%) as a yellow solid. Single crystals suitable for X-ray crystallographic analysis were grown by slow evaporation of solvent from petroleum ether (b.p. 333–363 K)-ethyl acetate (3/1 *v/v*).

### S3. Refinement

H atoms were positioned geometrically and refined using a riding model with C—H = 0.93–0.97 Å and with *U*<sub>iso</sub>(H) = 1.2 times *U*<sub>eq</sub>(C) or *U*<sub>iso</sub>(H) = 1.5 times *U*<sub>eq</sub>(C<sub>methyl</sub>).

**Figure 1**

The molecular structure of the title compound with 30% displacement ellipsoids.

**Figure 2**

Hydrogen-bonding of the title compound.

### 3,3-Dimethyl-1,2,3,4,6,11-hexahydrobenzo[*d*]naphtho[2,3-*b*]furan- 1,6,11-trione

#### Crystal data

$C_{18}H_{14}O_4$   
 $M_r = 294.29$   
Triclinic,  $P\bar{1}$   
Hall symbol: -P 1  
 $a = 5.8080 (12)$  Å  
 $b = 6.7510 (14)$  Å  
 $c = 18.332 (4)$  Å  
 $\alpha = 89.82 (3)^\circ$   
 $\beta = 81.32 (3)^\circ$   
 $\gamma = 78.18 (3)^\circ$   
 $V = 695.2 (2)$  Å<sup>3</sup>

$Z = 2$   
 $F(000) = 308$   
 $D_x = 1.406 \text{ Mg m}^{-3}$   
Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å  
Cell parameters from 810 reflections  
 $\theta = 2.5\text{--}28.0^\circ$   
 $\mu = 0.10 \text{ mm}^{-1}$   
 $T = 293$  K  
Block, yellow  
 $0.25 \times 0.18 \times 0.16$  mm

#### Data collection

Bruker APEX CCD  
diffractometer  
Radiation source: fine-focus sealed tube

Graphite monochromator  
 $\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
 (SADABS; Sheldrick, 2000)  
 $T_{\min} = 0.976$ ,  $T_{\max} = 0.984$   
 2958 measured reflections  
 2676 independent reflections  
 1845 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$   
 $\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 1.1^\circ$   
 $h = 0 \rightarrow 6$   
 $k = -8 \rightarrow 8$   
 $l = -21 \rightarrow 21$

### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.137$   
 $S = 1.08$   
 2676 reflections  
 199 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0661P)^2 + 0.1193P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	-0.1220 (3)	1.1681 (3)	0.34222 (12)	0.0843 (7)
O2	0.6416 (2)	0.9262 (2)	0.22339 (7)	0.0410 (4)
O3	0.0036 (3)	1.4623 (2)	0.22564 (10)	0.0632 (5)
O4	0.9115 (3)	1.1186 (2)	0.11462 (10)	0.0632 (5)
C1	0.2866 (5)	0.5132 (4)	0.41299 (15)	0.0674 (7)
H1A	0.4384	0.4486	0.4254	0.081*
H1B	0.2448	0.4330	0.3759	0.081*
H1C	0.1680	0.5262	0.4562	0.081*
C2	0.3705 (4)	0.8498 (4)	0.44217 (13)	0.0619 (7)
H2A	0.5219	0.7838	0.4545	0.074*
H2B	0.2523	0.8649	0.4856	0.074*
H2C	0.3812	0.9809	0.4232	0.074*
C3	0.3004 (4)	0.7223 (3)	0.38363 (12)	0.0463 (5)
C4	0.0571 (4)	0.8275 (4)	0.36332 (13)	0.0531 (6)
H4A	0.0099	0.7383	0.3296	0.064*
H4B	-0.0597	0.8444	0.4078	0.064*
C5	0.0481 (3)	1.0297 (4)	0.32872 (12)	0.0498 (6)
C6	0.2637 (3)	1.0420 (3)	0.27647 (11)	0.0406 (5)
C7	0.4580 (3)	0.8876 (3)	0.27275 (11)	0.0381 (5)

C8	0.4861 (3)	0.6996 (3)	0.31386 (11)	0.0413 (5)
H8A	0.6445	0.6671	0.3273	0.050*
H8B	0.4679	0.5893	0.2829	0.050*
C9	0.5599 (3)	1.1101 (3)	0.19527 (11)	0.0394 (5)
C10	0.3304 (3)	1.1890 (3)	0.22546 (11)	0.0394 (5)
C11	0.2091 (3)	1.3861 (3)	0.20166 (11)	0.0419 (5)
C12	0.3557 (3)	1.4922 (3)	0.14706 (11)	0.0407 (5)
C13	0.5943 (3)	1.4062 (3)	0.11906 (11)	0.0405 (5)
C14	0.7101 (3)	1.2020 (3)	0.14072 (12)	0.0430 (5)
C15	0.2569 (4)	1.6823 (3)	0.12476 (12)	0.0489 (6)
H15A	0.0991	1.7403	0.1428	0.059*
C16	0.3895 (4)	1.7868 (3)	0.07620 (13)	0.0556 (6)
H16A	0.3205	1.9141	0.0612	0.067*
C17	0.6245 (4)	1.7035 (3)	0.04972 (13)	0.0568 (6)
H17A	0.7148	1.7752	0.0176	0.068*
C18	0.7246 (4)	1.5134 (3)	0.07112 (12)	0.0497 (5)
H18A	0.8827	1.4568	0.0529	0.060*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0402 (10)	0.0837 (13)	0.1077 (16)	0.0104 (9)	0.0245 (10)	0.0054 (11)
O2	0.0272 (7)	0.0433 (8)	0.0473 (8)	-0.0004 (6)	0.0012 (6)	0.0034 (6)
O3	0.0309 (8)	0.0669 (11)	0.0782 (12)	0.0111 (7)	0.0054 (8)	0.0014 (8)
O4	0.0357 (8)	0.0517 (9)	0.0847 (12)	0.0102 (7)	0.0190 (8)	0.0122 (8)
C1	0.0565 (15)	0.0743 (18)	0.0711 (17)	-0.0243 (13)	0.0057 (13)	0.0130 (13)
C2	0.0604 (15)	0.0814 (18)	0.0472 (13)	-0.0245 (13)	-0.0052 (11)	-0.0070 (12)
C3	0.0344 (11)	0.0585 (13)	0.0481 (12)	-0.0174 (9)	-0.0019 (9)	-0.0011 (10)
C4	0.0285 (11)	0.0664 (15)	0.0640 (15)	-0.0170 (10)	0.0045 (10)	-0.0065 (11)
C5	0.0266 (11)	0.0655 (14)	0.0547 (13)	-0.0084 (10)	0.0004 (9)	-0.0064 (11)
C6	0.0267 (10)	0.0484 (12)	0.0454 (12)	-0.0058 (8)	-0.0037 (8)	-0.0069 (9)
C7	0.0270 (10)	0.0463 (11)	0.0404 (11)	-0.0089 (8)	-0.0017 (8)	-0.0037 (9)
C8	0.0311 (10)	0.0475 (12)	0.0456 (12)	-0.0102 (8)	-0.0043 (9)	-0.0014 (9)
C9	0.0290 (10)	0.0403 (11)	0.0444 (11)	0.0014 (8)	-0.0032 (8)	-0.0013 (9)
C10	0.0268 (10)	0.0455 (11)	0.0426 (11)	-0.0011 (8)	-0.0034 (8)	-0.0079 (9)
C11	0.0277 (10)	0.0472 (12)	0.0454 (12)	0.0046 (8)	-0.0052 (8)	-0.0104 (9)
C12	0.0334 (10)	0.0435 (11)	0.0416 (11)	0.0033 (8)	-0.0095 (9)	-0.0073 (9)
C13	0.0336 (10)	0.0414 (11)	0.0416 (11)	0.0030 (8)	-0.0047 (8)	-0.0030 (8)
C14	0.0301 (10)	0.0427 (11)	0.0501 (12)	0.0008 (8)	0.0013 (9)	-0.0024 (9)
C15	0.0392 (11)	0.0472 (12)	0.0535 (13)	0.0085 (9)	-0.0093 (10)	-0.0057 (10)
C16	0.0558 (14)	0.0440 (12)	0.0605 (15)	0.0064 (10)	-0.0111 (12)	0.0029 (10)
C17	0.0540 (14)	0.0496 (13)	0.0616 (15)	-0.0021 (11)	-0.0041 (12)	0.0078 (11)
C18	0.0384 (11)	0.0490 (12)	0.0548 (13)	0.0015 (9)	0.0006 (10)	0.0039 (10)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

O1—C5	1.208 (3)	C6—C10	1.432 (3)
O2—C7	1.358 (2)	C7—C8	1.467 (3)

O2—C9	1.364 (2)	C8—H8A	0.9700
O3—C11	1.214 (2)	C8—H8B	0.9700
O4—C14	1.216 (2)	C9—C10	1.362 (3)
C1—C3	1.522 (3)	C9—C14	1.449 (3)
C1—H1A	0.9600	C10—C11	1.470 (3)
C1—H1B	0.9600	C11—C12	1.492 (3)
C1—H1C	0.9600	C12—C15	1.382 (3)
C2—C3	1.530 (3)	C12—C13	1.406 (3)
C2—H2A	0.9600	C13—C18	1.374 (3)
C2—H2B	0.9600	C13—C14	1.486 (3)
C2—H2C	0.9600	C15—C16	1.377 (3)
C3—C8	1.530 (3)	C15—H15A	0.9300
C3—C4	1.545 (3)	C16—C17	1.379 (3)
C4—C5	1.498 (3)	C16—H16A	0.9300
C4—H4A	0.9700	C17—C18	1.378 (3)
C4—H4B	0.9700	C17—H17A	0.9300
C5—C6	1.474 (3)	C18—H18A	0.9300
C6—C7	1.364 (3)		
C7—O2—C9	105.92 (14)	C7—C8—H8A	109.6
C3—C1—H1A	109.5	C3—C8—H8A	109.6
C3—C1—H1B	109.5	C7—C8—H8B	109.6
H1A—C1—H1B	109.5	C3—C8—H8B	109.6
C3—C1—H1C	109.5	H8A—C8—H8B	108.1
H1A—C1—H1C	109.5	C10—C9—O2	111.59 (18)
H1B—C1—H1C	109.5	C10—C9—C14	126.95 (19)
C3—C2—H2A	109.5	O2—C9—C14	121.46 (16)
C3—C2—H2B	109.5	C9—C10—C6	105.23 (18)
H2A—C2—H2B	109.5	C9—C10—C11	119.84 (19)
C3—C2—H2C	109.5	C6—C10—C11	134.92 (17)
H2A—C2—H2C	109.5	O3—C11—C10	122.8 (2)
H2B—C2—H2C	109.5	O3—C11—C12	121.16 (19)
C1—C3—C8	109.05 (18)	C10—C11—C12	116.04 (16)
C1—C3—C2	109.7 (2)	C15—C12—C13	118.9 (2)
C8—C3—C2	109.88 (17)	C15—C12—C11	119.26 (18)
C1—C3—C4	109.92 (18)	C13—C12—C11	121.79 (18)
C8—C3—C4	108.39 (18)	C18—C13—C12	119.57 (19)
C2—C3—C4	109.86 (19)	C18—C13—C14	119.02 (18)
C5—C4—C3	115.97 (17)	C12—C13—C14	121.41 (19)
C5—C4—H4A	108.3	O4—C14—C9	122.99 (19)
C3—C4—H4A	108.3	O4—C14—C13	123.17 (19)
C5—C4—H4B	108.3	C9—C14—C13	113.84 (16)
C3—C4—H4B	108.3	C16—C15—C12	120.7 (2)
H4A—C4—H4B	107.4	C16—C15—H15A	119.6
O1—C5—C6	123.7 (2)	C12—C15—H15A	119.6
O1—C5—C4	122.6 (2)	C15—C16—C17	120.2 (2)
C6—C5—C4	113.70 (18)	C15—C16—H16A	119.9
C7—C6—C10	106.29 (17)	C17—C16—H16A	119.9

C7—C6—C5	118.9 (2)	C18—C17—C16	119.6 (2)
C10—C6—C5	134.80 (19)	C18—C17—H17A	120.2
O2—C7—C6	110.97 (18)	C16—C17—H17A	120.2
O2—C7—C8	120.29 (16)	C13—C18—C17	120.9 (2)
C6—C7—C8	128.74 (18)	C13—C18—H18A	119.5
C7—C8—C3	110.45 (17)	C17—C18—H18A	119.5
C1—C3—C4—C5	-175.99 (19)	C5—C6—C10—C11	-2.8 (4)
C8—C3—C4—C5	-56.9 (2)	C9—C10—C11—O3	178.7 (2)
C2—C3—C4—C5	63.2 (2)	C6—C10—C11—O3	-0.3 (4)
C3—C4—C5—O1	-142.1 (2)	C9—C10—C11—C12	-2.9 (3)
C3—C4—C5—C6	38.6 (3)	C6—C10—C11—C12	178.1 (2)
O1—C5—C6—C7	171.1 (2)	O3—C11—C12—C15	1.5 (3)
C4—C5—C6—C7	-9.6 (3)	C10—C11—C12—C15	-177.03 (17)
O1—C5—C6—C10	-6.8 (4)	O3—C11—C12—C13	179.7 (2)
C4—C5—C6—C10	172.5 (2)	C10—C11—C12—C13	1.2 (3)
C9—O2—C7—C6	-0.1 (2)	C15—C12—C13—C18	0.8 (3)
C9—O2—C7—C8	-179.77 (17)	C11—C12—C13—C18	-177.42 (19)
C10—C6—C7—O2	0.0 (2)	C15—C12—C13—C14	-179.75 (18)
C5—C6—C7—O2	-178.41 (17)	C11—C12—C13—C14	2.0 (3)
C10—C6—C7—C8	179.67 (19)	C10—C9—C14—O4	-177.5 (2)
C5—C6—C7—C8	1.2 (3)	O2—C9—C14—O4	3.3 (3)
O2—C7—C8—C3	159.23 (17)	C10—C9—C14—C13	1.8 (3)
C6—C7—C8—C3	-20.4 (3)	O2—C9—C14—C13	-177.40 (17)
C1—C3—C8—C7	164.09 (18)	C18—C13—C14—O4	-4.7 (3)
C2—C3—C8—C7	-75.6 (2)	C12—C13—C14—O4	175.9 (2)
C4—C3—C8—C7	44.4 (2)	C18—C13—C14—C9	176.00 (19)
C7—O2—C9—C10	0.1 (2)	C12—C13—C14—C9	-3.4 (3)
C7—O2—C9—C14	179.43 (18)	C13—C12—C15—C16	-0.3 (3)
O2—C9—C10—C6	-0.1 (2)	C11—C12—C15—C16	178.01 (19)
C14—C9—C10—C6	-179.36 (19)	C12—C15—C16—C17	-0.7 (3)
O2—C9—C10—C11	-179.37 (16)	C15—C16—C17—C18	1.2 (4)
C14—C9—C10—C11	1.4 (3)	C12—C13—C18—C17	-0.4 (3)
C7—C6—C10—C9	0.0 (2)	C14—C13—C18—C17	-179.8 (2)
C5—C6—C10—C9	178.1 (2)	C16—C17—C18—C13	-0.6 (4)
C7—C6—C10—C11	179.2 (2)		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C16—H16A···O4 <sup>i</sup>	0.93	2.54	3.177 (3)	126

Symmetry code: (i)  $x-1, y+1, z$ .