

# N'-(5-Bromo-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide dihydrate

Abeer A. Alhadi, Hapipah M. Ali, Subramaniam Puvaneswary, Ward T. Robinson and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 16 July 2008; accepted 19 July 2008

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$ ;  $R$  factor = 0.065;  $wR$  factor = 0.155; data-to-parameter ratio = 14.2.

The title compound,  $\text{C}_{14}\text{H}_{11}\text{BrN}_2\text{O}_5 \cdot 2\text{H}_2\text{O}$ , crystallizes as hydrogen-bonded sheets. The 2-hydroxy group on the benzylidene group forms an intramolecular hydrogen bond to the N atom of the  $\text{C}=\text{N}$  double bond. The amino N atom is a hydrogen-bond donor to a water molecule. The hydroxy group on the benzohydrazide group is a hydrogen-bond donor to one acceptor site, whereas each water molecule is a hydrogen-bond donor to two acceptor sites.

## Related literature

For the structure of a similar Schiff-base ligand, 5-bromo-salicylaldehyde benzoylhydrazone, see: Liu *et al.* (2006).



## Experimental

### Crystal data

$\text{C}_{14}\text{H}_{11}\text{BrN}_2\text{O}_5 \cdot 2\text{H}_2\text{O}$   
 $M_r = 403.19$   
Monoclinic,  $P2_1/c$   
 $a = 30.8424(8)\text{ \AA}$   
 $b = 3.7999(1)\text{ \AA}$   
 $c = 12.8484(4)\text{ \AA}$   
 $\beta = 90.280(2)^\circ$

$V = 1505.79(7)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 2.77\text{ mm}^{-1}$   
 $T = 100(2)\text{ K}$   
 $0.30 \times 0.03 \times 0.03\text{ mm}$

### Data collection

Bruker SMART APEX diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.658$ ,  $T_{\max} = 0.921$

9964 measured reflections  
3424 independent reflections  
2914 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.039$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.064$   
 $wR(F^2) = 0.155$   
 $S = 1.22$   
3424 reflections  
241 parameters  
10 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 1.08\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -1.82\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1o $\cdots$ N1	0.84 (1)	1.91 (5)	2.616 (6)	141 (7)
O3—H3o $\cdots$ O2w <sup>i</sup>	0.84 (1)	1.96 (4)	2.736 (6)	153 (7)
O4—H4o $\cdots$ O2w <sup>i</sup>	0.84 (1)	1.81 (3)	2.623 (8)	163 (9)
O5—H5o $\cdots$ O2 <sup>ii</sup>	0.84 (1)	1.93 (2)	2.764 (5)	171 (7)
O1w—H1w1 $\cdots$ O2 <sup>iii</sup>	0.84 (1)	1.98 (2)	2.812 (5)	170 (6)
O1w—H1w2 $\cdots$ O1 <sup>ii</sup>	0.84 (1)	2.09 (2)	2.914 (6)	167 (6)
O2w—H2w1 $\cdots$ O3 <sup>iv</sup>	0.84 (1)	2.13 (5)	2.845 (9)	142 (8)
O2w—H2w2 $\cdots$ O4 <sup>v</sup>	0.84 (1)	2.12 (4)	2.900 (8)	154 (8)

Symmetry codes: (i)  $-x, -y + 2, -z + 2$ ; (ii)  $x, -y + \frac{3}{2}, z + \frac{1}{2}$ ; (iii)  $x, -y + \frac{5}{2}, z + \frac{1}{2}$ ; (iv)  $-x, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (v)  $-x, -y + 3, -z + 2$ .

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

We thank the Science Fund (12-02-03-2031, 12-02-03-2051) and the University of Malaya (PJP) for supporting this study. We are grateful to the University of Malaya for the purchase of the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2751).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Liu, H.-Y., Wang, H.-Y., Gao, F., Lu, Z.-S. & Niu, D.-Z. (2006). *Acta Cryst. E62*, o4495–o4496.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2008). *publCIF*. In preparation.

# supporting information

*Acta Cryst.* (2008). E64, o1584 [doi:10.1107/S1600536808022708]

## N'-(5-Bromo-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide dihydrate

**Abeer A. Alhadi, Hapipah M. Ali, Subramaniam Puvaneswary, Ward T. Robinson and Seik Weng Ng**

### S1. Comment

This study extends the structural study on the Schiff base, 5-bromosalicylaldehyde benzoylhydrazone (Liu *et al.*, 2006) as the title compound (Scheme I, Fig. 1) has several hydroxy groups on one of the aromatic rings. The compound crystallizes with two lattice water molecules. Hydrogen bonding interactions (Table 1) give rise to a layer motif.


### S2. Experimental

3,4,5-Trihydroxybenzoylhydrazide (0.65 g, 3.5 mmol) and 5-bromo-2-hydroxybenzaldehyde (0.70 g, 3.5 mmol) were heated for 12 h in ethanol. The solvent was removed and the product recrystallized from ethanol.

### S3. Refinement

Carbon and nitrogen-bound H-atoms were placed in calculated positions (C—H 0.95 Å; N—H 0.88 Å) and were included in the refinement in the riding model approximation, with  $U_{\text{iso}}(\text{H})$  1.2  $U_{\text{eq}}(\text{C})$ . The hydroxy and water H-atoms were located in a difference Fourier map, and were refined with distance restraints of O—H 0.84±0.01 Å and H···H 1.37±0.01 Å.

The final difference Fourier map had a peak of 1.37eÅ<sup>-3</sup> at 0.69Å from Br1 and a hole of -1.81eÅ<sup>-3</sup> at 1.33Å from C2.



**Figure 1**

View (Barbour, 2001) of *N*-(5-bromo-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide with displacement ellipsoids at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

***N'*-(5-Bromo-2-hydroxybenzylidene)-3,4,5-trihydroxybenzohydrazide dihydrate***Crystal data* $M_r = 403.19$ Monoclinic,  $P2_1/c$ 

Hall symbol: -P 2ybc

 $a = 30.8424 (8) \text{ \AA}$  $b = 3.7999 (1) \text{ \AA}$  $c = 12.8484 (4) \text{ \AA}$  $\beta = 90.280 (2)^\circ$  $V = 1505.79 (7) \text{ \AA}^3$  $Z = 4$  $F(000) = 816$  $D_x = 1.779 \text{ Mg m}^{-3}$ Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$ 

Cell parameters from 2758 reflections

 $\theta = 3.2\text{--}27.4^\circ$  $\mu = 2.77 \text{ mm}^{-1}$  $T = 100 \text{ K}$ 

Needle, colorless

 $0.30 \times 0.03 \times 0.03 \text{ mm}$ *Data collection*

Bruker SMART APEX

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  scansAbsorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996) $T_{\min} = 0.658$ ,  $T_{\max} = 0.921$ 

9964 measured reflections

3424 independent reflections

2914 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.039$  $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 1.3^\circ$  $h = -38 \rightarrow 40$  $k = -4 \rightarrow 4$  $l = -15 \rightarrow 16$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.064$  $wR(F^2) = 0.155$  $S = 1.22$ 

3424 reflections

241 parameters

10 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

 $w = 1/[\sigma^2(F_o^2) + (0.0499P)^2 + 10.2476P]$   
where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 1.08 \text{ e \AA}^{-3}$  $\Delta\rho_{\min} = -1.82 \text{ e \AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.467910 (16)	0.31017 (16)	0.89799 (4)	0.01787 (17)
O1	0.32358 (12)	0.6446 (12)	0.5944 (3)	0.0201 (9)
H1O	0.3020 (15)	0.745 (18)	0.621 (5)	0.030*
O2	0.20078 (11)	0.9755 (11)	0.6545 (3)	0.0182 (8)
O3	0.05472 (13)	1.3768 (13)	0.7962 (4)	0.0295 (11)
H3O	0.0335 (16)	1.36 (2)	0.836 (5)	0.044*
O4	0.05208 (14)	1.1036 (15)	0.9894 (4)	0.0353 (12)

H4O	0.048 (3)	0.933 (15)	1.030 (6)	0.053*
O5	0.12324 (12)	0.7692 (12)	1.0737 (3)	0.0182 (9)
H5O	0.1471 (12)	0.682 (18)	1.092 (5)	0.027*
O1W	0.25912 (12)	1.3152 (13)	0.9988 (3)	0.0209 (9)
H1W1	0.2444 (17)	1.382 (18)	1.050 (3)	0.031*
H1W2	0.2806 (14)	1.199 (17)	1.020 (4)	0.031*
O2W	-0.02887 (15)	1.3435 (17)	0.8657 (5)	0.0506 (16)
H2W1	-0.042 (3)	1.30 (2)	0.810 (4)	0.076*
H2W2	-0.038 (3)	1.534 (14)	0.890 (7)	0.076*
N1	0.27850 (13)	0.8683 (12)	0.7531 (3)	0.0139 (9)
N2	0.24174 (13)	0.9859 (13)	0.8027 (3)	0.0138 (9)
H2N	0.2431	1.0511	0.8684	0.017*
C1	0.35563 (16)	0.5778 (15)	0.6648 (4)	0.0144 (10)
C2	0.39381 (16)	0.4251 (16)	0.6295 (4)	0.0173 (11)
H2	0.3970	0.3730	0.5576	0.021*
C3	0.42746 (17)	0.3479 (16)	0.6985 (4)	0.0193 (12)
H3	0.4535	0.2431	0.6743	0.023*
C4	0.42245 (16)	0.4256 (16)	0.8025 (4)	0.0160 (11)
C5	0.38470 (16)	0.5781 (15)	0.8398 (4)	0.0143 (10)
H5	0.3820	0.6294	0.9118	0.017*
C6	0.35065 (15)	0.6562 (15)	0.7712 (4)	0.0126 (10)
C7	0.31105 (16)	0.7988 (15)	0.8138 (4)	0.0130 (10)
H7	0.3089	0.8415	0.8865	0.016*
C8	0.20343 (16)	1.0010 (13)	0.7503 (4)	0.0108 (10)
C9	0.16507 (16)	1.0460 (15)	0.8180 (4)	0.0130 (10)
C10	0.12793 (16)	1.2046 (16)	0.7771 (4)	0.0163 (11)
H10	0.1281	1.2984	0.7086	0.020*
C11	0.09101 (17)	1.2242 (15)	0.8368 (4)	0.0176 (12)
C12	0.09010 (17)	1.0775 (17)	0.9364 (4)	0.0204 (12)
C13	0.12713 (17)	0.9146 (15)	0.9777 (4)	0.0152 (11)
C14	0.16479 (17)	0.9040 (15)	0.9188 (4)	0.0149 (11)
H14	0.1903	0.8006	0.9468	0.018*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0094 (2)	0.0195 (3)	0.0247 (3)	-0.0001 (2)	-0.00302 (17)	-0.0020 (3)
O1	0.0160 (18)	0.032 (3)	0.0125 (17)	0.0031 (18)	-0.0028 (14)	-0.0016 (17)
O2	0.0127 (17)	0.027 (2)	0.0148 (18)	-0.0039 (16)	-0.0015 (14)	0.0021 (17)
O3	0.0142 (19)	0.041 (3)	0.034 (2)	0.0120 (19)	-0.0051 (17)	-0.005 (2)
O4	0.016 (2)	0.060 (4)	0.030 (2)	0.009 (2)	0.0035 (18)	-0.013 (2)
O5	0.0129 (17)	0.031 (2)	0.0110 (17)	-0.0007 (17)	0.0027 (14)	-0.0045 (16)
O1W	0.0162 (18)	0.035 (2)	0.0113 (17)	0.0027 (18)	0.0015 (14)	-0.0044 (18)
O2W	0.016 (2)	0.049 (4)	0.087 (5)	0.010 (2)	0.010 (2)	0.013 (3)
N1	0.0107 (19)	0.014 (3)	0.017 (2)	-0.0016 (17)	0.0010 (16)	-0.0008 (18)
N2	0.0096 (19)	0.019 (3)	0.013 (2)	-0.0004 (18)	-0.0007 (15)	-0.0028 (18)
C1	0.014 (2)	0.016 (3)	0.014 (2)	-0.002 (2)	-0.0013 (19)	-0.002 (2)
C2	0.013 (2)	0.025 (3)	0.014 (2)	-0.001 (2)	0.0024 (19)	0.000 (2)

C3	0.013 (2)	0.020 (3)	0.025 (3)	0.001 (2)	0.006 (2)	-0.003 (2)
C4	0.012 (2)	0.014 (3)	0.021 (3)	-0.001 (2)	-0.003 (2)	0.003 (2)
C5	0.015 (2)	0.013 (3)	0.015 (2)	-0.001 (2)	0.0005 (19)	-0.002 (2)
C6	0.010 (2)	0.015 (3)	0.013 (2)	-0.003 (2)	0.0008 (18)	0.001 (2)
C7	0.013 (2)	0.015 (3)	0.012 (2)	-0.006 (2)	0.0022 (18)	-0.003 (2)
C8	0.013 (2)	0.004 (3)	0.015 (2)	-0.0007 (18)	-0.0019 (18)	0.0009 (19)
C9	0.011 (2)	0.013 (3)	0.015 (2)	0.000 (2)	-0.0027 (19)	-0.003 (2)
C10	0.013 (2)	0.016 (3)	0.020 (3)	0.001 (2)	-0.0034 (19)	0.000 (2)
C11	0.017 (2)	0.013 (3)	0.022 (3)	0.008 (2)	-0.007 (2)	-0.007 (2)
C12	0.012 (2)	0.027 (3)	0.022 (3)	0.002 (2)	0.001 (2)	-0.012 (2)
C13	0.016 (2)	0.017 (3)	0.013 (2)	0.001 (2)	0.0025 (19)	-0.008 (2)
C14	0.013 (2)	0.016 (3)	0.015 (2)	0.001 (2)	-0.0022 (19)	-0.006 (2)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

Br1—C4	1.909 (5)	C1—C6	1.408 (7)
O1—C1	1.361 (6)	C2—C3	1.393 (8)
O1—H1O	0.840 (10)	C2—H2	0.9500
O2—C8	1.236 (6)	C3—C4	1.378 (8)
O3—C11	1.362 (6)	C3—H3	0.9500
O3—H3O	0.838 (10)	C4—C5	1.388 (7)
O4—C12	1.362 (7)	C5—C6	1.400 (7)
O4—H4O	0.840 (10)	C5—H5	0.9500
O5—C13	1.357 (7)	C6—C7	1.446 (7)
O5—H5O	0.841 (10)	C7—H7	0.9500
O1W—H1W1	0.839 (10)	C8—C9	1.482 (7)
O1W—H1W2	0.838 (10)	C9—C10	1.395 (7)
O2W—H2W1	0.838 (10)	C9—C14	1.403 (7)
O2W—H2W2	0.839 (10)	C10—C11	1.378 (7)
N1—C7	1.296 (7)	C10—H10	0.9500
N1—N2	1.378 (6)	C11—C12	1.396 (8)
N2—C8	1.359 (6)	C12—C13	1.401 (8)
N2—H2N	0.8800	C13—C14	1.390 (7)
C1—C2	1.391 (7)	C14—H14	0.9500
C1—O1—H1O	113 (5)	C5—C6—C7	118.3 (4)
C11—O3—H3O	111 (6)	C1—C6—C7	122.9 (5)
C12—O4—H4O	112 (6)	N1—C7—C6	120.1 (4)
C13—O5—H5O	110 (5)	N1—C7—H7	120.0
H1W1—O1W—H1W2	109.6 (18)	C6—C7—H7	120.0
H2W1—O2W—H2W2	109.5 (18)	O2—C8—N2	122.9 (4)
C7—N1—N2	115.1 (4)	O2—C8—C9	123.0 (4)
C8—N2—N1	120.0 (4)	N2—C8—C9	114.1 (4)
C8—N2—H2N	120.0	C10—C9—C14	120.3 (5)
N1—N2—H2N	120.0	C10—C9—C8	119.0 (5)
O1—C1—C2	118.3 (5)	C14—C9—C8	120.4 (5)
O1—C1—C6	121.6 (5)	C11—C10—C9	119.5 (5)
C2—C1—C6	120.1 (5)	C11—C10—H10	120.2

C1—C2—C3	120.6 (5)	C9—C10—H10	120.2
C1—C2—H2	119.7	O3—C11—C10	119.3 (5)
C3—C2—H2	119.7	O3—C11—C12	120.1 (5)
C4—C3—C2	119.1 (5)	C10—C11—C12	120.6 (5)
C4—C3—H3	120.5	O4—C12—C11	116.7 (5)
C2—C3—H3	120.5	O4—C12—C13	123.0 (5)
C3—C4—C5	121.5 (5)	C11—C12—C13	120.3 (5)
C3—C4—Br1	119.2 (4)	O5—C13—C14	124.1 (5)
C5—C4—Br1	119.3 (4)	O5—C13—C12	116.7 (5)
C4—C5—C6	120.0 (5)	C14—C13—C12	119.2 (5)
C4—C5—H5	120.0	C13—C14—C9	120.0 (5)
C6—C5—H5	120.0	C13—C14—H14	120.0
C5—C6—C1	118.7 (5)	C9—C14—H14	120.0
C7—N1—N2—C8	-167.2 (5)	N2—C8—C9—C10	154.0 (5)
O1—C1—C2—C3	-179.3 (5)	O2—C8—C9—C14	147.8 (5)
C6—C1—C2—C3	0.1 (9)	N2—C8—C9—C14	-31.4 (7)
C1—C2—C3—C4	-0.1 (9)	C14—C9—C10—C11	0.6 (9)
C2—C3—C4—C5	0.1 (9)	C8—C9—C10—C11	175.2 (5)
C2—C3—C4—Br1	178.5 (4)	C9—C10—C11—O3	-179.5 (5)
C3—C4—C5—C6	0.0 (9)	C9—C10—C11—C12	-2.0 (9)
Br1—C4—C5—C6	-178.4 (4)	O3—C11—C12—O4	-1.4 (8)
C4—C5—C6—C1	-0.1 (8)	C10—C11—C12—O4	-178.9 (6)
C4—C5—C6—C7	177.0 (5)	O3—C11—C12—C13	178.9 (5)
O1—C1—C6—C5	179.4 (5)	C10—C11—C12—C13	1.4 (9)
C2—C1—C6—C5	0.1 (8)	O4—C12—C13—O5	2.2 (9)
O1—C1—C6—C7	2.5 (9)	C11—C12—C13—O5	-178.1 (5)
C2—C1—C6—C7	-176.9 (5)	O4—C12—C13—C14	-179.1 (6)
N2—N1—C7—C6	176.8 (5)	C11—C12—C13—C14	0.6 (9)
C5—C6—C7—N1	-178.7 (5)	O5—C13—C14—C9	176.7 (5)
C1—C6—C7—N1	-1.8 (8)	C12—C13—C14—C9	-1.9 (8)
N1—N2—C8—O2	-13.6 (8)	C10—C9—C14—C13	1.3 (8)
N1—N2—C8—C9	165.7 (5)	C8—C9—C14—C13	-173.2 (5)
O2—C8—C9—C10	-26.7 (8)		

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O1—H1o $\cdots$ N1	0.84 (1)	1.91 (5)	2.616 (6)	141 (7)
O3—H3o $\cdots$ O2w	0.84 (1)	1.96 (4)	2.736 (6)	153 (7)
O4—H4o $\cdots$ O2w <sup>i</sup>	0.84 (1)	1.81 (3)	2.623 (8)	163 (9)
O5—H5o $\cdots$ O2 <sup>ii</sup>	0.84 (1)	1.93 (2)	2.764 (5)	171 (7)
O1w—H1w1 $\cdots$ O2 <sup>iii</sup>	0.84 (1)	1.98 (2)	2.812 (5)	170 (6)
O1w—H1w2 $\cdots$ O1 <sup>ii</sup>	0.84 (1)	2.09 (2)	2.914 (6)	167 (6)
O2w—H2w1 $\cdots$ O3 <sup>iv</sup>	0.84 (1)	2.13 (5)	2.845 (9)	142 (8)
O2w—H2w2 $\cdots$ O4 <sup>v</sup>	0.84 (1)	2.12 (4)	2.900 (8)	154 (8)

Symmetry codes: (i)  $-x, -y+2, -z+2$ ; (ii)  $x, -y+3/2, z+1/2$ ; (iii)  $x, -y+5/2, z+1/2$ ; (iv)  $-x, y-1/2, -z+3/2$ ; (v)  $-x, -y+3, -z+2$ .