

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[μ_3 -2,2,4,4,6,6-Hexakis(3,5-dimethylpyrazol-1-yl)-2 λ^5 ,4 λ^5 ,6 λ^5 -1,3,5,2,4,6-triazatriphosphinine]tris[*cis*-dichlorido-palladium(II)]

Sung Yol Yun and Soon W. Lee*

Department of Chemistry (BK21), Sungkyunkwan University, Natural Science Campus, Suwon 440-746, Republic of Korea

Correspondence e-mail: soonwlee@skku.edu


Received 11 July 2008; accepted 23 July 2008

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.035; wR factor = 0.109; data-to-parameter ratio = 34.7.

The title complex, $[\text{Pd}_3\text{Cl}_6(\text{C}_{30}\text{H}_{42}\text{N}_{15}\text{P}_3)]$, possesses C_3 molecular symmetry. The P and N atoms of the cyclo-triphosphazene and the Pd atom are located on the crystallographic mirror plane. Each of the three symmetry-related Pd atoms is coordinated by two chloride ligands and two exocyclic pyrazolyl N atoms, but not by the cyclo-triphosphazene N atoms.

Related literature

For related literature, see: Chandrasekhar & Nagendran (2001); Gallicano & Paddock (1982).



Experimental

Crystal data

$[\text{Pd}_3\text{Cl}_6(\text{C}_{30}\text{H}_{42}\text{N}_{15}\text{P}_3)]$
 $M_r = 1237.60$
 Hexagonal, $P6_3/m$
 $a = 17.2989$ (3) Å
 $c = 14.4545$ (6) Å
 $V = 3746.02$ (18) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.02$ mm⁻¹
 $T = 296$ (2) K
 $0.24 \times 0.20 \times 0.16$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (North *et al.*, 1968)
 $T_{\min} = 0.792$, $T_{\max} = 0.854$

42917 measured reflections
 3157 independent reflections
 2098 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.108$
 $S = 1.07$
 3157 reflections

91 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.52$ e Å⁻³
 $\Delta\rho_{\min} = -0.35$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Pd1—N3	2.027 (2)	P1—N2	1.695 (2)
Pd1—Cl1	2.2642 (10)	N2—N3	1.384 (3)
N3—Pd1—N3 ⁱ	86.36 (14)	Cl1—Pd1—Cl1 ⁱ	88.95 (6)
N3—Pd1—Cl1	178.26 (8)	N1 ⁱⁱ —P1—N1	118.0 (2)
N3 ⁱ —Pd1—Cl1	92.34 (8)	N2 ⁱ —P1—N2	102.86 (17)

Symmetry codes: (i) $x, y, -z + \frac{3}{2}$; (ii) $-y + 1, x - y + 1, z$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINTE* (Bruker, 1997); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: KP2182).

References

- Bruker (1997). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Chandrasekhar, V. & Nagendran, S. (2001). *Chem. Soc. Rev.* **30**, 193–203.
 Gallicano, K. D. & Paddock, N. L. (1982). *Can. J. Chem.* **60**, 521–528.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2008). E64, m1084 [doi:10.1107/S1600536808023167]

[μ_3 -2,2,4,4,6,6-Hexakis(3,5-dimethylpyrazol-1-yl)-2 λ^5 ,4 λ^5 ,6 λ^5 -1,3,5,2,4,6-triaza-triphosphinine]tris[*cis*-dichloridopalladium(II)]

Sung Yol Yun and Soon W. Lee

S1. Comment

Various cyclotriphosphazene-based ligands have been designed and utilized to prepare coordination and organometallic complexes (Chandrasekhar & Nagendran, 2001). In particular, the 6-membered cyclic ligand $N_3P_3(3,5\text{-Me}_2\text{pz})_6$ (3,5-Me₂pz = 3,5-dimethylpyrazolyl) has many potential donor sites due to the exocyclic pyrazolyl nitrogen atoms in addition to the ring nitrogen and phosphorus atoms. This ligand was previously reported to react with $[\text{PdCl}_2(\text{PhCN})_2]$ to give the title complex, which was not structurally characterized by X-ray diffraction (Gallicano & Paddock, 1982). We chose the title complex to be used as a starting material with the C_3 -symmetry in preparing coordination polymers by treating it with organic linking ligands. In this context, we determined the three-dimensional structure of the title complex to confirm its molecular symmetry.

The central core has a perfectly planar hexagonal P_3N_3 unit, to which three surrounding square-planar palladium fragments (PdCl_2N_2) are perpendicular (Fig. 1, Table 1). The crystallographic mirror plane ($z = 3/4$) passes through the central cyclotriphosphazene ring (three P and three N atoms) and the three surrounding palladium atoms, and bisects the pendant germinal pyrazolyl ligands in each, symmetry related $\text{PdCl}_2(\text{pyrazolyl})_2$ unit. A space filling model of the title complex (Fig. 2) shows its C_3 -symmetry and close packing. As previously predicted by NMR and IR spectroscopy (Gallicano & Paddock, 1982), each palladium metal is coordinated by two chloro ligands and exocyclic pyrazolyl N atoms, but not to the cyclotriphosphazene N atoms, and lies 0.022 (1) Å below the Cl_2N_2 plane. Each phosphorus atom is bound to four N atoms: two central cyclotriphosphazene N atoms and two exocyclic pyrazolyl N atoms. Consistently with our expectation, the P1—N1 (cyclotriphosphazene) bond is significantly longer than P1—N2 (pyrazolyl) bond. All the Pd···Pd separations are equal (7.7538 (6) Å) due to the crystallographic symmetry.

S2. Experimental

The title complex was prepared by the literature method (Gallicano & Paddock, 1982). The product was recrystallized from a mixture of dichloromethane–hexane.

S3. Refinement

All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were generated in ideal positions and refined in a riding model.





Figure 1

Molecular structure showing the 50% probability displacement ellipsoids. H atoms are omitted for clarity.

**Figure 2**

A space filling model of the title complex showing its C_3 axis at the center of the cyclotriphosphazene ring: (a) red: Pd; green: Cl; orange: P; purple: N; grey: C; white, H.

[μ_3 -2,2,4,4,6,6-Hexakis(3,5-dimethylpyrazol-1-yl)-2,2 λ^5 ,4,4 λ^5 ,6,6 λ^5 -1,3,5,2,4,6-triazatriphosphinine]tris[*cis*-dichloridopalladium(II)]

Crystal data

[Pd₃Cl₆(C₃₀H₄₂N₁₅P₃)]

$M_r = 1237.60$

Hexagonal, $P6_3/m$

Hall symbol: -P6c

$a = 17.2989$ (3) Å

$c = 14.4545$ (6) Å

$V = 3746.02$ (18) Å³

$Z = 2$

$F(000) = 1224$

$D_x = 1.097$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9849 reflections

$\theta = 2.4$ – 27.2°

$\mu = 1.02$ mm⁻¹

$T = 296$ K

Block, yellow

$0.24 \times 0.20 \times 0.16$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(North *et al.*, 1968)

$T_{\min} = 0.792$, $T_{\max} = 0.854$

42917 measured reflections

3157 independent reflections

2098 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.048$
 $\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 3.6^\circ$
 $h = -23 \rightarrow 22$

$k = -19 \rightarrow 22$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.108$
 $S = 1.07$
 3157 reflections
 91 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0477P)^2 + 1.7582P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.002$
 $\Delta\rho_{\text{max}} = 0.52 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pd1	0.13572 (2)	0.37375 (2)	0.7500	0.05476 (14)
Cl1	0.02844 (7)	0.32969 (8)	0.64026 (8)	0.0961 (3)
P1	0.31769 (7)	0.56802 (7)	0.7500	0.0438 (2)
N1	0.2362 (2)	0.5861 (2)	0.7500	0.0472 (7)
N2	0.30675 (15)	0.50221 (15)	0.84168 (15)	0.0503 (5)
C1	0.3607 (2)	0.5175 (2)	0.9179 (2)	0.0697 (9)
C2	0.3213 (3)	0.4397 (3)	0.9672 (3)	0.0873 (12)
H2	0.3425	0.4290	1.0220	0.105*
N3	0.23411 (17)	0.41658 (16)	0.84595 (17)	0.0573 (6)
C3	0.2439 (2)	0.3792 (2)	0.9210 (3)	0.0743 (10)
C4	0.4433 (3)	0.6016 (3)	0.9398 (3)	0.1093 (17)
H4A	0.4555	0.6442	0.8914	0.164*
H4B	0.4921	0.5904	0.9449	0.164*
H4C	0.4359	0.6249	0.9974	0.164*
C5	0.1818 (3)	0.2838 (3)	0.9456 (4)	0.121 (2)
H5A	0.1345	0.2576	0.9008	0.181*
H5B	0.1570	0.2805	1.0058	0.181*
H5C	0.2139	0.2518	0.9458	0.181*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pd1	0.0519 (2)	0.0567 (2)	0.05085 (19)	0.02349 (16)	0.000	0.000
Cl1	0.0739 (6)	0.1049 (8)	0.0898 (7)	0.0300 (6)	-0.0295 (5)	-0.0085 (6)
P1	0.0508 (6)	0.0488 (6)	0.0333 (4)	0.0259 (5)	0.000	0.000
N1	0.0495 (18)	0.0505 (19)	0.0375 (15)	0.0218 (15)	0.000	0.000
N2	0.0538 (14)	0.0508 (13)	0.0433 (11)	0.0240 (12)	-0.0024 (10)	0.0062 (9)
C1	0.071 (2)	0.072 (2)	0.0544 (17)	0.0276 (18)	-0.0132 (15)	0.0103 (15)
C2	0.088 (3)	0.080 (2)	0.073 (2)	0.026 (2)	-0.019 (2)	0.0274 (19)
N3	0.0636 (16)	0.0503 (14)	0.0518 (13)	0.0238 (12)	-0.0023 (11)	0.0097 (10)
C3	0.074 (2)	0.064 (2)	0.068 (2)	0.0219 (18)	-0.0088 (17)	0.0191 (16)
C4	0.101 (3)	0.095 (3)	0.072 (2)	0.005 (2)	-0.041 (2)	0.024 (2)
C5	0.117 (4)	0.079 (3)	0.122 (4)	0.016 (3)	-0.028 (3)	0.047 (3)

Geometric parameters (\AA , $^\circ$)

Pd1—N3	2.027 (2)	C1—C4	1.476 (5)
Pd1—N3 ⁱ	2.027 (2)	C2—C3	1.389 (5)
Pd1—Cl1	2.2642 (10)	C2—H2	0.9300
Pd1—Cl1 ⁱ	2.2641 (10)	N3—C3	1.317 (4)
P1—N1 ⁱⁱ	1.557 (3)	C3—C5	1.494 (5)
P1—N1	1.589 (3)	C4—H4A	0.9600
P1—N2 ⁱ	1.695 (2)	C4—H4B	0.9600
P1—N2	1.695 (2)	C4—H4C	0.9600
N1—P1 ⁱⁱⁱ	1.557 (3)	C5—H5A	0.9600
N2—C1	1.381 (4)	C5—H5B	0.9600
N2—N3	1.384 (3)	C5—H5C	0.9600
C1—C2	1.367 (5)		
N3—Pd1—N3 ⁱ	86.36 (14)	C1—C2—H2	126.0
N3—Pd1—Cl1	178.26 (8)	C3—C2—H2	126.0
N3 ⁱ —Pd1—Cl1	92.34 (8)	C3—N3—N2	107.0 (2)
N3—Pd1—Cl1 ⁱ	92.34 (8)	C3—N3—Pd1	132.5 (2)
N3 ⁱ —Pd1—Cl1 ⁱ	178.26 (8)	N2—N3—Pd1	120.51 (16)
Cl1—Pd1—Cl1 ⁱ	88.95 (6)	N3—C3—C2	109.7 (3)
N1 ⁱⁱ —P1—N1	118.0 (2)	N3—C3—C5	122.7 (3)
N1 ⁱⁱ —P1—N2 ⁱ	108.78 (11)	C2—C3—C5	127.4 (3)
N1—P1—N2 ⁱ	108.66 (11)	C1—C4—H4A	109.5
N1 ⁱⁱ —P1—N2	108.78 (11)	C1—C4—H4B	109.5
N1—P1—N2	108.66 (11)	H4A—C4—H4B	109.5
N2 ⁱ —P1—N2	102.86 (17)	C1—C4—H4C	109.5
P1 ⁱⁱⁱ —N1—P1	122.0 (2)	H4A—C4—H4C	109.5
C1—N2—N3	109.5 (2)	H4B—C4—H4C	109.5
C1—N2—P1	131.1 (2)	C3—C5—H5A	109.5
N3—N2—P1	119.37 (17)	C3—C5—H5B	109.5
C2—C1—N2	105.7 (3)	H5A—C5—H5B	109.5
C2—C1—C4	128.3 (3)	C3—C5—H5C	109.5

N2—C1—C4	126.0 (3)	H5A—C5—H5C	109.5
C1—C2—C3	108.1 (3)	H5B—C5—H5C	109.5

Symmetry codes: (i) $x, y, -z+3/2$; (ii) $-y+1, x-y+1, z$; (iii) $-x+y, -x+1, z$.