

5-(1*H*-1,2,3-Benzotriazol-1-ylmethyl)-3-phenyl-1,2,4-oxadiazole

Shu-Qing Xu and Jia-Ming Li*

Laboratory of Beibu Gulf Marine Protection and Exploitation, Department of Chemistry and Biology, Qinzhou University, Qinzhou, Guangxi 535000, People's Republic of China
Correspondence e-mail: ljmimarise@163.com


Received 21 June 2008; accepted 30 June 2008

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.051; wR factor = 0.103; data-to-parameter ratio = 12.2.

In the title molecule, $\text{C}_{15}\text{H}_{11}\text{N}_5\text{O}$, the 1,2,3-benzotriazole and 3-phenyl-1,2,4-oxadiazole units are individually essentially planar and the dihedral angle between them is $80.2(2)^\circ$. In the crystal structure, molecules are connected via weak intermolecular $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds to form two-dimensional sheets.

Related literature

For related literature, see: Batista *et al.* (2000); Wardell *et al.* (2003); Srinivasan *et al.* (2007); Wang *et al.* (2004a,b,c, 2007); Romero (2001); Terashita *et al.* (2002); Zen *et al.* (1983).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{11}\text{N}_5\text{O}$
 $M_r = 277.29$
Monoclinic, $P2_1/c$
 $a = 4.7009(13)\text{ \AA}$
 $b = 11.100(3)\text{ \AA}$
 $c = 25.265(7)\text{ \AA}$
 $\beta = 95.234(6)^\circ$

$V = 1312.8(6)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 295\text{ K}$
 $0.18 \times 0.14 \times 0.12\text{ mm}$

Data collection

Bruker SMART diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.983$, $T_{\max} = 0.989$
6803 measured reflections
2322 independent reflections
1324 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.103$
 $S = 1.02$
190 parameters
H-atom parameters constrained
2322 reflections
 $\Delta\rho_{\max} = 0.13\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C9—H9B···N5 ⁱ	0.97	2.59	3.443 (3)	147
C9—H9A···N2 ⁱⁱ	0.97	2.60	3.466 (3)	149

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x - 1, y, z$.

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by a grant from the Qinzhou University Foundation of Guangxi Zhuang Autonomous Region of the People's Republic of China (grant No. 2007XJ15).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2650).

References

- Batista, H., Carpenter, G. B. & Srivastava, R. M. (2000). *J. Chem. Crystallogr.* **30**, 131–134.
- Bruker (2003). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Romero, J. R. (2001). *Expert Opin. Investig. Drugs*, **10**, 369–379.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Srinivasan, N., Nandhini, M. S., Ranjithkumar, R., Perumal, S. & Krishnakumar, R. V. (2007). *Acta Cryst. E* **63**, o3750.
- Terashita, Z., Naruo, K. & Morimoto, S. (2002). PCT Int. Appl. WO 02060439.
- Wang, H.-B., Chen, J.-H. & Wang, J.-T. (2004a). *Acta Cryst. E* **60**, o1478–o1480.
- Wang, H.-B., Chen, J.-H. & Wang, J.-T. (2004b). *Acta Cryst. E* **60**, o1709–o1711.
- Wang, H.-B., Chen, J.-H. & Wang, J.-T. (2004c). *Acta Cryst. E* **60**, o1917–o1918.
- Wang, P., Wang, H., Kang, S., Li, H. & Wu, W. (2007). *Acta Cryst. E* **63**, o4180.
- Wardell, S. M. S. V., Carvalho, C. E. M., Low, J. N. & Glidewell, C. (2003). *Acta Cryst. E* **59**, o1729–o1730.
- Zen, S., Nishino, T., Harada, K., Nakamura, H. & Iitaka, Y. (1983). *Chem. Pharm. Bull.* **31**, 4181–4184.

supporting information

Acta Cryst. (2008). E64, o1403 [doi:10.1107/S1600536808019879]

5-(1*H*-1,2,3-Benzotriazol-1-ylmethyl)-3-phenyl-1,2,4-oxadiazole

Shu-Qing Xu and Jia-Ming Li

S1. Comment


Synthesis of 1,2,4-oxadiazole derivatives has attracted a great interest due to their pharmacological properties such as intrinsic analgesic (Zen *et al.*, 1983; Terashita *et al.*, 2002) and antipicornaviral (Romero, 2001) effects. Wang *et al.* (2004a,b,c;2007) have described the synthesis and crystal structures of a series of these types of compounds. Herein, we report the synthesis and crystal structure of the title compound, (I), containing both 1,2,4-oxadiazole and 1,2,3-benzotriazole organic functional groups. The molecular structure of (I) is shown in Fig. 1. The title molecule can be considered as two rings systems: the 3-phenyl-1,2,4-oxadiazol-5-yl (A) and benzotriazole (B). All atoms in A, B are individually essentially planar with dihedral angles of 4.4 (2) ° and 0.5 (2) ° between the rings in each, respectively. The dihedral angle between A and B is 80.2 (2) °. This conformation presents no nonbonded interactions (Batista *et al.*, 2000). Molecules are connected *via* weak intermolecular C—H···N hydrogen bonds (Table. 1 and Fig. 2) to form two-dimensional sheets.

S2. Experimental


Reagents and solvents used were of commercially available quality. 1,2,3-Benzotriazole (1 mmol) was dissolved in acetonitrile (80 ml) and potassium carbonate (15 mmol) was added followed by 3-phenyl-5-chloromethyl-1,2,4-oxadiazole (1 mmol). The resulting mixture was refluxed for 10 h. After cooling and filtering, the crude title compound was obtained and purified by recrystallization from ethyl acetate. Crystals suitable for X-ray diffraction were obtained by slow evaporation of an ethanol solution of (I).

S3. Refinement

H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93–0.97 Å and with $U_{\text{iso}}(\text{H})$ = 1.2 times $U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure with displacement ellipsoids at the 30% probability level.

**Figure 2**

Part of the crystal structure showing hydrogen bonds as dashed lines. H atoms, except for those involved in hydrogen bonds, are not included.

5-(1*H*-1,2,3-Benzotriazol-1-ylmethyl)-3-phenyl-1,2,4-oxadiazole

Crystal data

C₁₅H₁₁N₅O
M_r = 277.29

Monoclinic, P2₁/c
Hall symbol: -P 2ybc

$a = 4.7009 (13)$ Å
 $b = 11.100 (3)$ Å
 $c = 25.265 (7)$ Å
 $\beta = 95.234 (6)^\circ$
 $V = 1312.8 (6)$ Å³
 $Z = 4$
 $F(000) = 576$
 $D_x = 1.403$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 760 reflections
 $\theta = 2.5\text{--}19.5^\circ$
 $\mu = 0.09$ mm⁻¹
 $T = 295$ K
Block, colorless
 $0.18 \times 0.14 \times 0.12$ mm

Data collection

Bruker SMART
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)
 $T_{\min} = 0.983$, $T_{\max} = 0.989$

6803 measured reflections
2322 independent reflections
1324 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.6^\circ$
 $h = -5 \rightarrow 5$
 $k = -13 \rightarrow 13$
 $l = -20 \rightarrow 29$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.103$
 $S = 1.02$
2322 reflections
190 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.036P)^2 + 0.0112P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.13$ e Å⁻³
 $\Delta\rho_{\min} = -0.19$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.1731 (4)	0.96111 (14)	0.37087 (7)	0.0620 (5)
N1	0.3748 (5)	0.94501 (18)	0.41561 (8)	0.0626 (6)
N2	0.4253 (4)	0.80085 (17)	0.35428 (7)	0.0477 (5)
N3	0.1502 (4)	0.80219 (18)	0.24592 (8)	0.0488 (5)
N4	0.1194 (5)	0.68098 (19)	0.23986 (9)	0.0626 (6)
N5	0.2711 (5)	0.64667 (18)	0.20147 (9)	0.0644 (6)
C1	0.5148 (5)	0.8490 (2)	0.40357 (9)	0.0443 (6)
C2	0.7410 (5)	0.7967 (2)	0.43988 (9)	0.0445 (6)
C3	0.8252 (5)	0.8510 (2)	0.48824 (10)	0.0587 (7)

H3	0.7393	0.9224	0.4976	0.070*
C4	1.0355 (6)	0.7994 (3)	0.52249 (10)	0.0669 (8)
H4	1.0927	0.8369	0.5546	0.080*
C5	1.1613 (6)	0.6933 (3)	0.50957 (11)	0.0675 (8)
H5	1.3019	0.6586	0.5330	0.081*
C6	1.0795 (6)	0.6382 (2)	0.46194 (11)	0.0662 (8)
H6	1.1652	0.5663	0.4532	0.079*
C7	0.8703 (5)	0.6893 (2)	0.42703 (10)	0.0550 (7)
H7	0.8157	0.6517	0.3948	0.066*
C8	0.2200 (5)	0.8714 (2)	0.33742 (9)	0.0460 (6)
C9	0.0215 (5)	0.8644 (2)	0.28801 (9)	0.0595 (7)
H9A	-0.1512	0.8225	0.2956	0.071*
H9B	-0.0313	0.9452	0.2763	0.071*
C10	0.3265 (5)	0.8462 (2)	0.21070 (9)	0.0428 (6)
C11	0.4018 (5)	0.7457 (2)	0.18258 (9)	0.0472 (6)
C12	0.5861 (5)	0.7564 (2)	0.14226 (10)	0.0578 (7)
H12	0.6374	0.6898	0.1229	0.069*
C13	0.6871 (5)	0.8689 (3)	0.13269 (10)	0.0630 (7)
H13	0.8125	0.8788	0.1066	0.076*
C14	0.6070 (5)	0.9691 (2)	0.16101 (10)	0.0594 (7)
H14	0.6790	1.0442	0.1529	0.071*
C15	0.4258 (5)	0.9607 (2)	0.20035 (10)	0.0515 (6)
H15	0.3724	1.0279	0.2190	0.062*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0714 (12)	0.0570 (11)	0.0567 (12)	0.0150 (9)	0.0015 (10)	-0.0011 (10)
N1	0.0734 (15)	0.0596 (14)	0.0533 (15)	0.0106 (12)	-0.0026 (12)	-0.0082 (12)
N2	0.0492 (12)	0.0528 (12)	0.0405 (13)	0.0061 (10)	0.0007 (10)	0.0001 (11)
N3	0.0531 (12)	0.0484 (13)	0.0431 (13)	0.0011 (10)	-0.0055 (10)	0.0021 (11)
N4	0.0712 (15)	0.0494 (15)	0.0642 (16)	-0.0082 (11)	-0.0104 (13)	0.0118 (13)
N5	0.0798 (17)	0.0458 (14)	0.0649 (16)	-0.0017 (12)	-0.0070 (13)	-0.0007 (13)
C1	0.0483 (14)	0.0460 (15)	0.0392 (16)	-0.0039 (12)	0.0066 (12)	0.0000 (13)
C2	0.0461 (14)	0.0511 (15)	0.0370 (14)	-0.0060 (12)	0.0069 (12)	-0.0030 (13)
C3	0.0609 (17)	0.0646 (17)	0.0502 (17)	-0.0045 (14)	0.0033 (14)	-0.0087 (15)
C4	0.0683 (19)	0.086 (2)	0.0451 (17)	-0.0147 (17)	-0.0028 (15)	-0.0076 (17)
C5	0.0631 (18)	0.084 (2)	0.0536 (19)	-0.0015 (17)	-0.0062 (15)	0.0128 (18)
C6	0.0714 (19)	0.0660 (18)	0.0596 (19)	0.0080 (15)	-0.0031 (15)	-0.0014 (16)
C7	0.0584 (16)	0.0633 (17)	0.0421 (15)	-0.0026 (14)	-0.0024 (13)	-0.0053 (14)
C8	0.0496 (15)	0.0492 (16)	0.0401 (16)	0.0014 (13)	0.0087 (13)	0.0022 (13)
C9	0.0555 (16)	0.0702 (18)	0.0519 (17)	0.0105 (13)	0.0005 (14)	0.0054 (15)
C10	0.0470 (14)	0.0412 (14)	0.0375 (14)	-0.0012 (12)	-0.0113 (12)	-0.0017 (13)
C11	0.0563 (15)	0.0371 (14)	0.0449 (15)	0.0025 (13)	-0.0138 (13)	0.0014 (13)
C12	0.0666 (17)	0.0570 (17)	0.0479 (16)	0.0153 (14)	-0.0052 (14)	-0.0071 (14)
C13	0.0647 (18)	0.073 (2)	0.0509 (18)	0.0072 (15)	0.0034 (14)	0.0064 (16)
C14	0.0647 (17)	0.0539 (17)	0.0584 (18)	-0.0072 (14)	-0.0015 (15)	0.0066 (15)
C15	0.0580 (16)	0.0431 (15)	0.0508 (17)	-0.0012 (13)	-0.0095 (13)	-0.0049 (13)

Geometric parameters (\AA , $\text{^{\circ}}$)

O1—C8	1.338 (3)	C5—H5	0.9300
O1—N1	1.419 (2)	C6—C7	1.382 (3)
N1—C1	1.303 (3)	C6—H6	0.9300
N2—C8	1.284 (3)	C7—H7	0.9300
N2—C1	1.385 (3)	C8—C9	1.491 (3)
N3—N4	1.360 (2)	C9—H9A	0.9700
N3—C10	1.361 (3)	C9—H9B	0.9700
N3—C9	1.446 (3)	C10—C11	1.386 (3)
N4—N5	1.312 (3)	C10—C15	1.387 (3)
N5—C11	1.366 (3)	C11—C12	1.401 (3)
C1—C2	1.460 (3)	C12—C13	1.365 (3)
C2—C3	1.387 (3)	C12—H12	0.9300
C2—C7	1.390 (3)	C13—C14	1.393 (3)
C3—C4	1.378 (3)	C13—H13	0.9300
C3—H3	0.9300	C14—C15	1.370 (3)
C4—C5	1.370 (3)	C14—H14	0.9300
C4—H4	0.9300	C15—H15	0.9300
C5—C6	1.373 (3)		
C8—O1—N1	105.82 (17)	C2—C7—H7	119.9
C1—N1—O1	103.41 (18)	N2—C8—O1	114.0 (2)
C8—N2—C1	102.7 (2)	N2—C8—C9	129.8 (2)
N4—N3—C10	110.2 (2)	O1—C8—C9	116.2 (2)
N4—N3—C9	120.5 (2)	N3—C9—C8	111.64 (19)
C10—N3—C9	129.1 (2)	N3—C9—H9A	109.3
N5—N4—N3	108.1 (2)	C8—C9—H9A	109.3
N4—N5—C11	108.5 (2)	N3—C9—H9B	109.3
N1—C1—N2	114.0 (2)	C8—C9—H9B	109.3
N1—C1—C2	122.2 (2)	H9A—C9—H9B	108.0
N2—C1—C2	123.8 (2)	N3—C10—C11	104.2 (2)
C3—C2—C7	118.8 (2)	N3—C10—C15	133.4 (2)
C3—C2—C1	120.9 (2)	C11—C10—C15	122.3 (2)
C7—C2—C1	120.2 (2)	N5—C11—C10	108.9 (2)
C4—C3—C2	120.2 (2)	N5—C11—C12	130.5 (2)
C4—C3—H3	119.9	C10—C11—C12	120.5 (2)
C2—C3—H3	119.9	C13—C12—C11	117.0 (2)
C5—C4—C3	120.5 (2)	C13—C12—H12	121.5
C5—C4—H4	119.7	C11—C12—H12	121.5
C3—C4—H4	119.7	C12—C13—C14	121.6 (2)
C4—C5—C6	119.9 (3)	C12—C13—H13	119.2
C4—C5—H5	120.0	C14—C13—H13	119.2
C6—C5—H5	120.0	C15—C14—C13	122.3 (2)
C5—C6—C7	120.2 (3)	C15—C14—H14	118.9
C5—C6—H6	119.9	C13—C14—H14	118.9
C7—C6—H6	119.9	C14—C15—C10	116.2 (2)
C6—C7—C2	120.3 (2)	C14—C15—H15	121.9

C6—C7—H7	119.9	C10—C15—H15	121.9
----------	-------	-------------	-------

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C9—H9B···N5 ⁱ	0.97	2.59	3.443 (3)	147
C9—H9A···N2 ⁱⁱ	0.97	2.60	3.466 (3)	149

Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $x-1, y, z$.