

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1-[2-(4-Fluorobenzoyloxy)-2-phenylethyl]-1*H*-benzimidazoleÖzden Özel Güven,^a Taner Erdoğan,^a Simon J. Coles^b and Tuncer Hökelek^{c*}

^aZonguldak Karaelmas University, Department of Chemistry, 67100 Zonguldak, Turkey, ^bSouthampton University, Department of Chemistry, Southampton SO17 1BJ, England, and ^cHacettepe University, Department of Physics, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 8 July 2008; accepted 9 July 2008

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.055; wR factor = 0.143; data-to-parameter ratio = 12.9.

The asymmetric unit of the title compound, $\text{C}_{22}\text{H}_{19}\text{FN}_2\text{O}$, contains two independent molecules. The planar benzimidazole ring systems are oriented with respect to the phenyl/fluorobenzene rings at dihedral angles of 31.10 (4)/ 45.17 (5) and 45.52 (5)/ 68.63 (5)°, respectively, for the two molecules. In the crystal structure, intermolecular $\text{C}-\text{H}\cdots\text{N}$ and intermolecular $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{F}$ hydrogen bonds link the molecules into a three-dimensional network. There are $\text{C}-\text{H}\cdots\pi$ contacts between the benzimidazole and fluorobenzene rings and a $\pi-\pi$ contact between the benzimidazole and phenyl ring systems [centroid-centroid distance = 4.575 (1) Å].

Related literature

For general background, see: Brammer & Feczko (1988); Özel Güven *et al.* (2007*a,b*). For related literature, see: Song & Shin (1998); Freer *et al.* (1986); Peeters *et al.* (1979*a,b*, 1996); Caira *et al.* (2004).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{19}\text{FN}_2\text{O}$
 $M_r = 346.39$
 Monoclinic, $P2_1/c$
 $a = 12.6946$ (2) Å
 $b = 18.1279$ (4) Å
 $c = 15.3547$ (3) Å
 $\beta = 95.747$ (1)°
 $V = 3515.76$ (12) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 120$ (2) K
 $0.55 \times 0.15 \times 0.14$ mm

Data collection

Bruker–Nonius KappaCCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2007)
 $T_{\min} = 0.953$, $T_{\max} = 0.988$
 41740 measured reflections
 8050 independent reflections
 5679 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.142$
 $S = 1.07$
 8050 reflections
 622 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.43$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.43$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg2 is the centroid of the N1'/N2'/C1'/C2'/C7' ring and Cg8 is the centroid of the C17'–C22' ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{N2}'$	0.97 (2)	2.62 (2)	3.498 (2)	150.5 (17)
$\text{C6}'-\text{H6A}\cdots\text{N2}'$	0.964 (19)	2.459 (19)	3.345 (2)	152.7 (16)
$\text{C13}-\text{H13}\cdots\text{F}^i$	1.00 (2)	2.54 (2)	3.493 (2)	159.6 (16)
$\text{C19}'-\text{H19A}\cdots\text{N2}''$	1.01 (2)	2.59 (2)	3.509 (2)	151.1 (16)
$\text{C5}-\text{H5}\cdots\text{Cg8}''''$	0.95 (2)	2.662 (19)	3.514 (2)	149.0 (15)
$\text{C22}'-\text{H22A}\cdots\text{Cg2}''$	0.98 (2)	2.62 (9)	3.518 (2)	141.9 (16)

Symmetry codes: (i) $x-1, y, z$; (ii) $x, -y+\frac{1}{2}, z-\frac{1}{2}$; (iii) $x, -y-\frac{1}{2}, z-\frac{3}{2}$; (iv) $-x+1, -y, -z+1$.

Data collection: COLLECT (Hooft, 1998); cell refinement: DENZO (Otwinowski & Minor, 1997) and COLLECT; data reduction: DENZO and COLLECT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON (Spek, 2003).

The authors acknowledge Zonguldak Karaelmas University Research Fund (grant No. 2004–13-02–16).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2436).

References

- Brammer, K. W. & Feczko, J. M. (1988). In *Antifungal Drugs*, edited by V. St. Georgiev, pp. 561–563. New York Academy of Science.
 Caira, M. R., Alkhamis, K. A. & Obaidat, R. M. (2004). *J. Pharm. Sci.* **93**, 601–611.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Freer, A. A., Pearson, A. & Salole, E. G. (1986). *Acta Cryst.* **C42**, 1350–1352.
 Hooft, R. W. W. (1998). COLLECT. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.

- Özel Güven, Ö., Erdoğan, T., Göker, H. & Yıldız, S. (2007a). *Bioorg. Med. Chem. Lett.* **17**, 2233–2236.
- Özel Güven, Ö., Erdoğan, T., Göker, H. & Yıldız, S. (2007b). *J. Heterocycl. Chem.* **44**, 731–734.
- Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1979a). *Acta Cryst.* **B35**, 2461–2464.
- Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1979b). *Bull. Soc. Chim. Belg.* **88**, 265–272.
- Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1996). *Acta Cryst.* **C52**, 2225–2229.
- Sheldrick, G. M. (2007). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Song, H. & Shin, H.-S. (1998). *Acta Cryst.* **C54**, 1675–1677.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supporting information

Acta Cryst. (2008). E64, o1496–o1497 [doi:10.1107/S1600536808021326]

1-[2-(4-Fluorobenzoyloxy)-2-phenylethyl]-1*H*-benzimidazole**Özden Özel Güven, Taner Erdoğan, Simon J. Coles and Tuncer Hökelek****S1. Comment**

In recent years, there has been increasing interest in synthesis of heterocyclic compounds having biological and commercial importances. Clotrimazole (Song & Shin, 1998), econazole (Freer *et al.*, 1986), ketoconazole (Peeters *et al.*, 1979*a*) and miconazole (Peeters *et al.*, 1979*b*) are well known imidazole ring containing, while itraconazole (Peeters *et al.*, 1996) and fluconazole (Caira *et al.*, 2004) are 1*H*-1,2,4-triazole ring containing,azole derivatives. They have been developed for clinical uses as antifungal agents (Brammer & Feczko, 1988). Lately, similar structures to miconazole and econazole have been reported to show antibacterial activity more than antifungal activity (Özel Güven *et al.*, 2007*a,b*). In these structures, benzimidazole ring has been found in place of the imidazole ring of miconazole and econazole. We report herein the crystal structure of title benzimidazole derivative.

The asymmetric unit of the title compound (Fig. 1) contains two independent molecules, in which the bond lengths and angles are generally within normal ranges. The planar benzimidazole ring systems are oriented with respect to the phenyl and fluorobenzene rings at dihedral angles of 31.10 (4)°, 45.17 (5)° and 45.52 (5)°, 68.63 (5)° for unprimed and primed molecules, respectively. Atoms C8, C9, C16 and C8', C9', C16' are -0.105 (2), 0.065 (2), -0.060 (2) and -0.163 (2), -0.088 (2), 0.009 (2) Å away from the ring planes of the corresponding benzimidazole, phenyl and fluorobenzene, respectively. So, they are nearly coplanar with the attached rings.

In the crystal structure, intramolecular C—H···N and intermolecular C—H···N and C—H···F hydrogen bonds (Table 1) link the molecules into a three dimensional network (Fig. 2), in which they may be effective in the stabilization of the structure. The C—H··· π contacts (Table 1) between the benzimidazole and the fluorobenzene rings and a π - π contact between the benzimidazole and phenyl ring systems $Cg1 \cdots Cg6^i$ [symmetry code: (i) 1 - x , 1/2 + y , 1/2 - z , where $Cg1$ and $Cg6$ are centroids of the rings (N1/N2/C1/C2/C7) and (C10'-C15'), respectively] further stabilize the structure, with centroid-centroid distance of 4.575 (1) Å.

S2. Experimental

The title compound was synthesized by the reaction of 2-(1*H*-benzimidazol-1-yl) -1-phenylethanol (Özel Güven *et al.*, 2007*a*) with NaH and appropriate benzyl halide. A solution of alcohol (300 mg, 1.259 mmol) in DMF (2.4 ml) was added to NaH (63 mg, 1.574 mmol) in small fractions. The appropriate benzyl halide (238 mg, 1.259 mmol) in DMF (1.2 ml) was added dropwise. The mixture was stirred at room temperature for 2 h, and the excess hydride was decomposed with a small amount of methyl alcohol. After evaporation to dryness under reduced pressure, the crude residue was suspended with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate, and then evaporated to dryness. The crude residue was purified by chromatography on a silica-gel column using chloroform-methanol as eluent. Crystals suitable for X-ray analysis were obtained by the recrystallization of the ether from a mixture of hexane/ethyl acetate (1:2) (yield; 364 mg, 83%).

S3. Refinement

H atoms were located in difference syntheses and refined isotropically [$C-H = 0.94(2)–1.05(2) \text{ \AA}$; $U_{iso}(H) = 0.023(5)–0.051(6) \text{ \AA}^2$].




Figure 1

The molecular structure of the title molecule with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bond is shown as double dashed line.




Figure 2

A packing diagram of (I). Hydrogen bonds are shown as dashed lines.

1-[2-(4-Fluorobenzyloxy)-2-phenylethyl]-1H-benzimidazole

Crystal data

C₂₂H₁₉FN₂O $M_r = 346.39$ Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

 $a = 12.6946$ (2) Å $b = 18.1279$ (4) Å $c = 15.3547$ (3) Å $\beta = 95.747$ (1)° $V = 3515.76$ (12) Å³ $Z = 8$ $F(000) = 1456$ $D_x = 1.309$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8094 reflections

 $\theta = 2.9$ – 27.5 ° $\mu = 0.09$ mm⁻¹ $T = 120$ K

Rod-shaped, colorless

 $0.55 \times 0.15 \times 0.14$ mm

Data collection

Bruker–Nonius KappaCCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 9.091 pixels mm⁻¹ φ and ω scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 2007)

 $T_{\min} = 0.953$, $T_{\max} = 0.988$

41740 measured reflections

8050 independent reflections

5679 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.065$ $\theta_{\max} = 27.5$ °, $\theta_{\min} = 3.0$ ° $h = -16$ → 14 $k = -23$ → 23 $l = -19$ → 19

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.055$ $wR(F^2) = 0.142$ $S = 1.07$

8050 reflections

622 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from

neighbouring sites

All H-atom parameters refined

 $w = 1/[\sigma^2(F_o^2) + (0.0671P)^2 + 0.7368P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.43$ e Å⁻³ $\Delta\rho_{\min} = -0.43$ e Å⁻³

Extinction correction: SHELXL97 (Sheldrick,

2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0237 (14)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F	1.32492 (8)	0.12305 (7)	1.08865 (8)	0.0437 (3)
O	0.89904 (9)	0.13207 (7)	0.84616 (8)	0.0271 (3)

N1	0.83956 (11)	0.19499 (8)	0.67743 (9)	0.0249 (3)
N2	0.98761 (11)	0.20522 (9)	0.61001 (10)	0.0288 (4)
C1	0.94433 (14)	0.21182 (11)	0.68318 (12)	0.0289 (4)
H1	0.9829 (14)	0.2262 (11)	0.7390 (12)	0.027 (5)*
C2	0.90500 (13)	0.18153 (9)	0.54985 (11)	0.0236 (4)
C3	0.90444 (14)	0.16519 (10)	0.46108 (12)	0.0280 (4)
H3	0.9686 (16)	0.1703 (12)	0.4301 (13)	0.038 (6)*
C4	0.81070 (14)	0.14160 (10)	0.41732 (12)	0.0292 (4)
H4	0.8070 (16)	0.1292 (12)	0.3561 (15)	0.039 (6)*
C5	0.71803 (14)	0.13502 (10)	0.45929 (12)	0.0283 (4)
H5	0.6542 (15)	0.1181 (11)	0.4275 (13)	0.030 (5)*
C6	0.71679 (13)	0.15155 (10)	0.54705 (12)	0.0255 (4)
H6	0.6538 (16)	0.1488 (12)	0.5780 (13)	0.036 (5)*
C7	0.81188 (13)	0.17443 (9)	0.59110 (11)	0.0227 (4)
C8	0.77179 (14)	0.19141 (11)	0.74888 (12)	0.0274 (4)
H81	0.7859 (15)	0.2378 (13)	0.7864 (13)	0.037 (6)*
H82	0.6946 (16)	0.1908 (11)	0.7251 (13)	0.032 (5)*
C9	0.79393 (13)	0.12350 (10)	0.80559 (11)	0.0257 (4)
H9	0.7892 (14)	0.0767 (12)	0.7672 (13)	0.030 (5)*
C10	0.71281 (13)	0.11680 (10)	0.87108 (11)	0.0239 (4)
C11	0.63294 (14)	0.06463 (11)	0.85919 (12)	0.0275 (4)
H11	0.6307 (14)	0.0317 (12)	0.8102 (13)	0.030 (5)*
C12	0.55543 (14)	0.06074 (11)	0.91698 (12)	0.0294 (4)
H12	0.5013 (16)	0.0250 (12)	0.9071 (13)	0.036 (5)*
C13	0.55847 (14)	0.10849 (11)	0.98734 (12)	0.0302 (4)
H13	0.5027 (16)	0.1064 (11)	1.0291 (13)	0.034 (5)*
C14	0.63851 (14)	0.16059 (11)	0.99970 (12)	0.0300 (4)
H14	0.6394 (16)	0.1942 (13)	1.0485 (14)	0.042 (6)*
C15	0.71528 (14)	0.16487 (10)	0.94216 (12)	0.0277 (4)
H15	0.7732 (15)	0.2003 (11)	0.9511 (12)	0.028 (5)*
C16	0.93679 (14)	0.06654 (11)	0.89152 (13)	0.0294 (4)
H161	0.9424 (17)	0.0256 (14)	0.8473 (15)	0.048 (6)*
H162	0.8831 (15)	0.0495 (11)	0.9330 (13)	0.030 (5)*
C17	1.04175 (13)	0.08265 (10)	0.94178 (11)	0.0259 (4)
C18	1.05046 (14)	0.13885 (11)	1.00380 (12)	0.0301 (4)
H18	0.9877 (18)	0.1682 (13)	1.0149 (14)	0.048 (6)*
C19	1.14550 (15)	0.15275 (11)	1.05316 (13)	0.0323 (4)
H19	1.1513 (16)	0.1897 (13)	1.0964 (14)	0.038 (6)*
C20	1.23161 (14)	0.10960 (11)	1.03963 (12)	0.0297 (4)
C21	1.22666 (15)	0.05391 (11)	0.97882 (12)	0.0317 (4)
H21	1.2875 (16)	0.0263 (12)	0.9714 (13)	0.037 (6)*
C22	1.13085 (14)	0.04091 (11)	0.92987 (12)	0.0295 (4)
H22	1.1258 (15)	-0.0001 (12)	0.8881 (14)	0.036 (5)*
F'	0.72117 (9)	0.07392 (7)	0.18637 (7)	0.0422 (3)
O'	0.35374 (9)	0.09943 (7)	0.42429 (9)	0.0319 (3)
N1'	0.32870 (11)	0.19095 (8)	0.57339 (9)	0.0250 (3)
N2'	0.49161 (11)	0.21227 (8)	0.63993 (10)	0.0274 (3)
C1'	0.43292 (13)	0.20470 (10)	0.56517 (12)	0.0265 (4)

H1A	0.4581 (15)	0.2084 (11)	0.5054 (13)	0.033 (5)*
C2'	0.42077 (13)	0.20278 (9)	0.70299 (12)	0.0242 (4)
C3'	0.43907 (14)	0.20280 (10)	0.79409 (12)	0.0277 (4)
H3A	0.5121 (16)	0.2104 (11)	0.8237 (13)	0.035 (5)*
C4'	0.35328 (14)	0.19125 (11)	0.84097 (12)	0.0290 (4)
H4A	0.3623 (15)	0.1912 (11)	0.9025 (14)	0.034 (5)*
C5'	0.25116 (14)	0.18014 (10)	0.79938 (12)	0.0284 (4)
H5A	0.1897 (14)	0.1719 (10)	0.8350 (12)	0.023 (5)*
C6'	0.23161 (13)	0.17905 (10)	0.70912 (12)	0.0252 (4)
H6A	0.1612 (15)	0.1685 (11)	0.6825 (12)	0.029 (5)*
C7'	0.31895 (13)	0.18983 (9)	0.66247 (11)	0.0234 (4)
C8'	0.24572 (14)	0.17291 (10)	0.50377 (12)	0.0267 (4)
H8A	0.2466 (15)	0.2088 (12)	0.4553 (13)	0.034 (5)*
H8B	0.1797 (16)	0.1747 (12)	0.5281 (13)	0.034 (5)*
C9'	0.25876 (14)	0.09658 (10)	0.46651 (12)	0.0282 (4)
H9A	0.2690 (15)	0.0595 (12)	0.5175 (13)	0.033 (5)*
C10'	0.16396 (13)	0.07542 (10)	0.40381 (12)	0.0278 (4)
C11'	0.10205 (14)	0.01572 (11)	0.42324 (13)	0.0311 (4)
H11A	0.1273 (17)	-0.0142 (13)	0.4771 (15)	0.046 (6)*
C12'	0.01201 (15)	-0.00234 (12)	0.36915 (13)	0.0356 (5)
H12A	-0.0308 (18)	-0.0463 (14)	0.3827 (15)	0.051 (6)*
C13'	-0.01693 (15)	0.03929 (12)	0.29548 (13)	0.0367 (5)
H13A	-0.0804 (18)	0.0252 (13)	0.2610 (14)	0.045 (6)*
C14'	0.04507 (16)	0.09780 (12)	0.27391 (13)	0.0380 (5)
H14A	0.0274 (17)	0.1268 (13)	0.2198 (15)	0.046 (6)*
C15'	0.13594 (16)	0.11604 (12)	0.32758 (13)	0.0334 (4)
H15A	0.1795 (16)	0.1575 (13)	0.3127 (14)	0.042 (6)*
C16'	0.38651 (15)	0.02982 (11)	0.39365 (14)	0.0313 (4)
H16A	0.3236 (17)	0.0054 (12)	0.3565 (14)	0.041 (6)*
H16B	0.4113 (16)	-0.0028 (13)	0.4487 (14)	0.041 (6)*
C17'	0.47598 (13)	0.04194 (10)	0.33826 (11)	0.0263 (4)
C18'	0.47972 (15)	0.10525 (11)	0.28687 (12)	0.0295 (4)
H18A	0.4234 (15)	0.1408 (12)	0.2871 (13)	0.031 (5)*
C19'	0.56238 (15)	0.11670 (11)	0.23553 (12)	0.0316 (4)
H19A	0.5682 (16)	0.1634 (13)	0.2001 (14)	0.040 (6)*
C20'	0.63892 (14)	0.06312 (11)	0.23568 (12)	0.0295 (4)
C21'	0.63728 (15)	-0.00091 (11)	0.28360 (13)	0.0330 (4)
H21A	0.6947 (18)	-0.0365 (14)	0.2804 (14)	0.049 (6)*
C22'	0.55530 (14)	-0.01061 (11)	0.33574 (12)	0.0300 (4)
H22A	0.5502 (16)	-0.0561 (13)	0.3701 (14)	0.039 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F	0.0279 (6)	0.0494 (8)	0.0519 (7)	-0.0033 (5)	-0.0053 (5)	0.0001 (6)
O	0.0248 (6)	0.0294 (7)	0.0266 (6)	0.0005 (5)	0.0007 (5)	0.0048 (5)
N1	0.0223 (7)	0.0297 (8)	0.0225 (7)	-0.0015 (6)	0.0014 (6)	0.0024 (6)
N2	0.0240 (7)	0.0334 (9)	0.0287 (8)	-0.0050 (6)	0.0007 (6)	0.0060 (7)

C1	0.0260 (9)	0.0331 (11)	0.0269 (10)	-0.0049 (7)	-0.0012 (8)	0.0041 (8)
C2	0.0214 (8)	0.0222 (9)	0.0272 (9)	-0.0001 (7)	0.0019 (7)	0.0047 (7)
C3	0.0272 (9)	0.0295 (10)	0.0281 (10)	0.0026 (8)	0.0073 (7)	0.0052 (8)
C4	0.0342 (10)	0.0290 (10)	0.0244 (9)	0.0038 (8)	0.0024 (7)	-0.0012 (8)
C5	0.0266 (9)	0.0277 (10)	0.0295 (10)	-0.0008 (7)	-0.0022 (7)	-0.0001 (8)
C6	0.0223 (8)	0.0254 (9)	0.0286 (9)	-0.0003 (7)	0.0025 (7)	0.0022 (7)
C7	0.0248 (8)	0.0220 (9)	0.0212 (8)	0.0021 (7)	0.0017 (7)	0.0027 (7)
C8	0.0262 (9)	0.0321 (10)	0.0241 (9)	0.0010 (8)	0.0045 (7)	0.0021 (8)
C9	0.0237 (9)	0.0295 (10)	0.0235 (9)	0.0007 (7)	0.0013 (7)	0.0003 (8)
C10	0.0223 (8)	0.0270 (9)	0.0221 (9)	0.0022 (7)	0.0010 (6)	0.0026 (7)
C11	0.0292 (9)	0.0285 (10)	0.0242 (9)	-0.0014 (7)	-0.0004 (7)	0.0001 (8)
C12	0.0238 (9)	0.0318 (10)	0.0318 (10)	-0.0051 (8)	-0.0010 (7)	0.0077 (8)
C13	0.0271 (9)	0.0342 (11)	0.0297 (10)	0.0036 (8)	0.0054 (8)	0.0082 (8)
C14	0.0323 (10)	0.0306 (10)	0.0279 (10)	0.0013 (8)	0.0069 (8)	0.0010 (8)
C15	0.0285 (9)	0.0269 (10)	0.0277 (9)	-0.0036 (8)	0.0026 (7)	0.0003 (8)
C16	0.0293 (9)	0.0268 (10)	0.0315 (10)	0.0040 (7)	0.0006 (8)	0.0030 (8)
C17	0.0269 (9)	0.0259 (9)	0.0252 (9)	0.0017 (7)	0.0045 (7)	0.0053 (7)
C18	0.0280 (9)	0.0291 (10)	0.0340 (10)	0.0032 (8)	0.0065 (8)	-0.0001 (8)
C19	0.0344 (10)	0.0299 (11)	0.0326 (10)	-0.0023 (8)	0.0029 (8)	-0.0018 (9)
C20	0.0236 (9)	0.0330 (10)	0.0319 (10)	-0.0035 (8)	-0.0003 (7)	0.0071 (8)
C21	0.0285 (9)	0.0352 (11)	0.0322 (10)	0.0083 (8)	0.0072 (8)	0.0070 (8)
C22	0.0343 (10)	0.0318 (10)	0.0224 (9)	0.0069 (8)	0.0039 (7)	0.0034 (8)
F'	0.0391 (6)	0.0524 (8)	0.0372 (6)	0.0111 (5)	0.0142 (5)	0.0048 (6)
O'	0.0283 (7)	0.0285 (7)	0.0400 (8)	-0.0007 (5)	0.0087 (6)	-0.0077 (6)
N1'	0.0237 (7)	0.0277 (8)	0.0237 (8)	-0.0014 (6)	0.0032 (6)	-0.0005 (6)
N2'	0.0243 (7)	0.0276 (8)	0.0307 (8)	-0.0024 (6)	0.0043 (6)	-0.0004 (6)
C1'	0.0255 (9)	0.0256 (9)	0.0289 (10)	-0.0011 (7)	0.0056 (7)	-0.0002 (7)
C2'	0.0235 (8)	0.0197 (9)	0.0296 (9)	0.0003 (7)	0.0034 (7)	-0.0005 (7)
C3'	0.0257 (9)	0.0253 (10)	0.0310 (10)	0.0012 (7)	-0.0018 (7)	-0.0015 (8)
C4'	0.0329 (10)	0.0288 (10)	0.0251 (10)	0.0055 (8)	0.0020 (8)	0.0003 (8)
C5'	0.0282 (9)	0.0269 (10)	0.0310 (10)	0.0029 (7)	0.0077 (8)	0.0018 (8)
C6'	0.0216 (8)	0.0241 (9)	0.0298 (10)	0.0009 (7)	0.0030 (7)	-0.0016 (7)
C7'	0.0250 (8)	0.0191 (8)	0.0261 (9)	0.0026 (7)	0.0020 (7)	-0.0005 (7)
C8'	0.0249 (9)	0.0305 (10)	0.0244 (9)	-0.0005 (7)	0.0007 (7)	-0.0009 (8)
C9'	0.0256 (9)	0.0286 (10)	0.0308 (10)	-0.0014 (7)	0.0041 (7)	0.0003 (8)
C10'	0.0265 (9)	0.0299 (10)	0.0269 (9)	0.0031 (7)	0.0023 (7)	-0.0064 (8)
C11'	0.0317 (10)	0.0282 (10)	0.0333 (10)	0.0015 (8)	0.0025 (8)	-0.0043 (8)
C12'	0.0332 (10)	0.0344 (11)	0.0399 (11)	-0.0034 (9)	0.0060 (8)	-0.0106 (9)
C13'	0.0291 (10)	0.0464 (13)	0.0337 (11)	0.0032 (9)	-0.0011 (8)	-0.0144 (9)
C14'	0.0444 (12)	0.0448 (13)	0.0241 (10)	0.0092 (10)	0.0001 (8)	-0.0034 (9)
C15'	0.0384 (11)	0.0324 (11)	0.0302 (10)	-0.0002 (9)	0.0074 (8)	-0.0035 (8)
C16'	0.0297 (9)	0.0264 (10)	0.0381 (11)	0.0013 (8)	0.0049 (8)	-0.0026 (8)
C17'	0.0263 (9)	0.0253 (9)	0.0265 (9)	0.0006 (7)	-0.0011 (7)	-0.0049 (7)
C18'	0.0295 (9)	0.0300 (10)	0.0286 (10)	0.0074 (8)	0.0011 (7)	-0.0026 (8)
C19'	0.0365 (10)	0.0316 (11)	0.0265 (10)	0.0066 (8)	0.0032 (8)	0.0013 (8)
C20'	0.0283 (9)	0.0362 (11)	0.0245 (9)	0.0031 (8)	0.0048 (7)	-0.0029 (8)
C21'	0.0323 (10)	0.0306 (11)	0.0359 (11)	0.0080 (8)	0.0028 (8)	-0.0049 (8)
C22'	0.0336 (10)	0.0237 (10)	0.0323 (10)	0.0027 (8)	0.0006 (8)	-0.0013 (8)

Geometric parameters (Å, °)

F—C20	1.360 (2)	F'—C20'	1.364 (2)
O—C9	1.423 (2)	O'—C9'	1.426 (2)
O—C16	1.435 (2)	O'—C16'	1.423 (2)
N1—C1	1.359 (2)	N1'—C1'	1.365 (2)
N1—C7	1.388 (2)	N1'—C7'	1.386 (2)
N1—C8	1.462 (2)	N1'—C8'	1.461 (2)
N2—C1	1.305 (2)	N2'—C1'	1.312 (2)
N2—C2	1.394 (2)	N2'—C2'	1.397 (2)
C1—H1	0.980 (19)	C1'—H1A	1.00 (2)
C2—C3	1.394 (3)	C2'—C7'	1.397 (2)
C2—C7	1.402 (2)	C3'—C2'	1.395 (3)
C3—H3	0.99 (2)	C3'—C4'	1.380 (3)
C4—C3	1.375 (3)	C3'—H3A	1.00 (2)
C4—C5	1.402 (3)	C4'—C5'	1.401 (3)
C4—H4	0.96 (2)	C4'—H4A	0.94 (2)
C5—H5	0.95 (2)	C5'—C6'	1.383 (3)
C6—C5	1.382 (3)	C5'—H5A	1.008 (18)
C6—H6	0.97 (2)	C6'—H6A	0.964 (19)
C7—C6	1.387 (2)	C7'—C6'	1.393 (2)
C8—H81	1.03 (2)	C8'—C9'	1.513 (3)
C8—H82	1.011 (19)	C8'—H8A	0.99 (2)
C9—C8	1.518 (3)	C8'—H8B	0.95 (2)
C9—C10	1.514 (2)	C9'—C10'	1.513 (2)
C9—H9	1.03 (2)	C9'—H9A	1.03 (2)
C10—C15	1.395 (3)	C10'—C11'	1.387 (3)
C11—C10	1.385 (3)	C10'—C15'	1.398 (3)
C11—C12	1.391 (3)	C11'—C12'	1.383 (3)
C11—H11	0.96 (2)	C11'—H11A	1.01 (2)
C12—C13	1.382 (3)	C12'—H12A	1.00 (2)
C12—H12	0.95 (2)	C13'—C12'	1.379 (3)
C13—H13	1.00 (2)	C13'—C14'	1.381 (3)
C14—C13	1.386 (3)	C13'—H13A	0.95 (2)
C14—C15	1.382 (3)	C14'—H14A	0.99 (2)
C14—H14	0.97 (2)	C15'—C14'	1.389 (3)
C15—H15	0.97 (2)	C15'—H15A	0.97 (2)
C16—H161	1.01 (2)	C16'—H16A	1.03 (2)
C16—H162	1.03 (2)	C16'—H16B	1.05 (2)
C17—C16	1.500 (2)	C17'—C16'	1.502 (3)
C17—C18	1.391 (3)	C17'—C18'	1.396 (3)
C17—C22	1.388 (2)	C17'—C22'	1.389 (3)
C18—C19	1.382 (3)	C18'—C19'	1.390 (3)
C18—H18	0.99 (2)	C18'—H18A	0.96 (2)
C19—H19	0.94 (2)	C19'—H19A	1.01 (2)
C20—C19	1.377 (3)	C20'—C19'	1.374 (3)
C20—C21	1.372 (3)	C20'—C21'	1.376 (3)
C21—C22	1.384 (3)	C21'—C22'	1.386 (3)

C21—H21	0.94 (2)	C21'—H21A	0.98 (2)
C22—H22	0.98 (2)	C22'—H22A	0.98 (2)
C9—O—C16	112.02 (13)	C16'—O'—C9'	113.99 (14)
C1—N1—C7	105.96 (14)	C1'—N1'—C7'	106.09 (14)
C1—N1—C8	127.37 (15)	C1'—N1'—C8'	127.44 (15)
C7—N1—C8	126.37 (14)	C7'—N1'—C8'	126.15 (14)
C1—N2—C2	104.19 (14)	C1'—N2'—C2'	104.19 (14)
N1—C1—H1	121.1 (11)	N1'—C1'—H1A	119.8 (11)
N2—C1—N1	114.66 (16)	N2'—C1'—N1'	114.16 (16)
N2—C1—H1	124.2 (11)	N2'—C1'—H1A	126.0 (11)
N2—C2—C7	109.96 (15)	N2'—C2'—C7'	110.07 (15)
C3—C2—N2	130.02 (16)	C3'—C2'—N2'	129.88 (16)
C3—C2—C7	120.02 (15)	C3'—C2'—C7'	120.03 (16)
C2—C3—H3	121.8 (12)	C2'—C3'—H3A	120.6 (11)
C4—C3—C2	117.60 (16)	C4'—C3'—C2'	117.52 (16)
C4—C3—H3	120.6 (12)	C4'—C3'—H3A	121.8 (11)
C3—C4—C5	121.79 (17)	C3'—C4'—C5'	121.77 (17)
C3—C4—H4	120.3 (12)	C3'—C4'—H4A	120.1 (12)
C5—C4—H4	117.9 (12)	C5'—C4'—H4A	118.2 (12)
C4—C5—H5	120.1 (12)	C4'—C5'—H5A	120.3 (10)
C6—C5—C4	121.45 (16)	C6'—C5'—C4'	121.58 (17)
C6—C5—H5	118.5 (12)	C6'—C5'—H5A	118.1 (10)
C5—C6—C7	116.49 (16)	C5'—C6'—C7'	116.17 (16)
C5—C6—H6	123.8 (12)	C5'—C6'—H6A	119.6 (11)
C7—C6—H6	119.7 (12)	C7'—C6'—H6A	124.2 (11)
N1—C7—C2	105.22 (14)	N1'—C7'—C6'	131.61 (15)
C6—C7—N1	132.15 (16)	N1'—C7'—C2'	105.49 (14)
C6—C7—C2	122.63 (16)	C6'—C7'—C2'	122.90 (16)
N1—C8—C9	112.17 (15)	N1'—C8'—C9'	112.54 (15)
N1—C8—H81	107.6 (11)	N1'—C8'—H8A	110.0 (11)
N1—C8—H82	110.6 (11)	N1'—C8'—H8B	107.7 (12)
C9—C8—H81	109.3 (11)	C9'—C8'—H8A	107.9 (12)
C9—C8—H82	108.4 (12)	C9'—C8'—H8B	108.2 (13)
H81—C8—H82	108.7 (16)	H8A—C8'—H8B	110.5 (17)
O—C9—C8	105.91 (14)	O'—C9'—C8'	105.57 (14)
O—C9—C10	112.84 (14)	O'—C9'—C10'	112.01 (15)
O—C9—H9	109.7 (10)	O'—C9'—H9A	108.7 (11)
C8—C9—H9	110.0 (11)	C8'—C9'—C10'	111.24 (15)
C10—C9—C8	110.29 (14)	C8'—C9'—H9A	108.7 (11)
C10—C9—H9	108.0 (11)	C10'—C9'—H9A	110.4 (11)
C11—C10—C9	120.31 (16)	C11'—C10'—C15'	119.25 (17)
C11—C10—C15	119.24 (16)	C11'—C10'—C9'	119.50 (17)
C15—C10—C9	120.40 (16)	C15'—C10'—C9'	121.22 (17)
C10—C11—C12	120.38 (17)	C10'—C11'—H11A	117.0 (13)
C10—C11—H11	119.5 (12)	C12'—C11'—C10'	120.49 (19)
C12—C11—H11	120.1 (12)	C12'—C11'—H11A	122.5 (13)
C11—C12—H12	118.8 (13)	C11'—C12'—H12A	119.9 (13)

C13—C12—C11	120.05 (17)	C13'—C12'—C11'	120.0 (2)
C13—C12—H12	121.2 (13)	C13'—C12'—H12A	120.1 (13)
C12—C13—C14	119.77 (17)	C12'—C13'—C14'	120.38 (18)
C12—C13—H13	120.6 (12)	C12'—C13'—H13A	116.8 (14)
C14—C13—H13	119.6 (12)	C14'—C13'—H13A	122.8 (14)
C13—C14—H14	119.2 (13)	C13'—C14'—C15'	120.01 (19)
C15—C14—C13	120.33 (18)	C13'—C14'—H14A	121.5 (13)
C15—C14—H14	120.5 (13)	C15'—C14'—H14A	118.5 (13)
C10—C15—H15	118.6 (11)	C10'—C15'—H15A	120.0 (13)
C14—C15—C10	120.22 (17)	C14'—C15'—C10'	119.85 (19)
C14—C15—H15	121.2 (11)	C14'—C15'—H15A	120.1 (12)
O—C16—C17	108.96 (15)	O'—C16'—C17'	108.67 (15)
O—C16—H161	108.9 (13)	O'—C16'—H16A	109.1 (12)
O—C16—H162	109.9 (11)	O'—C16'—H16B	107.8 (12)
C17—C16—H162	110.5 (10)	C17'—C16'—H16A	109.7 (12)
C17—C16—H161	111.6 (12)	C17'—C16'—H16B	110.8 (11)
H162—C16—H161	106.9 (17)	H16A—C16'—H16B	110.7 (17)
C18—C17—C16	120.02 (16)	C18'—C17'—C16'	120.72 (16)
C22—C17—C18	118.64 (17)	C22'—C17'—C18'	118.82 (17)
C22—C17—C16	121.30 (17)	C22'—C17'—C16'	120.45 (17)
C17—C18—H18	120.2 (13)	C17'—C18'—H18A	118.8 (12)
C19—C18—C17	120.93 (17)	C19'—C18'—C17'	120.91 (17)
C19—C18—H18	118.8 (13)	C19'—C18'—H18A	120.3 (12)
C18—C19—H19	121.3 (13)	C18'—C19'—H19A	121.9 (12)
C20—C19—C18	118.45 (19)	C20'—C19'—C18'	117.87 (18)
C20—C19—H19	120.2 (13)	C20'—C19'—H19A	120.2 (12)
F—C20—C19	118.38 (17)	F'—C20'—C19'	118.57 (17)
F—C20—C21	119.09 (16)	F'—C20'—C21'	118.13 (16)
C21—C20—C19	122.53 (17)	C19'—C20'—C21'	123.30 (18)
C20—C21—C22	118.19 (17)	C20'—C21'—C22'	117.92 (18)
C20—C21—H21	119.6 (13)	C20'—C21'—H21A	118.6 (14)
C22—C21—H21	122.2 (13)	C22'—C21'—H21A	123.5 (14)
C17—C22—H22	119.9 (12)	C17'—C22'—H22A	118.2 (12)
C21—C22—C17	121.26 (18)	C21'—C22'—C17'	121.14 (18)
C21—C22—H22	118.8 (12)	C21'—C22'—H22A	120.6 (12)
C16—O—C9—C8	-171.09 (14)	C16'—O'—C9'—C8'	172.99 (15)
C16—O—C9—C10	68.15 (19)	C16'—O'—C9'—C10'	-65.8 (2)
C9—O—C16—C17	-172.87 (14)	C9'—O'—C16'—C17'	170.86 (14)
C7—N1—C1—N2	0.7 (2)	C7'—N1'—C1'—N2'	0.3 (2)
C8—N1—C1—N2	174.73 (17)	C8'—N1'—C1'—N2'	-173.60 (17)
C1—N1—C7—C2	-0.70 (18)	C1'—N1'—C7'—C6'	179.34 (18)
C1—N1—C7—C6	179.78 (19)	C8'—N1'—C7'—C6'	-6.7 (3)
C8—N1—C7—C2	-174.84 (16)	C1'—N1'—C7'—C2'	-0.44 (18)
C8—N1—C7—C6	5.6 (3)	C8'—N1'—C7'—C2'	173.54 (16)
C1—N1—C8—C9	-73.6 (2)	C1'—N1'—C8'—C9'	70.5 (2)
C7—N1—C8—C9	99.33 (19)	C7'—N1'—C8'—C9'	-102.2 (2)
C2—N2—C1—N1	-0.3 (2)	C2'—N2'—C1'—N1'	0.0 (2)

C1—N2—C2—C3	-179.88 (19)	C1'—N2'—C2'—C3'	177.92 (19)
C1—N2—C2—C7	-0.15 (19)	C1'—N2'—C2'—C7'	-0.30 (19)
N2—C2—C3—C4	179.03 (18)	C3'—C2'—C7'—N1'	-177.96 (16)
C7—C2—C3—C4	-0.7 (3)	N2'—C2'—C7'—N1'	0.47 (19)
N2—C2—C7—N1	0.54 (19)	C3'—C2'—C7'—C6'	2.2 (3)
N2—C2—C7—C6	-179.88 (16)	N2'—C2'—C7'—C6'	-179.33 (16)
C3—C2—C7—N1	-179.70 (16)	C4'—C3'—C2'—N2'	-179.40 (17)
C3—C2—C7—C6	-0.1 (3)	C4'—C3'—C2'—C7'	-1.3 (3)
C5—C4—C3—C2	0.9 (3)	C2'—C3'—C4'—C5'	-0.2 (3)
C3—C4—C5—C6	-0.4 (3)	C3'—C4'—C5'—C6'	1.0 (3)
C7—C6—C5—C4	-0.4 (3)	C4'—C5'—C6'—C7'	-0.2 (3)
N1—C7—C6—C5	-179.90 (18)	N1'—C7'—C6'—C5'	178.82 (18)
C2—C7—C6—C5	0.6 (3)	C2'—C7'—C6'—C5'	-1.4 (3)
O—C9—C8—N1	65.00 (18)	N1'—C8'—C9'—O'	-66.22 (19)
C10—C9—C8—N1	-172.60 (14)	N1'—C8'—C9'—C10'	172.07 (15)
O—C9—C10—C11	-136.59 (17)	O'—C9'—C10'—C11'	125.14 (18)
O—C9—C10—C15	45.8 (2)	C8'—C9'—C10'—C11'	-116.98 (19)
C8—C9—C10—C11	105.18 (19)	O'—C9'—C10'—C15'	-56.5 (2)
C8—C9—C10—C15	-72.4 (2)	C8'—C9'—C10'—C15'	61.4 (2)
C9—C10—C15—C14	177.27 (16)	C9'—C10'—C11'—C12'	176.72 (17)
C11—C10—C15—C14	-0.3 (3)	C15'—C10'—C11'—C12'	-1.7 (3)
C12—C11—C10—C9	-176.91 (16)	C9'—C10'—C15'—C14'	-176.33 (17)
C12—C11—C10—C15	0.7 (3)	C11'—C10'—C15'—C14'	2.1 (3)
C10—C11—C12—C13	-0.8 (3)	C10'—C11'—C12'—C13'	-0.2 (3)
C11—C12—C13—C14	0.5 (3)	C14'—C13'—C12'—C11'	1.8 (3)
C15—C14—C13—C12	-0.1 (3)	C12'—C13'—C14'—C15'	-1.4 (3)
C13—C14—C15—C10	0.1 (3)	C10'—C15'—C14'—C13'	-0.5 (3)
C18—C17—C16—O	57.8 (2)	C18'—C17'—C16'—O'	-33.7 (2)
C22—C17—C16—O	-124.31 (18)	C22'—C17'—C16'—O'	147.83 (17)
C16—C17—C18—C19	177.53 (17)	C16'—C17'—C18'—C19'	179.98 (17)
C22—C17—C18—C19	-0.4 (3)	C22'—C17'—C18'—C19'	-1.5 (3)
C16—C17—C22—C21	-177.36 (17)	C18'—C17'—C22'—C21'	0.2 (3)
C18—C17—C22—C21	0.6 (3)	C16'—C17'—C22'—C21'	178.70 (17)
C17—C18—C19—C20	-0.1 (3)	C17'—C18'—C19'—C20'	1.3 (3)
F—C20—C19—C18	-179.59 (16)	F'—C20'—C19'—C18'	-179.30 (16)
C21—C20—C19—C18	0.4 (3)	C21'—C20'—C19'—C18'	0.3 (3)
F—C20—C21—C22	179.72 (16)	F'—C20'—C21'—C22'	178.03 (16)
C19—C20—C21—C22	-0.3 (3)	C19'—C20'—C21'—C22'	-1.6 (3)
C20—C21—C22—C17	-0.2 (3)	C20'—C21'—C22'—C17'	1.3 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6 \cdots N2'	0.97 (2)	2.62 (2)	3.498 (2)	150.5 (17)
C6'—H6A \cdots N2 ⁱ	0.964 (19)	2.459 (19)	3.345 (2)	152.7 (16)
C13—H13 \cdots F ⁱ	1.00 (2)	2.54 (2)	3.493 (2)	159.6 (16)
C19'—H19A \cdots N2 ⁱⁱ	1.01 (2)	2.59 (2)	3.509 (2)	151.1 (16)

C5—H5...Cg8 ⁱⁱⁱ	0.95 (2)	2.662 (19)	3.514 (2)	149.0 (15)
C22'—H22A...Cg2 ^{iv}	0.98 (2)	2.62 (9)	3.518 (2)	141.9 (16)

Symmetry codes: (i) $x-1, y, z$; (ii) $x, -y+1/2, z-1/2$; (iii) $x, -y-1/2, z-3/2$; (iv) $-x+1, -y, -z+1$.