

(E)-1,2-Bis(4-fluorophenyl)ethane-1,2-dione

Hoong-Kun Fun* and Reza Kia†

X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia
Correspondence e-mail: hkfun@usm.my


Received 22 July 2008; accepted 24 July 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002 \text{ \AA}$; R factor = 0.043; wR factor = 0.107; data-to-parameter ratio = 21.4.

The title compound, $\text{C}_{14}\text{H}_8\text{F}_2\text{O}_2$, is a substituted benzil with an *s-trans* conformation of the dicarbonyl unit. This conformation is also shown by the $\text{O}-\text{C}-\text{C}-\text{O}$ torsion angle of $-110.65 (12)^\circ$. An unusual feature of the structure is the length, $1.536 (2) \text{ \AA}$, of the central $\text{C}-\text{C}$ bond connecting the carbonyl units, which is significantly longer than a normal $\text{Csp}^2-\text{Csp}^2$ single bond. This is probably the result of decreasing the unfavourable vicinal dipole–dipole interactions by increasing the distance between the two electronegative O atoms [$\text{O}\cdots\text{O} = 3.1867 (12) \text{ \AA}$] and allowing orbital overlap of the dione with the π system of the benzene rings. The dihedral angle between the aromatic rings is $64.74 (5)^\circ$. In the crystal structure, neighbouring molecules are linked together by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ ($\times 2$) hydrogen bonds. In addition, the crystal structure is further stabilized by intermolecular $\pi-\pi$ interactions with centroid–centroid distances in the range $3.6416 (6)$ – $3.7150 (7) \text{ \AA}$.

Related literature

For bond-length data, see Allen *et al.* (1987). For carbonyl–carbonyl interaction, see Allen *et al.* (1998). For related structures and applications see, for example: Kaftory & Rubin (1983); Frey *et al.* (1995); Crowley *et al.* (1983); More *et al.* (1987); Brown *et al.* (1965); Gabe *et al.* (1981); Kimura *et al.* (1979); Stevens & Dubois (1962); Shimizu & Bartlett (1976); Rubin (1978).

**Experimental***Crystal data*

$\text{C}_{14}\text{H}_8\text{F}_2\text{O}_2$	$V = 1099.16 (3) \text{ \AA}^3$
$M_r = 246.20$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 12.1351 (2) \text{ \AA}$	$\mu = 0.12 \text{ mm}^{-1}$
$b = 7.3500 (1) \text{ \AA}$	$T = 100.0 (1) \text{ K}$
$c = 13.1572 (2) \text{ \AA}$	$0.39 \times 0.30 \times 0.28 \text{ mm}$
$\beta = 110.507 (1)^\circ$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	14891 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	3489 independent reflections
$T_{\min} = 0.903$, $T_{\max} = 0.967$	2846 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	163 parameters
$wR(F^2) = 0.107$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.39 \text{ e \AA}^{-3}$
3489 reflections	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$

Table 1
Selected centroid–centroid distances (\AA).

$Cg1\cdots Cg1^i$	3.6416 (6)	$Cg2\cdots Cg2^{ii}$	3.7150 (7)
-------------------	------------	----------------------	------------

Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $-x + 2, -y + 2, -z + 1$. $Cg1$ and $Cg2$ are the centroids of the C1–C6 and C9–C14 benzene rings, respectively.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\cdots \text{O}1^{iii}$	0.93	2.40	3.2648 (15)	155
$\text{C}11-\text{H}11\text{A}\cdots \text{O}2^{iv}$	0.93	2.51	3.3098 (16)	145

Symmetry codes: (iii) $x, y - 1, z$; (iv) $x, -y + \frac{5}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2003).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/

‡ Additional correspondence author, e-mail: zsrkk@yahoo.com.

PFIZIK/613312. RK thanks Universiti Sains Malaysia for the award of a post-doctoral research fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2128).

References

- Allen, F. H., Baalham, C. A., Lommerse, J. P. M. & Raithby, P. R. (1998). *Acta Cryst. B* **54**, 320–329.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Brown, C. J. & Sadanaga, R. (1965). *Acta Cryst. A* **18**, 158–164.
- Bruker (2005). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Crowley, J. I., Balanson, R. D. & Mayerle, J. J. (1983). *J. Am. Chem. Soc.* **105**, 6416–6422.
- Frey, J., Faraggi, E., Rappoport, Z. & Kaftory, M. (1995). *J. Chem. Soc. Perkin Trans. 2*, pp. 1745–1748.
- Gabe, E. J., Le Page, Y., Lee, F. L. & Barclay, L. R. C. (1981). *Acta Cryst. B* **37**, 197–200.
- Kaftory, M. & Rubin, M. B. (1983). *J. Chem. Soc. Perkin Trans. 2*, pp. 149–154.
- Kimura, M., McCluney, R. E. & Watson, W. H. (1979). *Acta Cryst. B* **35**, 483–484.
- More, M., Odou, G. & Lefebvre, J. (1987). *Acta Cryst. B* **43**, 398–405.
- Rubin, M. B. (1978). *Chem. Rev.* **78**, 1121–1164.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shimizu, N. & Bartlett, P. D. (1976). *J. Am. Chem. Soc.* **98**, 4193–4200.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Stevens, B. & Dubois, J. T. (1962). *J. Chem. Soc.* pp. 2813–2815.

supporting information

Acta Cryst. (2008). E64, o1617–o1618 [doi:10.1107/S1600536808023350]

(*E*)-1,2-Bis(4-fluorophenyl)ethane-1,2-dione

Hoong-Kun Fun and Reza Kia

S1. Comment

Investigation of photophysical properties of α -dicarbonyls has focused on the intramolecular carbonyl group electronic interaction as a function of their geometrical relationship. In previous extensive studies of the photochemistry of these compounds (Stevens & Dubois, 1962; Shimizu & Bartlett, 1976), biacetyl and benzil were the exclusive experimental vehicles for photophysical study. The structure of vicinal di- and polycarbonyl compounds have been of interest for many years (Rubin, 1978; Crowley *et al.*, 1983; Kaftory *et al.*, 1983; Frey *et al.*, 1995; Kimura *et al.*, 1979). Only a limited amount of data has been gathered from solid-state configurations such as in single crystals or as inclusion dopants in host crystals.


In the title compound (**I**, Fig. 1), bond lengths, bond angles, and torsion angles of the dicarbonyl unit deviate significantly from normal values (Allen *et al.*, 1987) in order to minimize the repulsive interactions resulting from juxtaposition of dipolar carbonyl groups (Allen *et al.*, 1987). The C7–C8 bond distance connecting the carbonyl units is longer than those in normal sp^2 – sp^2 single bonds, such as in butadiene. This is probably the result of decreasing the unfavourable vicinal dipole-dipole interactions. The dicarbonyl unit has an *s-trans* conformation as indicated by the torsion angle of O1–C7–C6–C5, and O2–C8–C9–C10 being -174.32 (11) and 174.49 (11) $^\circ$, respectively. This conformation is substantiated by the torsion angle of O–C–C–O, being -110.65 (12) $^\circ$. The overall effect is to maximize the distance between the two electronegative oxygen atoms [$O1 \cdots O2 = 3.1867$ (12) Å] and to allow orbital overlap of the dione with the π system of the benzene rings. The dihedral angle between two phenyl rings is 64.74 (5) $^\circ$. In the crystal structure, neighbouring molecules are linked together by weak intermolecular C—H \cdots O (2 \times) hydrogen bonds. The packing mode (Fig. 2) tends to be dominated by van der Waals close packing interactions and the preference for aligning the substituted phenyl rings parallel to each other along the *c* axis with centroid to centroid distances of the π rings of 3.6416 (6) to 3.7150 (7) Å.

S2. Experimental


The synthetic method has been described earlier (Frey *et al.*, 1995). Single crystals suitable for X-ray diffraction were obtained by evaporation of a methanol solution at room temperature.

S3. Refinement

All of the hydrogen atoms were positioned geometrically and refined using a riding model with isotropic thermal parameters 1.2 times that of the parent atom.

**Figure 1**

The molecular structure of (I) with atom labels and 50% probability ellipsoids for non-H atoms.

**Figure 2**

The crystal packing, showing parallel aligning of the benzene rings along the *c* axis. Intermolecular C—H···O interactions are shown as dashed lines.

(E)-1,2-Bis(4-fluorophenyl)ethane-1,2-dione

Crystal data

$C_{14}H_8F_2O_2$
 $M_r = 246.20$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 12.1351 (2) \text{ \AA}$
 $b = 7.3500 (1) \text{ \AA}$
 $c = 13.1572 (2) \text{ \AA}$
 $\beta = 110.507 (1)^\circ$

$V = 1099.16 (3) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 504$
 $D_x = 1.488 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 6651 reflections
 $\theta = 3.2\text{--}30.8^\circ$
 $\mu = 0.12 \text{ mm}^{-1}$

$T = 100\text{ K}$
Block, pale-yellow

$0.39 \times 0.30 \times 0.28\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)
 $T_{\min} = 0.903$, $T_{\max} = 0.967$

14891 measured reflections
3489 independent reflections
2846 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\max} = 31.0^\circ$, $\theta_{\min} = 3.2^\circ$
 $h = -15 \rightarrow 17$
 $k = -8 \rightarrow 10$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.107$
 $S = 1.06$
3489 reflections
163 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0458P)^2 + 0.3386P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.39\text{ e }{\text{\AA}}^{-3}$
 $\Delta\rho_{\min} = -0.21\text{ e }{\text{\AA}}^{-3}$

Special details

Experimental. The low-temperature data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.
Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.35237 (6)	0.66325 (10)	0.08245 (6)	0.03021 (18)
F2	1.21050 (6)	1.20663 (11)	0.59266 (6)	0.03299 (19)
O1	0.69635 (7)	1.33900 (11)	0.19570 (7)	0.0273 (2)
O2	0.85192 (7)	1.05531 (12)	0.12153 (7)	0.02583 (19)
C1	0.64396 (10)	0.85311 (15)	0.16653 (9)	0.0200 (2)
H1A	0.7218	0.8152	0.1869	0.024*
C2	0.55476 (10)	0.72510 (16)	0.14036 (9)	0.0217 (2)
H2A	0.5712	0.6012	0.1434	0.026*
C3	0.44035 (10)	0.78766 (16)	0.10962 (9)	0.0215 (2)
C4	0.41085 (10)	0.96950 (17)	0.10511 (9)	0.0223 (2)
H4A	0.3327	1.0060	0.0844	0.027*
C5	0.50092 (10)	1.09585 (16)	0.13226 (8)	0.0202 (2)
H5A	0.4836	1.2193	0.1304	0.024*

C6	0.61811 (9)	1.03855 (15)	0.16255 (8)	0.0183 (2)
C7	0.71259 (9)	1.17601 (15)	0.18983 (9)	0.0201 (2)
C8	0.83778 (9)	1.11508 (14)	0.20254 (9)	0.0195 (2)
C9	0.93465 (9)	1.14174 (14)	0.30709 (9)	0.0186 (2)
C10	0.91694 (10)	1.22467 (15)	0.39561 (9)	0.0206 (2)
H10A	0.8422	1.2644	0.3896	0.025*
C11	1.01017 (11)	1.24806 (15)	0.49231 (10)	0.0231 (2)
H11A	0.9995	1.3047	0.5514	0.028*
C12	1.11907 (10)	1.18496 (16)	0.49832 (9)	0.0230 (2)
C13	1.14030 (10)	1.09925 (16)	0.41347 (10)	0.0228 (2)
H13A	1.2149	1.0563	0.4212	0.027*
C14	1.04700 (9)	1.07948 (15)	0.31680 (9)	0.0204 (2)
H14A	1.0589	1.0245	0.2578	0.025*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0221 (3)	0.0307 (4)	0.0367 (4)	-0.0068 (3)	0.0089 (3)	-0.0029 (3)
F2	0.0255 (4)	0.0401 (5)	0.0272 (4)	-0.0061 (3)	0.0014 (3)	0.0002 (3)
O1	0.0255 (4)	0.0194 (4)	0.0371 (5)	0.0035 (3)	0.0110 (4)	0.0006 (3)
O2	0.0237 (4)	0.0296 (4)	0.0263 (4)	-0.0006 (3)	0.0114 (3)	-0.0042 (3)
C1	0.0174 (5)	0.0227 (5)	0.0199 (5)	0.0039 (4)	0.0064 (4)	0.0032 (4)
C2	0.0225 (5)	0.0202 (5)	0.0227 (5)	0.0026 (4)	0.0082 (4)	0.0026 (4)
C3	0.0189 (5)	0.0256 (6)	0.0200 (5)	-0.0022 (4)	0.0068 (4)	-0.0002 (4)
C4	0.0169 (5)	0.0291 (6)	0.0207 (5)	0.0041 (4)	0.0062 (4)	0.0011 (4)
C5	0.0199 (5)	0.0220 (5)	0.0190 (5)	0.0051 (4)	0.0070 (4)	0.0014 (4)
C6	0.0179 (5)	0.0207 (5)	0.0166 (4)	0.0024 (4)	0.0065 (4)	0.0016 (4)
C7	0.0192 (5)	0.0216 (5)	0.0200 (5)	0.0023 (4)	0.0075 (4)	0.0014 (4)
C8	0.0189 (5)	0.0162 (5)	0.0251 (5)	0.0003 (4)	0.0097 (4)	0.0009 (4)
C9	0.0184 (5)	0.0148 (5)	0.0238 (5)	-0.0002 (4)	0.0089 (4)	0.0014 (4)
C10	0.0199 (5)	0.0183 (5)	0.0264 (5)	0.0001 (4)	0.0118 (4)	0.0011 (4)
C11	0.0272 (6)	0.0203 (5)	0.0243 (5)	-0.0031 (4)	0.0121 (4)	-0.0010 (4)
C12	0.0215 (5)	0.0212 (5)	0.0240 (5)	-0.0046 (4)	0.0050 (4)	0.0033 (4)
C13	0.0177 (5)	0.0213 (5)	0.0302 (6)	0.0005 (4)	0.0093 (4)	0.0045 (4)
C14	0.0207 (5)	0.0172 (5)	0.0263 (5)	0.0013 (4)	0.0118 (4)	0.0020 (4)

Geometric parameters (\AA , $^\circ$)

F1—C3	1.3551 (13)	C6—C7	1.4751 (15)
F2—C12	1.3536 (14)	C7—C8	1.5358 (15)
O1—C7	1.2209 (14)	C8—C9	1.4764 (15)
O2—C8	1.2193 (13)	C9—C10	1.3962 (15)
C1—C2	1.3836 (16)	C9—C14	1.4009 (14)
C1—C6	1.3955 (15)	C10—C11	1.3854 (17)
C1—H1A	0.9300	C10—H10A	0.9300
C2—C3	1.3818 (15)	C11—C12	1.3764 (17)
C2—H2A	0.9300	C11—H11A	0.9300
C3—C4	1.3795 (17)	C12—C13	1.3824 (17)

C4—C5	1.3825 (16)	C13—C14	1.3826 (16)
C4—H4A	0.9300	C13—H13A	0.9300
C5—C6	1.4005 (14)	C14—H14A	0.9300
C5—H5A	0.9300		
Cg1···Cg1 ⁱ	3.6416 (6)	Cg2···Cg2 ⁱⁱ	3.7150 (7)
C2—C1—C6	120.58 (10)	O2—C8—C9	123.53 (10)
C2—C1—H1A	119.7	O2—C8—C7	116.53 (10)
C6—C1—H1A	119.7	C9—C8—C7	119.81 (9)
C3—C2—C1	117.70 (11)	C10—C9—C14	119.72 (10)
C3—C2—H2A	121.1	C10—C9—C8	122.02 (9)
C1—C2—H2A	121.1	C14—C9—C8	118.26 (9)
F1—C3—C4	118.28 (10)	C11—C10—C9	120.35 (10)
F1—C3—C2	118.08 (10)	C11—C10—H10A	119.8
C4—C3—C2	123.64 (11)	C9—C10—H10A	119.8
C3—C4—C5	118.03 (10)	C12—C11—C10	118.01 (10)
C3—C4—H4A	121.0	C12—C11—H11A	121.0
C5—C4—H4A	121.0	C10—C11—H11A	121.0
C4—C5—C6	120.24 (10)	F2—C12—C11	118.37 (11)
C4—C5—H5A	119.9	F2—C12—C13	118.00 (10)
C6—C5—H5A	119.9	C11—C12—C13	123.63 (11)
C1—C6—C5	119.80 (10)	C12—C13—C14	117.81 (10)
C1—C6—C7	120.96 (10)	C12—C13—H13A	121.1
C5—C6—C7	119.24 (10)	C14—C13—H13A	121.1
O1—C7—C6	123.95 (10)	C13—C14—C9	120.44 (10)
O1—C7—C8	117.05 (10)	C13—C14—H14A	119.8
C6—C7—C8	118.80 (9)	C9—C14—H14A	119.8
C6—C1—C2—C3	0.53 (16)	O1—C7—C8—C9	65.39 (14)
C1—C2—C3—F1	179.13 (9)	C6—C7—C8—C9	-119.59 (11)
C1—C2—C3—C4	-0.89 (17)	O2—C8—C9—C10	174.49 (11)
F1—C3—C4—C5	-179.60 (9)	C7—C8—C9—C10	-1.26 (15)
C2—C3—C4—C5	0.41 (17)	O2—C8—C9—C14	-6.17 (16)
C3—C4—C5—C6	0.42 (15)	C7—C8—C9—C14	178.08 (9)
C2—C1—C6—C5	0.26 (16)	C14—C9—C10—C11	0.97 (16)
C2—C1—C6—C7	-179.72 (10)	C8—C9—C10—C11	-179.71 (10)
C4—C5—C6—C1	-0.75 (15)	C9—C10—C11—C12	-0.93 (16)
C4—C5—C6—C7	179.23 (10)	C10—C11—C12—F2	-179.73 (10)
C1—C6—C7—O1	-174.32 (10)	C10—C11—C12—C13	-0.22 (17)
C5—C6—C7—O1	5.70 (16)	F2—C12—C13—C14	-179.18 (10)
C1—C6—C7—C8	11.03 (15)	C11—C12—C13—C14	1.31 (17)
C5—C6—C7—C8	-168.95 (9)	C12—C13—C14—C9	-1.24 (16)
O1—C7—C8—O2	-110.65 (12)	C10—C9—C14—C13	0.15 (16)
C6—C7—C8—O2	64.37 (13)	C8—C9—C14—C13	-179.20 (10)

Symmetry codes: (i) $-x+1, -y+2, -z$; (ii) $-x+2, -y+2, -z+1$.

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C2—H2A···O1 ⁱⁱⁱ	0.93	2.40	3.2648 (15)	155
C11—H11A···O2 ^{iv}	0.93	2.51	3.3098 (16)	145

Symmetry codes: (iii) $x, y-1, z$; (iv) $x, -y+5/2, z+1/2$.