

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-Methyl N'-[1-(2-hydroxyphenyl)ethylidene]hydrazinecarboxylate

Bin Zhu and Xiang-Wei Cheng*

Zhejiang Police College Experience Center, Zhejiang Police College, Hangzhou 310053, People's Republic of China
Correspondence e-mail: zpccxw@126.com


Received 27 July 2008; accepted 30 July 2008

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.036; wR factor = 0.107; data-to-parameter ratio = 12.9.

The molecule of the title compound, $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_3$, adopts a *trans* configuration with respect to the $\text{C}=\text{N}$ bond. The dihedral angle between the benzene ring and the hydrazinecarboxylate plane is $8.98(7)^\circ$. Intramolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds are observed. Molecules are linked into chains along the c axis by $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds. In addition, $\text{C}-\text{H}\cdots\pi$ interactions are observed.

Related literature

For general background, see: Parashar *et al.* (1988); Hadjoudis *et al.* (1987); Borg *et al.* (1999). For a related structure, see: Cheng (2008).



Experimental

Crystal data

$\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_3$
 $M_r = 208.22$
Monoclinic, $P2_1/c$
 $a = 8.6432(8)$ Å

$b = 12.6696(11)$ Å
 $c = 9.9810(9)$ Å
 $\beta = 109.837(3)^\circ$
 $V = 1028.12(16)$ Å³

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 273(2)$ K
 $0.28 \times 0.24 \times 0.23$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2002)
 $T_{\min} = 0.973$, $T_{\max} = 0.979$

10601 measured reflections
1809 independent reflections
1587 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.019$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.107$
 $S = 1.05$
1809 reflections

140 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.26$ e Å⁻³
 $\Delta\rho_{\min} = -0.12$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{N1}$	0.82	1.85	2.5625 (14)	145
$\text{N2}-\text{H2A}\cdots\text{O2}^i$	0.86	2.25	3.0550 (14)	156
$\text{C8}-\text{H8A}\cdots\text{N2}$	0.96	2.47	2.820 (2)	101
$\text{C8}-\text{H8C}\cdots\text{Cg1}^{ii}$	0.96	2.93	3.803 (2)	151

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x + 1, -y + 1, -z + 1$. Cg1 is the centroid of the $\text{C1}-\text{C6}$ ring.

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors acknowledge financial support from Zhejiang Police College, China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2645).

References

- Borg, S., Vollinga, R. C., Labarre, M., Payza, K., Terenius, L. & Luthman, K. (1999). *J. Med. Chem.* **42**, 4331–4342.
Bruker (2002). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Cheng, X.-W. (2008). *Acta Cryst.* **E64**, o1384.
Hadjoudis, E., Vittorakis, M. & Moustakali-Mavridis, J. (1987). *Tetrahedron*, **43**, 1345–1360.
Parashar, R. K., Sharma, R. C., Kumar, A. & Mohanm, G. (1988). *Inorg. Chim. Acta*, **151**, 201–208.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o1748 [doi:10.1107/S1600536808024434]

(*E*)-Methyl *N'*-[1-(2-hydroxyphenyl)ethylidene]hydrazinecarboxylate

B. Zhu and X.-W. Cheng

Comment

Benzaldehydehydrazone compounds have received considerable attention for a long time due to their pharmacological activity (Parashar *et al.*, 1988) and their photochromic properties (Hadjoudis *et al.*, 1987). They are important intermediates of 1,3,4-oxadiazoles, which have been reported to be versatile compounds with many properties (Borg *et al.*, 1999). As a further investigation of this type of derivatives, the crystal structure of the title compound is reported here.

The title molecule (Fig.1) adopts a *trans* configuration with respect to the C=N bond. The C9/C10/N1/N2/O2/O3 plane of the hydrazine carboxylic acid methyl ester group is slightly twisted away from the attached ring. The dihedral angle between the C1–C6 ring and the C9/C10/N1/N2/O2/O3 plane is 8.98 (7)°. The bond lengths and angles agree with those observed for methyl *N'*-((*E*)-1-phenylethylidene)hydrazinecarboxylate (Cheng, 2008). Intramolecular O—H···N and C—H···N hydrogen bonds are observed.

The molecules are linked into chains along the *c* axis by N—H···O hydrogen bonds. In addition, C—H··· π interactions are observed (Table 1, Fig.2).

Experimental

2-Hydroxyacetophenone (1.36 g, 0.01 mol) and methyl hydrazinecarboxylate (0.9 g, 0.01 mol) were dissolved in stirred methanol (15 ml) and left for 2 h at room temperature. The resulting solid was filtered off and recrystallized from ethanol to give the title compound in 85% yield. Single crystals suitable for X-ray analysis were obtained by slow evaporation of an ethanol solution at room temperature (m.p. 465–467 K).

Refinement

H atoms were positioned geometrically (O-H = 0.82 Å, N-H = 0.86 Å and C-H = 0.93 or 0.96 Å) and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$. A rotating group model was used for the methyl groups.

Figures




Fig. 1. Molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atomic numbering. Dashed lines indicate intramolecular hydrogen bonds.




Fig. 2. Crystal packing of the title compound. Dashed lines indicate intermolecular hydrogen bonds.

(E)-Methyl N'-[1-(2-hydroxyphenyl)ethylidene]hydrazinecarboxylate

Crystal data

$C_{10}H_{12}N_2O_3$	$F_{000} = 440$
$M_r = 208.22$	$D_x = 1.345 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 8.6432 (8) \text{ \AA}$	Cell parameters from 1809 reflections
$b = 12.6696 (11) \text{ \AA}$	$\theta = 2.5\text{--}25.0^\circ$
$c = 9.9810 (9) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$\beta = 109.837 (3)^\circ$	$T = 273 (2) \text{ K}$
$V = 1028.12 (16) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.28 \times 0.24 \times 0.23 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer	1809 independent reflections
Radiation source: fine-focus sealed tube	1587 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.019$
$T = 273(2) \text{ K}$	$\theta_{\text{max}} = 25.1^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2002)	$h = -10 \rightarrow 9$
$T_{\text{min}} = 0.973$, $T_{\text{max}} = 0.979$	$k = -14 \rightarrow 13$
10601 measured reflections	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.035$	$w = 1/[\sigma^2(F_o^2) + (0.0557P)^2 + 0.2233P]$
$wR(F^2) = 0.107$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.05$	$(\Delta/\sigma)_{\text{max}} = 0.001$
1809 reflections	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$
140 parameters	$\Delta\rho_{\text{min}} = -0.12 \text{ e \AA}^{-3}$
	Extinction correction: SHELXL97 (Sheldrick, 2008),
	$F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Primary atom site location: structure-invariant direct methods
 Extinction coefficient: 0.017 (3)
 Secondary atom site location: difference Fourier map

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.62350 (16)	-0.02927 (10)	0.76460 (14)	0.0409 (3)
C6	0.67306 (16)	-0.04195 (10)	0.91401 (14)	0.0403 (3)
C7	0.79166 (16)	0.03028 (10)	1.01395 (13)	0.0404 (3)
C9	1.01051 (17)	0.25416 (11)	0.97214 (14)	0.0430 (3)
C2	0.51172 (17)	-0.09885 (12)	0.67429 (16)	0.0497 (4)
H2	0.4799	-0.0896	0.5762	0.060*
C3	0.44767 (19)	-0.18132 (12)	0.72853 (18)	0.0577 (4)
H3	0.3740	-0.2279	0.6671	0.069*
C5	0.60373 (19)	-0.12639 (12)	0.96486 (17)	0.0538 (4)
H5	0.6335	-0.1365	1.0627	0.065*
C4	0.4927 (2)	-0.19499 (14)	0.87417 (19)	0.0623 (5)
H4	0.4484	-0.2502	0.9109	0.075*
C10	1.1944 (2)	0.39775 (14)	1.01329 (19)	0.0650 (5)
H10A	1.2220	0.3668	0.9365	0.098*
H10B	1.2933	0.4178	1.0883	0.098*
H10C	1.1272	0.4591	0.9794	0.098*
C8	0.8508 (2)	0.00907 (13)	1.17083 (15)	0.0595 (4)
H8A	0.9553	0.0427	1.2149	0.089*
H8B	0.8622	-0.0657	1.1871	0.089*
H8C	0.7727	0.0365	1.2110	0.089*
O1	0.68111 (13)	0.04936 (8)	0.70165 (10)	0.0535 (3)
H1	0.7464	0.0857	0.7631	0.080*
O2	0.98628 (13)	0.25492 (8)	0.84561 (10)	0.0539 (3)
O3	1.10557 (14)	0.32236 (9)	1.06669 (11)	0.0603 (3)
N1	0.83941 (13)	0.10946 (9)	0.95665 (11)	0.0416 (3)
N2	0.94739 (14)	0.18236 (10)	1.04034 (11)	0.0470 (3)
H2A	0.9736	0.1824	1.1315	0.056*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0454 (7)	0.0377 (7)	0.0412 (7)	0.0050 (5)	0.0167 (6)	0.0020 (5)
C6	0.0439 (7)	0.0363 (7)	0.0421 (7)	0.0051 (5)	0.0167 (6)	0.0023 (5)
C7	0.0460 (7)	0.0404 (7)	0.0361 (7)	0.0071 (6)	0.0156 (6)	0.0029 (5)
C9	0.0477 (8)	0.0444 (8)	0.0381 (7)	-0.0012 (6)	0.0163 (6)	-0.0035 (6)
C2	0.0521 (8)	0.0504 (9)	0.0447 (8)	0.0007 (6)	0.0138 (6)	-0.0056 (6)
C3	0.0532 (9)	0.0511 (9)	0.0655 (10)	-0.0078 (7)	0.0156 (8)	-0.0084 (7)
C5	0.0632 (9)	0.0506 (9)	0.0495 (8)	-0.0015 (7)	0.0216 (7)	0.0089 (7)
C4	0.0659 (10)	0.0522 (9)	0.0710 (11)	-0.0131 (7)	0.0260 (8)	0.0056 (8)
C10	0.0688 (10)	0.0671 (11)	0.0637 (10)	-0.0237 (9)	0.0284 (8)	-0.0088 (8)
C8	0.0833 (11)	0.0526 (9)	0.0384 (8)	-0.0026 (8)	0.0151 (7)	0.0045 (7)
O1	0.0721 (7)	0.0501 (6)	0.0362 (5)	-0.0107 (5)	0.0155 (5)	0.0007 (4)
O2	0.0721 (7)	0.0544 (7)	0.0368 (6)	-0.0094 (5)	0.0206 (5)	-0.0010 (4)
O3	0.0716 (7)	0.0673 (7)	0.0464 (6)	-0.0274 (6)	0.0258 (5)	-0.0137 (5)
N1	0.0460 (6)	0.0423 (6)	0.0358 (6)	-0.0029 (5)	0.0130 (5)	-0.0021 (5)
N2	0.0563 (7)	0.0524 (7)	0.0319 (6)	-0.0097 (5)	0.0145 (5)	-0.0034 (5)

Geometric parameters (\AA , $^\circ$)

C1—O1	1.3592 (16)	C5—C4	1.381 (2)
C1—C2	1.390 (2)	C5—H5	0.93
C1—C6	1.4141 (19)	C4—H4	0.93
C6—C5	1.4027 (19)	C10—O3	1.4364 (19)
C6—C7	1.4796 (19)	C10—H10A	0.96
C7—N1	1.2899 (17)	C10—H10B	0.96
C7—C8	1.4974 (18)	C10—H10C	0.96
C9—O2	1.2081 (16)	C8—H8A	0.96
C9—O3	1.3364 (17)	C8—H8B	0.96
C9—N2	1.3567 (17)	C8—H8C	0.96
C2—C3	1.377 (2)	O1—H1	0.82
C2—H2	0.93	N1—N2	1.3757 (16)
C3—C4	1.382 (2)	N2—H2A	0.86
C3—H3	0.93		
O1—C1—C2	116.62 (12)	C5—C4—C3	119.71 (15)
O1—C1—C6	122.91 (12)	C5—C4—H4	120.1
C2—C1—C6	120.46 (13)	C3—C4—H4	120.1
C5—C6—C1	117.03 (13)	O3—C10—H10A	109.5
C5—C6—C7	120.73 (12)	O3—C10—H10B	109.5
C1—C6—C7	122.23 (12)	H10A—C10—H10B	109.5
N1—C7—C6	115.77 (11)	O3—C10—H10C	109.5
N1—C7—C8	123.80 (13)	H10A—C10—H10C	109.5
C6—C7—C8	120.43 (12)	H10B—C10—H10C	109.5
O2—C9—O3	125.45 (13)	C7—C8—H8A	109.5
O2—C9—N2	124.98 (13)	C7—C8—H8B	109.5
O3—C9—N2	109.56 (11)	H8A—C8—H8B	109.5

C3—C2—C1	120.68 (14)	C7—C8—H8C	109.5
C3—C2—H2	119.7	H8A—C8—H8C	109.5
C1—C2—H2	119.7	H8B—C8—H8C	109.5
C2—C3—C4	120.09 (14)	C1—O1—H1	109.5
C2—C3—H3	120.0	C9—O3—C10	116.48 (11)
C4—C3—H3	120.0	C7—N1—N2	120.41 (11)
C4—C5—C6	122.01 (14)	C9—N2—N1	116.81 (11)
C4—C5—H5	119.0	C9—N2—H2A	121.6
C6—C5—H5	119.0	N1—N2—H2A	121.6
O1—C1—C6—C5	179.88 (12)	C1—C6—C5—C4	0.4 (2)
C2—C1—C6—C5	-0.42 (19)	C7—C6—C5—C4	179.96 (14)
O1—C1—C6—C7	0.3 (2)	C6—C5—C4—C3	0.2 (3)
C2—C1—C6—C7	-179.99 (12)	C2—C3—C4—C5	-0.8 (3)
C5—C6—C7—N1	-175.42 (12)	O2—C9—O3—C10	-4.6 (2)
C1—C6—C7—N1	4.12 (19)	N2—C9—O3—C10	174.30 (13)
C5—C6—C7—C8	5.3 (2)	C6—C7—N1—N2	178.87 (11)
C1—C6—C7—C8	-175.19 (13)	C8—C7—N1—N2	-1.8 (2)
O1—C1—C2—C3	179.57 (13)	O2—C9—N2—N1	-4.4 (2)
C6—C1—C2—C3	-0.2 (2)	O3—C9—N2—N1	176.72 (11)
C1—C2—C3—C4	0.8 (2)	C7—N1—N2—C9	170.54 (12)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N1	0.82	1.85	2.5625 (14)	145
N2—H2A \cdots O2 ⁱ	0.86	2.25	3.0550 (14)	156
C8—H8A \cdots N2	0.96	2.47	2.820 (2)	101
C8—H8C \cdots Cg1 ⁱⁱ	0.96	2.93	3.803 (2)	151

Symmetry codes: (i) $x, -y+1/2, z+1/2$; (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

