

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Ethyl 4-{2,6-dichloro-4-[3-(2,6-difluorobenzoyl)ureido]phenoxy}butanoate

Yin-Hong Liu, Fang-Shi Li,* Yi Li, Da-Sheng Yu and Chui Lu


 Department of Applied Chemistry, College of Science, Nanjing University of Technology, Xinmofan Road No. 5, Nanjing 210009, People's Republic of China
 Correspondence e-mail: fangshi.li@njut.edu.cn

Received 16 July 2008; accepted 3 August 2008

 Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.082; wR factor = 0.290; data-to-parameter ratio = 14.7.

The title compound, $\text{C}_{20}\text{H}_{18}\text{Cl}_2\text{F}_2\text{N}_2\text{O}_5$, is considered to belong to the fourth generation of insecticides with properties such as high selectivity, low acute toxicity for mammals and high biological activity. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond results in the formation of a six-membered ring. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{F}$ hydrogen bonds link the molecules.

Related literature

 For related literature, see: Wang *et al.* (1998, 1999). For bond-length data, see: Allen *et al.* (1987).


Experimental

Crystal data

 $\text{C}_{20}\text{H}_{18}\text{Cl}_2\text{F}_2\text{N}_2\text{O}_5$
 $M_r = 475.26$
 Monoclinic, $P2_1/n$
 $a = 11.262$ (2) Å
 $b = 10.463$ (2) Å
 $c = 18.613$ (4) Å

 $\beta = 98.78$ (3)°
 $V = 2167.5$ (8) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

 $\mu = 0.35$ mm⁻¹
 $T = 293$ (2) K
 $0.40 \times 0.30 \times 0.20$ mm

Data collection

 Enraf–Nonius CAD-4 diffractometer
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.872$, $T_{\max} = 0.933$
 4157 measured reflections

 3944 independent reflections
 2385 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 3 standard reflections every 200 reflections
 intensity decay: none

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.082$
 $wR(F^2) = 0.290$
 $S = 1.13$
 3944 reflections

 268 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.77$ e Å⁻³
 $\Delta\rho_{\min} = -0.99$ e Å⁻³

 Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2A}\cdots\text{O4}^i$	0.86	2.00	2.856 (6)	173
$\text{C5}-\text{H5A}\cdots\text{F2}^{ii}$	0.97	2.44	3.201 (8)	135
$\text{N1}-\text{H1A}\cdots\text{O5}$	0.86	1.97	2.675 (7)	138

 Symmetry codes: (i) $-x - 1, -y, -z$; (ii) $-x, -y, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXL97*; software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CS2088).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, S., Allan, R. D., Skerritt, J. H. & Kennedy, I. R. (1998). *J. Agric. Food Chem.* **46**, 3330–3338.
- Wang, S., Allan, R. D., Skerritt, J. H. & Kennedy, I. R. (1999). *J. Agric. Food Chem.* **47**, 3416–3424.

supplementary materials

Acta Cryst. (2008). E64, o1729 [doi:10.1107/S1600536808024987]

Ethyl 4-{2,6-dichloro-4-[3-(2,6-difluorobenzoyl)ureido]phenoxy}butanoate

Y.-H. Liu, F.-S. Li, Y. Li, D.-S. Yu and C. Lu

Comment

The title compound is generally recognized as an insect growth regulator that interferes with chitin synthesis in target pests, causing death or abortive development (Wang *et al.* 1998). Bonding dimensions conform to expected values (Allen *et al.*, 1987).

Experimental

The title compound was prepared according to a literature method (Wang *et al.* 1999). The crystals suitable for X-ray analysis were obtained by dissolving it (0.1 g) in acetonitrile (25 ml) and evaporating the solvent slowly at room temperature for about 6 d.

Refinement

H atoms were positioned geometrically, with C—H distances of 0.93 and of 0.97 Å for aromatic and methyl H atoms and with N—H = 0.86 Å for amido H atoms. All H atoms were constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{N})$, where $x = 1.2$ for all H atoms.

Figures




Fig. 1. The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. N—H···O intramolecular hydrogen bond is shown by dashed line.




Fig. 2. A packing diagram detail from the crystal structure. N—H...O and C—H...F intermolecular hydrogen bonds are shown by dashed lines.

Ethyl 4-{2,6-dichloro-4-[3-(2,6-difluorobenzoyl)ureido]phenoxy}butanoate

Crystal data

$C_{20}H_{18}Cl_2F_2N_2O_5$

$M_r = 475.26$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 11.262\ (2)\ \text{\AA}$

$b = 10.463\ (2)\ \text{\AA}$

$c = 18.613\ (4)\ \text{\AA}$

$\beta = 98.78\ (3)^\circ$

$V = 2167.5\ (8)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 976$

$D_x = 1.456\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$

$\mu = 0.35\ \text{mm}^{-1}$

$T = 293\ (2)\ \text{K}$

Block, colourless

$0.40 \times 0.30 \times 0.20\ \text{mm}$

Data collection

Enraf-Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293\ (2)\ \text{K}$

$\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.872$, $T_{\max} = 0.933$

4157 measured reflections

3944 independent reflections

2385 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.041$

$\theta_{\max} = 25.3^\circ$

$\theta_{\min} = 2.0^\circ$

$h = -13 \rightarrow 13$

$k = 0 \rightarrow 12$

$l = 0 \rightarrow 22$

3 standard reflections

every 200 reflections

intensity decay: none

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.082$	H-atom parameters constrained
$wR(F^2) = 0.290$	$w = 1/[\sigma^2(F_o^2) + (0.1372P)^2 + 4.2262P]$
$S = 1.13$	where $P = (F_o^2 + 2F_c^2)/3$
3944 reflections	$(\Delta/\sigma)_{\max} < 0.001$
268 parameters	$\Delta\rho_{\max} = 0.77 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.99 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Experimental. (North *et al.*, 1968)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.23061 (14)	0.09615 (16)	0.16895 (10)	0.0611 (5)
C12	0.02268 (15)	-0.30602 (15)	0.02019 (10)	0.0672 (6)
F1	-0.5092 (6)	0.3204 (6)	0.2109 (3)	0.127 (2)
F2	-0.5604 (4)	0.3418 (6)	-0.0403 (2)	0.0995 (17)
N1	-0.2177 (4)	0.0877 (5)	0.0826 (3)	0.0473 (12)
H1A	-0.2132	0.1606	0.1042	0.057*
O1	0.5929 (5)	0.0111 (5)	0.2331 (3)	0.0813 (16)
C1	0.7375 (8)	0.1501 (10)	0.2970 (5)	0.095
H1B	0.7622	0.1938	0.3421	0.142*
H1C	0.7173	0.2116	0.2588	0.142*
H1D	0.8019	0.0969	0.2861	0.142*
N2	-0.4207 (4)	0.1420 (5)	0.0532 (3)	0.0481 (12)
H2A	-0.4908	0.1202	0.0316	0.058*
O2	0.4801 (5)	-0.1199 (6)	0.2895 (3)	0.092
C2	0.6371 (8)	0.0743 (9)	0.3027 (4)	0.092 (3)
H2B	0.5738	0.1271	0.3170	0.110*
H2C	0.6583	0.0100	0.3400	0.110*
O3	0.2254 (3)	-0.1568 (4)	0.09703 (19)	0.0458 (10)
C3	0.5158 (6)	-0.0854 (7)	0.2341 (3)	0.0544 (16)
O4	-0.3529 (4)	-0.0477 (4)	0.0170 (2)	0.0578 (12)
C4	0.4795 (5)	-0.1407 (6)	0.1610 (4)	0.0545 (16)
H4A	0.5512	-0.1638	0.1411	0.065*
H4B	0.4376	-0.0757	0.1297	0.065*
O5	-0.3196 (4)	0.3054 (5)	0.1175 (3)	0.0775 (16)
C5	0.3991 (5)	-0.2578 (6)	0.1597 (4)	0.0513 (15)
H5A	0.4037	-0.3058	0.1156	0.062*

supplementary materials

H5B	0.4293	-0.3121	0.2006	0.062*
C6	0.2705 (5)	-0.2267 (6)	0.1628 (3)	0.0436 (13)
H6A	0.2246	-0.3046	0.1655	0.052*
H6B	0.2640	-0.1751	0.2053	0.052*
C7	0.1154 (5)	-0.1029 (6)	0.0964 (3)	0.0419 (13)
C8	0.1028 (5)	0.0189 (6)	0.1269 (3)	0.0431 (13)
C9	-0.0055 (5)	0.0791 (6)	0.1230 (3)	0.0455 (14)
H9A	-0.0108	0.1591	0.1440	0.055*
C10	-0.1091 (5)	0.0182 (5)	0.0867 (3)	0.0440 (14)
C11	-0.1009 (5)	-0.1023 (5)	0.0560 (3)	0.0453 (14)
H11A	-0.1688	-0.1437	0.0323	0.054*
C12	0.0113 (5)	-0.1586 (5)	0.0618 (3)	0.0464 (14)
C13	-0.3273 (5)	0.0529 (6)	0.0489 (3)	0.0428 (13)
C14	-0.4138 (6)	0.2591 (6)	0.0874 (3)	0.0509 (15)
C15	-0.5308 (5)	0.3250 (5)	0.0855 (3)	0.0431 (13)
C16	-0.5738 (7)	0.3548 (7)	0.1491 (4)	0.0622 (18)
C17	-0.6834 (8)	0.4151 (7)	0.1493 (5)	0.076 (2)
H17A	-0.7108	0.4325	0.1930	0.091*
C18	-0.7493 (7)	0.4481 (7)	0.0858 (5)	0.074 (2)
H18A	-0.8236	0.4871	0.0854	0.089*
C19	-0.7083 (6)	0.4251 (7)	0.0223 (4)	0.071 (2)
H19A	-0.7535	0.4498	-0.0215	0.085*
C20	-0.6010 (6)	0.3657 (7)	0.0230 (3)	0.0574 (17)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0473 (9)	0.0581 (10)	0.0746 (11)	-0.0088 (7)	-0.0015 (7)	-0.0205 (8)
Cl2	0.0649 (11)	0.0426 (9)	0.0892 (13)	0.0079 (8)	-0.0042 (9)	-0.0222 (8)
F1	0.181 (6)	0.146 (5)	0.057 (3)	0.081 (5)	0.027 (3)	0.020 (3)
F2	0.104 (3)	0.145 (5)	0.048 (2)	0.052 (3)	0.006 (2)	-0.015 (3)
N1	0.047 (3)	0.041 (3)	0.050 (3)	0.010 (2)	-0.007 (2)	-0.010 (2)
O1	0.092 (4)	0.072 (3)	0.068 (3)	-0.033 (3)	-0.024 (3)	0.011 (3)
C1	0.095	0.095	0.095	0.000	0.014	0.000
N2	0.048 (3)	0.038 (3)	0.056 (3)	0.009 (2)	0.001 (2)	-0.010 (2)
O2	0.092	0.092	0.092	0.000	0.014	0.000
C2	0.109 (7)	0.085 (6)	0.068 (5)	-0.039 (5)	-0.029 (5)	-0.001 (4)
O3	0.042 (2)	0.058 (3)	0.038 (2)	0.0023 (19)	0.0072 (16)	-0.0027 (18)
C3	0.055 (4)	0.060 (4)	0.047 (4)	-0.001 (3)	0.006 (3)	0.015 (3)
O4	0.048 (2)	0.041 (2)	0.077 (3)	0.0065 (19)	-0.015 (2)	-0.021 (2)
C4	0.045 (3)	0.056 (4)	0.063 (4)	-0.008 (3)	0.009 (3)	-0.001 (3)
O5	0.053 (3)	0.058 (3)	0.111 (4)	0.000 (2)	-0.020 (3)	-0.034 (3)
C5	0.038 (3)	0.052 (4)	0.063 (4)	0.002 (3)	0.008 (3)	-0.001 (3)
C6	0.043 (3)	0.042 (3)	0.045 (3)	0.003 (3)	0.004 (2)	0.001 (3)
C7	0.041 (3)	0.048 (3)	0.036 (3)	0.002 (3)	0.004 (2)	0.004 (3)
C8	0.046 (3)	0.043 (3)	0.040 (3)	-0.009 (3)	0.004 (2)	-0.003 (3)
C9	0.052 (3)	0.043 (3)	0.039 (3)	0.002 (3)	0.001 (3)	-0.008 (3)
C10	0.047 (3)	0.040 (3)	0.041 (3)	0.002 (3)	-0.004 (2)	-0.001 (3)

C11	0.045 (3)	0.038 (3)	0.051 (3)	0.002 (3)	-0.001 (3)	-0.006 (3)
C12	0.054 (3)	0.035 (3)	0.049 (3)	0.001 (3)	0.005 (3)	0.000 (3)
C13	0.040 (3)	0.043 (3)	0.043 (3)	0.011 (3)	-0.004 (2)	-0.003 (3)
C14	0.053 (4)	0.046 (4)	0.051 (4)	0.001 (3)	0.003 (3)	-0.002 (3)
C15	0.044 (3)	0.037 (3)	0.049 (3)	-0.005 (2)	0.008 (3)	-0.001 (3)
C16	0.091 (5)	0.051 (4)	0.049 (4)	0.010 (4)	0.024 (4)	0.011 (3)
C17	0.101 (6)	0.055 (4)	0.085 (6)	0.026 (4)	0.056 (5)	0.005 (4)
C18	0.060 (4)	0.050 (4)	0.114 (7)	0.015 (3)	0.020 (4)	-0.022 (4)
C19	0.062 (4)	0.073 (5)	0.074 (5)	0.016 (4)	0.001 (4)	-0.019 (4)
C20	0.059 (4)	0.061 (4)	0.050 (4)	0.007 (3)	-0.002 (3)	-0.018 (3)

Geometric parameters (Å, °)

C11—C8	1.730 (6)	C4—H4B	0.9700
C12—C12	1.739 (6)	O5—C14	1.222 (7)
F1—C16	1.314 (9)	C5—C6	1.494 (8)
F2—C20	1.350 (7)	C5—H5A	0.9700
N1—C13	1.347 (7)	C5—H5B	0.9700
N1—C10	1.414 (7)	C6—H6A	0.9700
N1—H1A	0.8600	C6—H6B	0.9700
O1—C3	1.333 (8)	C7—C12	1.379 (8)
O1—C2	1.472 (9)	C7—C8	1.410 (8)
C1—C2	1.399 (11)	C8—C9	1.365 (8)
C1—H1B	0.9600	C9—C10	1.408 (8)
C1—H1C	0.9600	C9—H9A	0.9300
C1—H1D	0.9600	C10—C11	1.394 (8)
N2—C14	1.378 (8)	C11—C12	1.383 (8)
N2—C13	1.417 (7)	C11—H11A	0.9300
N2—H2A	0.8600	C14—C15	1.482 (8)
O2—C3	1.217 (8)	C15—C20	1.372 (9)
C2—H2B	0.9700	C15—C16	1.381 (8)
C2—H2C	0.9700	C16—C17	1.386 (10)
O3—C7	1.360 (6)	C17—C18	1.342 (11)
O3—C6	1.449 (7)	C17—H17A	0.9300
C3—C4	1.477 (9)	C18—C19	1.354 (10)
O4—C13	1.223 (7)	C18—H18A	0.9300
C4—C5	1.522 (9)	C19—C20	1.358 (9)
C4—H4A	0.9700	C19—H19A	0.9300
C13—N1—C10	127.6 (5)	O3—C7—C8	121.3 (5)
C13—N1—H1A	116.2	C12—C7—C8	116.1 (5)
C10—N1—H1A	116.2	C9—C8—C7	122.6 (5)
C3—O1—C2	117.4 (6)	C9—C8—C11	118.9 (4)
C2—C1—H1B	109.5	C7—C8—C11	118.4 (4)
C2—C1—H1C	109.5	C8—C9—C10	119.1 (5)
H1B—C1—H1C	109.5	C8—C9—H9A	120.5
C2—C1—H1D	109.5	C10—C9—H9A	120.5
H1B—C1—H1D	109.5	C11—C10—C9	120.2 (5)
H1C—C1—H1D	109.5	C11—C10—N1	123.7 (5)
C14—N2—C13	128.4 (5)	C9—C10—N1	116.1 (5)

supplementary materials

C14—N2—H2A	115.8	C12—C11—C10	118.1 (5)
C13—N2—H2A	115.8	C12—C11—H11A	121.0
C1—C2—O1	110.9 (8)	C10—C11—H11A	121.0
C1—C2—H2B	109.5	C7—C12—C11	123.9 (5)
O1—C2—H2B	109.5	C7—C12—C12	117.9 (5)
C1—C2—H2C	109.5	C11—C12—C12	118.1 (5)
O1—C2—H2C	109.5	O4—C13—N1	126.2 (5)
H2B—C2—H2C	108.1	O4—C13—N2	118.2 (5)
C7—O3—C6	114.7 (4)	N1—C13—N2	115.6 (5)
O2—C3—O1	122.5 (7)	O5—C14—N2	123.3 (6)
O2—C3—C4	125.6 (6)	O5—C14—C15	122.2 (6)
O1—C3—C4	111.9 (5)	N2—C14—C15	114.5 (5)
C3—C4—C5	114.3 (6)	C20—C15—C16	115.1 (6)
C3—C4—H4A	108.7	C20—C15—C14	124.0 (5)
C5—C4—H4A	108.7	C16—C15—C14	120.8 (6)
C3—C4—H4B	108.7	F1—C16—C15	117.9 (6)
C5—C4—H4B	108.7	F1—C16—C17	119.7 (6)
H4A—C4—H4B	107.6	C15—C16—C17	122.3 (7)
C6—C5—C4	113.7 (5)	C18—C17—C16	119.2 (7)
C6—C5—H5A	108.8	C18—C17—H17A	120.4
C4—C5—H5A	108.8	C16—C17—H17A	120.4
C6—C5—H5B	108.8	C17—C18—C19	120.4 (7)
C4—C5—H5B	108.8	C17—C18—H18A	119.8
H5A—C5—H5B	107.7	C19—C18—H18A	119.8
O3—C6—C5	107.1 (5)	C18—C19—C20	119.7 (7)
O3—C6—H6A	110.3	C18—C19—H19A	120.1
C5—C6—H6A	110.3	C20—C19—H19A	120.1
O3—C6—H6B	110.3	F2—C20—C19	119.8 (6)
C5—C6—H6B	110.3	F2—C20—C15	117.1 (6)
H6A—C6—H6B	108.5	C19—C20—C15	123.2 (6)
O3—C7—C12	122.4 (5)		
C3—O1—C2—C1	165.2 (8)	C10—C11—C12—C7	0.4 (9)
C2—O1—C3—O2	1.5 (10)	C10—C11—C12—C12	-177.4 (4)
C2—O1—C3—C4	-179.5 (7)	C10—N1—C13—O4	2.0 (10)
O2—C3—C4—C5	-5.6 (10)	C10—N1—C13—N2	-179.6 (5)
O1—C3—C4—C5	175.5 (5)	C14—N2—C13—O4	178.4 (6)
C3—C4—C5—C6	79.5 (7)	C14—N2—C13—N1	-0.2 (9)
C7—O3—C6—C5	-169.7 (5)	C13—N2—C14—O5	3.4 (10)
C4—C5—C6—O3	64.3 (7)	C13—N2—C14—C15	-176.1 (5)
C6—O3—C7—C12	-99.1 (6)	O5—C14—C15—C20	116.7 (8)
C6—O3—C7—C8	85.8 (6)	N2—C14—C15—C20	-63.8 (8)
O3—C7—C8—C9	175.6 (5)	O5—C14—C15—C16	-60.2 (9)
C12—C7—C8—C9	0.3 (8)	N2—C14—C15—C16	119.3 (6)
O3—C7—C8—C11	-2.8 (7)	C20—C15—C16—F1	-178.7 (7)
C12—C7—C8—C11	-178.2 (4)	C14—C15—C16—F1	-1.5 (10)
C7—C8—C9—C10	-0.5 (9)	C20—C15—C16—C17	3.5 (10)
C11—C8—C9—C10	177.9 (4)	C14—C15—C16—C17	-179.3 (6)
C8—C9—C10—C11	0.7 (9)	F1—C16—C17—C18	-179.1 (8)
C8—C9—C10—N1	-178.0 (5)	C15—C16—C17—C18	-1.3 (12)

C13—N1—C10—C11	-1.2 (9)	C16—C17—C18—C19	-1.3 (12)
C13—N1—C10—C9	177.4 (6)	C17—C18—C19—C20	1.4 (12)
C9—C10—C11—C12	-0.6 (9)	C18—C19—C20—F2	179.5 (7)
N1—C10—C11—C12	178.0 (5)	C18—C19—C20—C15	1.1 (11)
O3—C7—C12—C11	-175.5 (5)	C16—C15—C20—F2	178.1 (6)
C8—C7—C12—C11	-0.2 (9)	C14—C15—C20—F2	1.0 (10)
O3—C7—C12—C12	2.2 (8)	C16—C15—C20—C19	-3.4 (10)
C8—C7—C12—C12	177.5 (4)	C14—C15—C20—C19	179.5 (6)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2A...O4 ⁱ	0.86	2.00	2.856 (6)	173
C5—H5A...F2 ⁱⁱ	0.97	2.44	3.201 (8)	135
N1—H1A...O5	0.86	1.97	2.675 (7)	138

Symmetry codes: (i) $-x-1, -y, -z$; (ii) $-x, -y, -z$.

Fig. 1




Fig. 2

