

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2'-Dimethyl-7,7'-(methylenediimino)di-1,8-naphthyridin-1-ium bis(perchlorate)

Juan Mo,* Jian-Hua Liu, Yu-Shan Pan, Su-Mei Zhang and Xiang-Dang Du

College of Animal Husbandry and Veterinary Studies, Henan Agricultural University, Zhengzhou, Henan Province 450002, People's Republic of China

Correspondence e-mail: mojuan52@126.com


Received 14 June 2008; accepted 31 July 2008

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in solvent or counterion; R factor = 0.060; wR factor = 0.134; data-to-parameter ratio = 12.6.

In the title salt, $\text{C}_{19}\text{H}_{20}\text{N}_6^{2+} \cdot 2\text{ClO}_4^-$, the two planar 1,8-naphthyridine systems are linked by a methylenediamine group with a dihedral angle of $60.6(1)^\circ$ between the two systems. The crystal structure involves extensive $\text{N}-\text{H} \cdots \text{O}$ and $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonding.

Related literature

For related literature, see: Baker & Norman (2004); Gavrilova & Bosnich (2004); Nakatani *et al.* (2000, 2001); Stadie *et al.* (2007); Ferrarini *et al.* (1997).



Experimental

Crystal data

$\text{C}_{19}\text{H}_{20}\text{N}_6^{2+} \cdot 2\text{ClO}_4^-$

$M_r = 531.31$

Orthorhombic, $Pbca$

$a = 8.191(1)$ Å

$b = 19.325(2)$ Å

$c = 27.885(2)$ Å

$V = 4413.9(5)$ Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 0.36$ mm⁻¹

$T = 113(2)$ K

$0.34 \times 0.16 \times 0.14$ mm

Data collection

Bruker SMART CCD area-detector diffractometer

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.908$, $T_{\max} = 0.952$

31220 measured reflections
3882 independent reflections

3598 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.061$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$

$wR(F^2) = 0.134$

$S = 1.16$

3882 reflections

308 parameters

16 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.35$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.41$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1} \cdots \text{O6}^i$	0.88 (4)	1.92	2.794 (4)	168
$\text{N3}-\text{H3A} \cdots \text{O3}$	0.80 (3)	2.23	2.990 (6)	159
$\text{N4}-\text{H4A} \cdots \text{O5}^i$	0.71 (4)	2.56	3.233 (5)	160
$\text{N4}-\text{H4A} \cdots \text{O7}^i$	0.71 (4)	2.52	3.133 (5)	145
$\text{N6}-\text{H6} \cdots \text{O1}$	0.84 (3)	2.00	2.838 (8)	178
$\text{C1}-\text{H1A} \cdots \text{O7}^{ii}$	0.98	2.54	3.501 (3)	167
$\text{C1}-\text{H1B} \cdots \text{O4}^{iii}$	0.98	2.33	3.078 (4)	132
$\text{C4}-\text{H4} \cdots \text{O8}^{iv}$	0.95	2.52	3.392 (7)	152
$\text{C10}-\text{H10B} \cdots \text{O4}^{iv}$	0.99	2.35	3.082 (5)	130
$\text{C13}-\text{H13} \cdots \text{O2}^i$	0.95	2.41	3.259 (3)	149
$\text{C19}-\text{H19B} \cdots \text{O5}$	0.98	2.57	3.351 (6)	136
$\text{C19}-\text{H19C} \cdots \text{O7}^v$	0.98	2.58	3.207 (7)	122
$\text{C19}-\text{H19C} \cdots \text{O8}^v$	0.98	2.57	3.548 (4)	172

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x + \frac{3}{2}, -y, z - \frac{1}{2}$; (iii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (v) $x - 1, y, z$.

Data collection: *SMART* (Bruker, 1997); cell refinement: *SAINT* (Bruker, 1997); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *XP* in *SHELXTL*.

We thank Henan Agricultural University for the generous support of this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2415).

References

- Baker, R. S. & Norman, R. E. (2004). *Acta Cryst.* **E60**, m1761–m1763.
 Bruker (1997). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Ferrarini, P. L., Mori, C., Badawneh, M., Manera, C., Martinelli, A., Miceli, M., Ramagnoli, F. & Saccomanni, G. (1997). *J. Heterocycl. Chem.* **34**, 1501–1504.
 Gavrilova, E. L. & Bosnich, B. (2004). *Chem. Rev.* **104**, 349–383.
 Nakatani, K., Sando, S. & Saito, I. (2000). *J. Am. Chem. Soc.* **122**, 2172–2178.
 Nakatani, K., Sando, S. & Saito, I. (2001). *Nat. Biotechnol.* **19**, 51–55.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stadie, N. P., Sanchez-Smith, R. & Groy, T. L. (2007). *Acta Cryst.* **E63**, m2153–m2154.

supplementary materials

Acta Cryst. (2008). E64, o1702 [doi:10.1107/S1600536808024616]

2,2'-Dimethyl-7,7'-(methylenediimino)di-1,8-naphthyridin-1-ium bis(perchlorate)

J. Mo, J.-H. Liu, Y.-S. Pan, S.-M. Zhang and X.-D. Du

Comment

1,8-Naphthyridine and its derivatives are used for binding of mismatched guanine or used as versatile ligands which are able to form metal aggregates with monodentate fashion or chelating bidentate fashion (Nakatani *et al.*, 2000; Nakatani *et al.*, 2001; Ferrarini *et al.*, 1997; Gavrilova & Bosnich, 2004; Baker & Norman, 2004; Stadie *et al.*, 2007). We report here a new 1,8-Naphthyridine compound (Fig. 1).

The title compound reveals 1,8-naphthyridine rings are linked by methenediamine with a dihedral angle between two 1,8-naphthyridine rings of $60.6 (1)^\circ$. Each 1,8-naphthyridine ring is almost planar in which the ten atoms forming the 1,8-naphthyridine ring have mean deviation of 0.03 \AA from the least-squares plane calculated using the ten atoms. To balance hydrogen ion charge of two 1,8-naphthyridine rings, there are two perchlorate groups in crystal cell. From the packing diagram (Fig. 2), it seems that the intramolecular $\text{N-H}\cdots\text{O}$ and $\text{C-H}\cdots\text{O}$ and hydrogen bonds are effective in the stabilization of the crystal structure.

Experimental

To the solution of 2-amino-7-methyl-1,8-naphthyridine (3.18 g, 0.02 mol) in mixed solvent of water (28 mL) and ethanol (2 mL), 37% formaldehyde solution (0.86 mL, 0.01 mol) was added dropwise at 0°C and the reaction mixture was stirred at room temperature for 24h. The white precipitate formed was filtered, washed several times with water and then with diethyl ether and dried. Yield: 55% (1.81 g). FTIR (KBr) cm^{-1} : ν_{NH} 3389, 3266; ν_{CH} 3026. Anal. Calc. For $\text{C}_{19}\text{H}_{18}\text{N}_6$: C, 69.07; H, 5.49; N, 25.44. Found: C, 68.86; H, 5.56; N, 25.37. Single crystals of (I) suitable for an X-ray study were obtained by slow evaporation of an aqueous ethanol solution (30% v/v) under the conditions in the presence of perchloric acid at 293 K over a period of one month.

Refinement

Hydrogen atoms of NH (naphthyridine and amine) were located in a Fourier map and refined freely. All the other hydrogen atoms were generated geometrically (C—H bond lengths of methyl group fixed at 0.98 \AA , C—H bond lengths of naphthyridine fixed at 0.95 \AA) assigned appropriate isotropic thermal parameters, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. Each perchlorate anion is disordered over two different orientations. The Cl—O distances were restrained to $1.43 (4) \text{ \AA}$.

Figures




Fig. 1. Molecular structure of the cation of the title compound showing the atom-numbering scheme and displacement ellipsoids drawn at the 40% probability level.




Fig. 2. Unit-cell packing diagram as viewed down the *c*-direction. Hydrogen bonds are shown as dashed lines.

2,2'-Dimethyl-7,7'-(methylenediimino)di-1,8-naphthyridin-1-ium bis(perchlorate)

Crystal data

$C_{19}H_{20}N_6^{2+} \cdot 2ClO_4^-$

$M_r = 531.31$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 8.1910 (5) \text{ \AA}$

$b = 19.3250 (12) \text{ \AA}$

$c = 27.8850 (19) \text{ \AA}$

$V = 4413.9 (5) \text{ \AA}^3$

$Z = 8$

$F_{000} = 2192$

$D_x = 1.599 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71070 \text{ \AA}$

Cell parameters from 7263 reflections

$\theta = 2.1\text{--}28.0^\circ$

$\mu = 0.36 \text{ mm}^{-1}$

$T = 113 (2) \text{ K}$

Prism, colorless

$0.34 \times 0.16 \times 0.14 \text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: $7.31 \text{ pixels mm}^{-1}$

$T = 113(2) \text{ K}$

φ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.909$, $T_{\max} = 0.952$

31220 measured reflections

3882 independent reflections

3598 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.061$

$\theta_{\max} = 25.0^\circ$

$\theta_{\min} = 2.1^\circ$

$h = -9 \rightarrow 9$

$k = -22 \rightarrow 22$

$l = -33 \rightarrow 32$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.060$

$wR(F^2) = 0.134$

$S = 1.16$

3882 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0498P)^2 + 5.3196P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.003$

$\Delta\rho_{\max} = 0.35 \text{ e \AA}^{-3}$

308 parameters

$$\Delta\rho_{\min} = -0.41 \text{ e } \text{\AA}^{-3}$$

16 restraints

Extinction correction: none

Primary atom site location: structure-invariant direct methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.24246 (10)	0.22574 (4)	0.56855 (3)	0.0360 (2)	
C12	1.00163 (9)	0.01682 (3)	0.69459 (3)	0.0323 (2)	
O1	0.1693 (6)	0.1730 (2)	0.59780 (16)	0.0534 (14)	0.747 (5)
O2	0.3894 (5)	0.2014 (2)	0.5483 (2)	0.0758 (17)	0.747 (5)
O3	0.1302 (6)	0.2438 (2)	0.53084 (12)	0.0738 (15)	0.747 (5)
O4	0.2651 (6)	0.28664 (19)	0.59698 (15)	0.0700 (14)	0.747 (5)
O5	0.8509 (4)	0.02894 (17)	0.67091 (14)	0.0664 (12)	0.870 (6)
O6	0.9850 (5)	-0.03196 (14)	0.73264 (10)	0.0657 (13)	0.870 (6)
O7	1.1129 (3)	-0.01123 (15)	0.65972 (10)	0.0490 (10)	0.870 (6)
O8	1.0661 (6)	0.0802 (2)	0.7134 (2)	0.0444 (13)	0.870 (6)
O1'	0.1358 (14)	0.1676 (5)	0.5766 (4)	0.039 (3)	0.253 (5)
O2'	0.3710 (11)	0.2233 (6)	0.6037 (3)	0.062 (4)	0.253 (5)
O3'	0.3135 (14)	0.2178 (6)	0.5220 (3)	0.062 (4)	0.253 (5)
O4'	0.1558 (13)	0.2875 (4)	0.5708 (4)	0.052 (3)	0.253 (5)
O5'	0.8301 (12)	0.0031 (10)	0.7058 (7)	0.053 (6)	0.130 (6)
O6'	1.0870 (19)	-0.0453 (6)	0.7069 (7)	0.048 (6)	0.130 (6)
O7'	1.012 (3)	0.0307 (10)	0.6446 (4)	0.062 (7)	0.130 (6)
O8'	1.056 (3)	0.0738 (10)	0.7228 (8)	0.039 (10)	0.130 (6)
N1	0.1466 (3)	0.16498 (13)	0.27420 (9)	0.0328 (6)	
N2	0.1460 (3)	0.16453 (12)	0.35633 (8)	0.0293 (6)	
N3	0.1396 (3)	0.16601 (15)	0.43819 (10)	0.0353 (6)	
N6	0.3295 (3)	0.04505 (14)	0.58097 (9)	0.0336 (6)	
N5	0.2041 (3)	0.04253 (13)	0.50731 (9)	0.0327 (6)	
N4	0.0922 (4)	0.04231 (16)	0.43264 (11)	0.0417 (7)	
C2	0.1739 (4)	0.19127 (16)	0.23034 (10)	0.0377 (8)	
C3	0.2463 (5)	0.25570 (18)	0.22665 (11)	0.0435 (9)	
H3	0.2634	0.2759	0.1960	0.052*	
C4	0.2935 (4)	0.29061 (16)	0.26746 (11)	0.0390 (8)	
H4	0.3453	0.3344	0.2647	0.047*	

supplementary materials

C5	0.2662 (4)	0.26239 (14)	0.31277 (10)	0.0285 (7)
C6	0.1858 (4)	0.19774 (14)	0.31595 (10)	0.0268 (6)
C7	0.1866 (4)	0.19636 (15)	0.39716 (10)	0.0288 (7)
C8	0.2786 (4)	0.25928 (15)	0.39855 (10)	0.0314 (7)
H8	0.3132	0.2779	0.4284	0.038*
C9	0.3159 (4)	0.29193 (15)	0.35722 (10)	0.0324 (7)
H9	0.3748	0.3343	0.3578	0.039*
C1	0.1243 (6)	0.14783 (18)	0.18868 (12)	0.0518 (10)
H1A	0.2123	0.1154	0.1808	0.078*
H1B	0.1026	0.1776	0.1610	0.078*
H1C	0.0253	0.1219	0.1968	0.078*
C16	0.4320 (4)	0.02351 (16)	0.61578 (11)	0.0360 (8)
C17	0.5112 (4)	-0.03943 (17)	0.60916 (12)	0.0402 (8)
H17	0.5862	-0.0556	0.6326	0.048*
C18	0.4812 (4)	-0.07835 (17)	0.56878 (12)	0.0398 (8)
H18	0.5322	-0.1222	0.5653	0.048*
C14	0.3773 (4)	-0.05443 (15)	0.53293 (12)	0.0357 (7)
C15	0.3015 (4)	0.01043 (16)	0.53928 (11)	0.0316 (7)
C11	0.1802 (4)	0.00934 (17)	0.46592 (11)	0.0368 (8)
C12	0.2463 (4)	-0.05808 (16)	0.45608 (12)	0.0424 (8)
H12	0.2221	-0.0808	0.4267	0.051*
C13	0.3428 (4)	-0.08878 (16)	0.48880 (12)	0.0411 (8)
H13	0.3878	-0.1332	0.4825	0.049*
C19	0.4554 (5)	0.06921 (19)	0.65809 (11)	0.0446 (8)
H19A	0.4863	0.1156	0.6473	0.067*
H19B	0.5418	0.0503	0.6785	0.067*
H19C	0.3533	0.0718	0.6763	0.067*
C10	0.0224 (4)	0.10976 (18)	0.43970 (12)	0.0431 (8)
H10A	-0.0336	0.1106	0.4712	0.052*
H10B	-0.0612	0.1178	0.4147	0.052*
H3A	0.163 (4)	0.1865 (17)	0.4619 (12)	0.035 (10)*
H1	0.101 (5)	0.124 (2)	0.2764 (13)	0.055 (11)*
H4A	0.081 (5)	0.0269 (19)	0.4097 (13)	0.040 (12)*
H6	0.281 (4)	0.0828 (18)	0.5853 (12)	0.035 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0433 (5)	0.0317 (4)	0.0331 (4)	-0.0003 (3)	0.0046 (4)	-0.0008 (3)
C12	0.0360 (5)	0.0256 (4)	0.0353 (4)	-0.0007 (3)	0.0018 (3)	-0.0012 (3)
O1	0.064 (3)	0.046 (2)	0.050 (3)	0.016 (2)	0.026 (2)	0.022 (2)
O2	0.055 (3)	0.049 (2)	0.124 (4)	-0.002 (2)	0.053 (3)	-0.011 (3)
O3	0.118 (4)	0.059 (2)	0.045 (2)	0.031 (2)	-0.025 (2)	0.0033 (18)
O4	0.089 (3)	0.052 (2)	0.070 (3)	-0.001 (2)	0.002 (3)	-0.027 (2)
O5	0.0421 (19)	0.063 (2)	0.095 (3)	0.0139 (15)	-0.0291 (19)	-0.019 (2)
O6	0.126 (4)	0.0396 (16)	0.0313 (15)	-0.0408 (19)	-0.0042 (19)	0.0050 (12)
O7	0.0410 (17)	0.0542 (18)	0.0519 (18)	-0.0075 (14)	0.0127 (14)	-0.0213 (14)
O8	0.055 (3)	0.0232 (18)	0.055 (2)	-0.0038 (16)	-0.002 (2)	-0.0076 (17)

O1'	0.036 (6)	0.030 (5)	0.051 (7)	-0.003 (4)	0.014 (5)	0.002 (5)
O2'	0.049 (6)	0.086 (7)	0.052 (6)	-0.022 (5)	-0.013 (5)	0.031 (5)
O3'	0.056 (7)	0.087 (7)	0.041 (6)	-0.016 (6)	0.023 (5)	-0.020 (5)
O4'	0.063 (6)	0.027 (4)	0.065 (6)	0.017 (4)	0.012 (5)	0.007 (4)
O5'	0.030 (8)	0.067 (9)	0.061 (10)	0.008 (7)	0.004 (7)	0.003 (8)
O6'	0.046 (9)	0.033 (8)	0.064 (10)	0.009 (7)	0.000 (8)	0.003 (7)
O7'	0.085 (11)	0.060 (10)	0.041 (9)	-0.012 (8)	0.014 (8)	-0.003 (7)
O8'	0.053 (14)	0.032 (13)	0.032 (12)	0.002 (8)	-0.006 (8)	-0.010 (8)
N1	0.0448 (17)	0.0233 (13)	0.0302 (14)	-0.0046 (12)	-0.0070 (12)	0.0000 (10)
N2	0.0321 (14)	0.0275 (12)	0.0283 (13)	-0.0027 (11)	-0.0009 (11)	0.0035 (10)
N3	0.0381 (16)	0.0419 (15)	0.0260 (14)	0.0023 (13)	-0.0004 (12)	0.0032 (12)
N6	0.0353 (15)	0.0344 (15)	0.0310 (14)	0.0059 (13)	0.0054 (12)	0.0103 (12)
N5	0.0314 (14)	0.0360 (14)	0.0306 (13)	-0.0023 (11)	0.0053 (11)	0.0101 (11)
N4	0.0479 (19)	0.0467 (18)	0.0306 (16)	-0.0128 (14)	-0.0024 (15)	0.0084 (14)
C2	0.053 (2)	0.0331 (16)	0.0266 (16)	0.0011 (15)	-0.0066 (15)	-0.0008 (13)
C3	0.064 (2)	0.0381 (18)	0.0287 (16)	-0.0036 (17)	-0.0008 (16)	0.0063 (14)
C4	0.050 (2)	0.0317 (16)	0.0349 (17)	-0.0098 (15)	0.0017 (15)	0.0043 (13)
C5	0.0341 (17)	0.0212 (14)	0.0302 (15)	-0.0031 (12)	-0.0020 (13)	-0.0004 (12)
C6	0.0325 (16)	0.0223 (13)	0.0257 (14)	-0.0012 (12)	-0.0021 (13)	-0.0003 (11)
C7	0.0299 (16)	0.0301 (15)	0.0264 (15)	0.0067 (13)	0.0012 (13)	0.0022 (12)
C8	0.0359 (18)	0.0293 (15)	0.0290 (15)	0.0050 (13)	-0.0035 (13)	-0.0066 (12)
C9	0.0374 (18)	0.0262 (15)	0.0335 (16)	-0.0018 (13)	-0.0029 (14)	-0.0030 (12)
C1	0.083 (3)	0.0408 (19)	0.0316 (18)	-0.0031 (19)	-0.0128 (19)	-0.0052 (15)
C16	0.0290 (17)	0.0429 (18)	0.0362 (17)	-0.0019 (14)	0.0067 (14)	0.0172 (14)
C17	0.0313 (18)	0.0449 (19)	0.0444 (19)	0.0015 (15)	0.0056 (15)	0.0222 (16)
C18	0.0337 (18)	0.0324 (17)	0.053 (2)	0.0026 (14)	0.0174 (16)	0.0202 (15)
C14	0.0355 (18)	0.0285 (15)	0.0430 (18)	-0.0041 (14)	0.0150 (15)	0.0115 (14)
C15	0.0287 (17)	0.0350 (16)	0.0309 (16)	-0.0028 (13)	0.0097 (13)	0.0104 (13)
C11	0.0381 (19)	0.0409 (18)	0.0314 (16)	-0.0138 (15)	0.0053 (15)	0.0099 (14)
C12	0.052 (2)	0.0331 (17)	0.0417 (19)	-0.0172 (16)	0.0116 (17)	0.0006 (15)
C13	0.048 (2)	0.0273 (16)	0.048 (2)	-0.0080 (15)	0.0165 (17)	0.0061 (14)
C19	0.042 (2)	0.058 (2)	0.0341 (17)	-0.0008 (17)	0.0011 (16)	0.0121 (16)
C10	0.0343 (19)	0.061 (2)	0.0343 (18)	-0.0038 (17)	0.0030 (15)	0.0176 (16)

Geometric parameters (Å, °)

C11—O4'	1.390 (7)	C2—C3	1.383 (5)
C11—O2	1.410 (4)	C2—C1	1.490 (4)
C11—O3'	1.430 (7)	C3—C4	1.378 (4)
C11—O4	1.431 (3)	C3—H3	0.9500
C11—O1	1.436 (4)	C4—C5	1.394 (4)
C11—O2'	1.439 (7)	C4—H4	0.9500
C11—O3	1.440 (3)	C5—C6	1.415 (4)
C11—O1'	1.441 (8)	C5—C9	1.424 (4)
C12—O5	1.420 (3)	C7—C8	1.431 (4)
C12—O7'	1.423 (9)	C8—C9	1.349 (4)
C12—O8'	1.424 (10)	C8—H8	0.9500
C12—O6	1.426 (3)	C9—H9	0.9500
C12—O6'	1.432 (9)	C1—H1A	0.9800

supplementary materials

C12—O8	1.433 (3)	C1—H1B	0.9800
C12—O7	1.439 (3)	C1—H1C	0.9800
C12—O5'	1.464 (9)	C16—C17	1.390 (5)
N1—C2	1.343 (4)	C16—C19	1.486 (5)
N1—C6	1.364 (4)	C17—C18	1.376 (5)
N1—H1	0.88 (4)	C17—H17	0.9500
N2—C7	1.336 (4)	C18—C14	1.392 (5)
N2—C6	1.336 (4)	C18—H18	0.9500
N3—C7	1.342 (4)	C14—C15	1.410 (4)
N3—C10	1.451 (4)	C14—C13	1.427 (5)
N3—H3A	0.80 (3)	C11—C12	1.437 (5)
N6—C16	1.350 (4)	C12—C13	1.345 (5)
N6—C15	1.360 (4)	C12—H12	0.9500
N6—H6	0.84 (3)	C13—H13	0.9500
N5—C11	1.335 (4)	C19—H19A	0.9800
N5—C15	1.348 (4)	C19—H19B	0.9800
N4—C11	1.337 (4)	C19—H19C	0.9800
N4—C10	1.437 (5)	C10—H10A	0.9900
N4—H4A	0.71 (4)	C10—H10B	0.9900
O4'—C11—O2	137.8 (4)	N1—C2—C3	118.7 (3)
O4'—C11—O3'	109.9 (6)	N1—C2—C1	116.8 (3)
O4'—C11—O4	48.3 (4)	C3—C2—C1	124.5 (3)
O2—C11—O4	112.7 (3)	C4—C3—C2	120.0 (3)
O3'—C11—O4	122.6 (5)	C4—C3—H3	120.0
O4'—C11—O1	111.8 (5)	C2—C3—H3	120.0
O2—C11—O1	110.3 (3)	C3—C4—C5	120.8 (3)
O3'—C11—O1	127.5 (5)	C3—C4—H4	119.6
O4—C11—O1	108.9 (3)	C5—C4—H4	119.6
O4'—C11—O2'	111.7 (6)	C4—C5—C6	118.5 (3)
O2—C11—O2'	68.8 (5)	C4—C5—C9	125.9 (3)
O3'—C11—O2'	108.5 (6)	C6—C5—C9	115.6 (3)
O4—C11—O2'	63.6 (5)	N2—C6—N1	116.0 (2)
O1—C11—O2'	84.0 (4)	N2—C6—C5	126.2 (3)
O4'—C11—O3	59.9 (5)	N1—C6—C5	117.8 (3)
O2—C11—O3	109.5 (3)	N2—C7—N3	117.0 (3)
O3'—C11—O3	67.9 (5)	N2—C7—C8	123.0 (3)
O4—C11—O3	106.7 (2)	N3—C7—C8	120.0 (3)
O1—C11—O3	108.7 (3)	C9—C8—C7	119.6 (3)
O2'—C11—O3	166.5 (4)	C9—C8—H8	120.2
O4'—C11—O1'	110.7 (6)	C7—C8—H8	120.2
O2—C11—O1'	108.6 (6)	C8—C9—C5	119.4 (3)
O3'—C11—O1'	107.7 (6)	C8—C9—H9	120.3
O4—C11—O1'	129.3 (5)	C5—C9—H9	120.3
O2'—C11—O1'	108.2 (6)	C2—C1—H1A	109.5
O3—C11—O1'	85.2 (5)	C2—C1—H1B	109.5
O5—C12—O7'	64.3 (8)	H1A—C1—H1B	109.5
O5—C12—O8'	113.6 (11)	C2—C1—H1C	109.5
O7'—C12—O8'	112.1 (10)	H1A—C1—H1C	109.5
O5—C12—O6	111.8 (2)	H1B—C1—H1C	109.5

O7'—C12—O6	149.3 (8)	N6—C16—C17	117.7 (3)
O8'—C12—O6	97.5 (11)	N6—C16—C19	117.9 (3)
O5—C12—O6'	132.4 (7)	C17—C16—C19	124.4 (3)
O7'—C12—O6'	111.3 (8)	C18—C17—C16	120.2 (3)
O8'—C12—O6'	111.4 (10)	C18—C17—H17	119.9
O6—C12—O6'	46.6 (7)	C16—C17—H17	119.9
O5—C12—O8	110.5 (2)	C17—C18—C14	121.0 (3)
O7'—C12—O8	100.1 (8)	C17—C18—H18	119.5
O6—C12—O8	109.1 (3)	C14—C18—H18	119.5
O6'—C12—O8	116.7 (8)	C18—C14—C15	118.3 (3)
O5—C12—O7	107.38 (19)	C18—C14—C13	125.9 (3)
O7'—C12—O7	50.9 (8)	C15—C14—C13	115.8 (3)
O8'—C12—O7	117.9 (11)	N5—C15—N6	116.0 (3)
O6—C12—O7	108.3 (2)	N5—C15—C14	125.9 (3)
O6'—C12—O7	62.4 (8)	N6—C15—C14	118.1 (3)
O8—C12—O7	109.6 (3)	N5—C11—N4	116.8 (3)
O5—C12—O5'	45.2 (7)	N5—C11—C12	123.1 (3)
O7'—C12—O5'	107.7 (8)	N4—C11—C12	120.2 (3)
O8'—C12—O5'	108.7 (10)	C13—C12—C11	119.5 (3)
O6—C12—O5'	68.2 (8)	C13—C12—H12	120.3
O6'—C12—O5'	105.3 (8)	C11—C12—H12	120.3
O8—C12—O5'	115.5 (8)	C12—C13—C14	119.8 (3)
O7—C12—O5'	133.2 (8)	C12—C13—H13	120.1
C2—N1—C6	124.2 (3)	C14—C13—H13	120.1
C2—N1—H1	118 (2)	C16—C19—H19A	109.5
C6—N1—H1	117 (2)	C16—C19—H19B	109.5
C7—N2—C6	115.9 (2)	H19A—C19—H19B	109.5
C7—N3—C10	122.8 (3)	C16—C19—H19C	109.5
C7—N3—H3A	115 (2)	H19A—C19—H19C	109.5
C10—N3—H3A	121 (2)	H19B—C19—H19C	109.5
C16—N6—C15	124.6 (3)	N4—C10—N3	114.4 (3)
C16—N6—H6	117 (2)	N4—C10—H10A	108.7
C15—N6—H6	118 (2)	N3—C10—H10A	108.7
C11—N5—C15	116.0 (3)	N4—C10—H10B	108.7
C11—N4—C10	123.5 (3)	N3—C10—H10B	108.7
C11—N4—H4A	120 (3)	H10A—C10—H10B	107.6
C10—N4—H4A	117 (3)		
C6—N1—C2—C3	-0.1 (5)	C15—N6—C16—C19	176.9 (3)
C6—N1—C2—C1	179.5 (3)	N6—C16—C17—C18	-1.5 (4)
N1—C2—C3—C4	2.1 (5)	C19—C16—C17—C18	179.8 (3)
C1—C2—C3—C4	-177.4 (4)	C16—C17—C18—C14	2.8 (5)
C2—C3—C4—C5	-1.4 (6)	C17—C18—C14—C15	-0.8 (4)
C3—C4—C5—C6	-1.3 (5)	C17—C18—C14—C13	177.1 (3)
C3—C4—C5—C9	177.0 (3)	C11—N5—C15—N6	178.3 (3)
C7—N2—C6—N1	179.2 (3)	C11—N5—C15—C14	-0.7 (4)
C7—N2—C6—C5	0.2 (4)	C16—N6—C15—N5	-175.3 (3)
C2—N1—C6—N2	178.4 (3)	C16—N6—C15—C14	3.8 (4)
C2—N1—C6—C5	-2.5 (5)	C18—C14—C15—N5	176.7 (3)
C4—C5—C6—N2	-177.9 (3)	C13—C14—C15—N5	-1.5 (4)

supplementary materials

C9—C5—C6—N2	3.7 (5)	C18—C14—C15—N6	-2.3 (4)
C4—C5—C6—N1	3.1 (4)	C13—C14—C15—N6	179.5 (3)
C9—C5—C6—N1	-175.3 (3)	C15—N5—C11—N4	-176.0 (3)
C6—N2—C7—N3	176.2 (3)	C15—N5—C11—C12	3.0 (4)
C6—N2—C7—C8	-4.9 (4)	C10—N4—C11—N5	-0.9 (5)
C10—N3—C7—N2	-12.6 (4)	C10—N4—C11—C12	-179.9 (3)
C10—N3—C7—C8	168.5 (3)	N5—C11—C12—C13	-2.9 (5)
N2—C7—C8—C9	5.7 (5)	N4—C11—C12—C13	176.0 (3)
N3—C7—C8—C9	-175.5 (3)	C11—C12—C13—C14	0.5 (5)
C7—C8—C9—C5	-1.4 (5)	C18—C14—C13—C12	-176.5 (3)
C4—C5—C9—C8	178.9 (3)	C15—C14—C13—C12	1.5 (4)
C6—C5—C9—C8	-2.8 (4)	C11—N4—C10—N3	74.2 (4)
C15—N6—C16—C17	-1.9 (4)	C7—N3—C10—N4	85.6 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O6 ⁱ	0.88 (4)	1.92	2.794 (4)	168
N3—H3A \cdots O3	0.80 (3)	2.23	2.990 (6)	159
N4—H4A \cdots O5 ⁱ	0.71 (4)	2.56	3.233 (5)	160
N4—H4A \cdots O7 ⁱ	0.71 (4)	2.52	3.133 (5)	145
N6—H6 \cdots O1	0.84 (3)	2.00	2.838 (8)	178
C1—H1A \cdots O7 ⁱⁱ	0.98	2.54	3.501 (3)	167
C1—H1B \cdots O4 ⁱⁱⁱ	0.98	2.33	3.078 (4)	132
C4—H4 \cdots O8 ^{iv}	0.95	2.52	3.392 (7)	152
C10—H10B \cdots O4 ^{iv}	0.99	2.35	3.082 (5)	130
C13—H13 \cdots O2 ⁱ	0.95	2.41	3.259 (3)	149
C19—H19B \cdots O5	0.98	2.57	3.351 (6)	136
C19—H19C \cdots O7 ^v	0.98	2.58	3.207 (7)	122
C19—H19C \cdots O8 ^v	0.98	2.57	3.548 (4)	172

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x+3/2, -y, z-1/2$; (iii) $x, -y+1/2, z-1/2$; (iv) $x-1/2, -y+1/2, -z+1$; (v) $x-1, y, z$.

Fig. 1




Fig. 2

