

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

{2-[(2,5-Dimethylphenyl)iminomethyl]-pyridine- κ^2N,N' }diiodidozinc(II)

 Mohamad Reza Talei Olyai,^{a*} Saeed Dehghanpour,^b Bitā Hoormehr,^c Fahimeh Gholami^c and Hamid Reza Khavasi^d
^aDepartment of Chemistry, Islamic Azad University, South Tehran Branch, Tehran, Iran, ^bDepartment of Chemistry, Alzahra University, Vanak, Tehran, Iran,

^cDepartment of Chemistry, Islamic Azad University, Karaj Branch, Tehran, Iran, and

^dDepartment of Chemistry, Shahid Beheshti University, Evin, Tehran 1983963113, Iran

Correspondence e-mail: talei3@yahoo.com


Received 30 June 2008; accepted 2 July 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(C-C) = 0.009$ Å; R factor = 0.072; wR factor = 0.230; data-to-parameter ratio = 26.2.

In the molecule of the title compound, $[ZnI_2(C_{14}H_{14}N_2)]$, the Zn atom is four-coordinated in a distorted tetrahedral geometry by two N atoms of the Schiff base ligand and by two I atoms. The benzene and pyridine rings are oriented at a dihedral angle of $70.75(3)^\circ$. The five-membered ring has an envelope conformation. There is a weak $\pi-\pi$ interaction between benzene rings, with a centroid-to-centroid distance of $3.975(4)$ Å.

Related literature

For general background, see: Gibson *et al.* (2007); Ittel *et al.* (2000); Gibson & Spitzmesser (2003); Bart *et al.* (2004); Sugiyama *et al.* (2004); Kooistra *et al.* (2004); Bouwkamp *et al.* (2006). For related literature, see: Dehghanpour *et al.* (2007).



Experimental

Crystal data

 $[ZnI_2(C_{14}H_{14}N_2)]$
 $M_r = 529.46$

 Monoclinic, $P2_1/c$
 $a = 11.467(5)$ Å

 $b = 9.627(4)$ Å
 $c = 15.868(6)$ Å
 $\beta = 103.88(3)^\circ$
 $V = 1700.6(12)$ Å³
 $Z = 4$

 Mo $K\alpha$ radiation
 $\mu = 5.06$ mm⁻¹
 $T = 298(2)$ K
 $0.5 \times 0.4 \times 0.25$ mm

Data collection

 Stoe IPDSII diffractometer
 Absorption correction: numerical
 [$X-RED32$ and $X-SHAPE$ (Stoe & Cie, 2005)]
 $T_{min} = 0.100$, $T_{max} = 0.280$

 10947 measured reflections
 4498 independent reflections
 4035 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.098$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.071$
 $wR(F^2) = 0.230$
 $S = 1.12$
 4498 reflections

 172 parameters
 H-atom parameters constrained
 $\Delta\rho_{max} = 1.97$ e Å⁻³
 $\Delta\rho_{min} = -1.66$ e Å⁻³
Table 1

Selected geometric parameters (Å, °).

Zn1–I1	2.5201 (11)	Zn1–N1	2.059 (5)
Zn1–I2	2.5517 (12)	Zn1–N2	2.104 (5)
I1–Zn1–I2	120.26 (4)	N2–Zn1–I1	107.27 (13)
N1–Zn1–N2	80.16 (19)	N1–Zn1–I2	107.58 (15)
N1–Zn1–I1	112.45 (14)	N2–Zn1–I2	121.99 (13)

Data collection: $X-AREA$ (Stoe & Cie, 2005); cell refinement: $X-AREA$; data reduction: $X-RED32$ (Stoe & Cie, 2005); program(s) used to solve structure: $SHELXS97$ (Sheldrick, 2008); program(s) used to refine structure: $SHELXL97$ (Sheldrick, 2008); molecular graphics: $ORTEP-3$ for Windows (Farrugia, 1997); software used to prepare material for publication: $WinGX$ (Farrugia, 1999).

The authors acknowledge Isfahan University of Technology and Shahid Beheshti University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2486).

References

- Bart, S. C., Lobkovsky, E. & Chirik, P. J. (2004). *J. Am. Chem. Soc.* **126**, 13794–13807.
- Bouwkamp, M. W., Lobkovsky, E. & Chirik, P. J. (2006). *Inorg. Chem.* **45**, 2–4.
- Dehghanpour, S., Mahmoudi, A., Khalaj, M., Salmanpour, S. & Adib, M. (2007). *Acta Cryst.* **E63**, m2841.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gibson, V. C., Redshaw, C. & Solan, G. A. (2007). *Chem. Rev.* **107**, 1745–1776.
- Gibson, V. C. & Spitzmesser, S. K. (2003). *Chem. Rev.* **103**, 283–316.
- Ittel, S. D., Johnson, L. K. & Brookhart, M. (2000). *Chem. Rev.* **100**, 1169–1203.
- Kooistra, T. M., Hettterscheid, D. G. H., Schwartz, E., Knijnenburg, Q., Budzelaar, P. H. M. & Gal, A. W. (2004). *Inorg. Chim. Acta*, **357**, 2945–2952.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Stoe & Cie (2005). $X-AREA$, $X-RED32$ and $X-SHAPE$. Stoe & Cie, Darmstadt, Germany.
- Sugiyama, I., Korobkov, I., Gambarotta, S., Mueller, A. & Budzelaar, P. H. M. (2004). *Inorg. Chem.* **43**, 5771–5779.

supplementary materials

Acta Cryst. (2008). E64, m1191 [doi:10.1107/S1600536808020230]

{2-[(2,5-Dimethylphenyl)iminomethyl]pyridine- κ^2N,N' }diiodidozinc(II)

M. R. Talei Babil Olyai, S. Dehghanpour, B. Hoormehr, F. Gholami and H. R. Khavasi

Comment

Transition metal compounds containing Schiff base ligands have been of great interest for many years. These compounds play an important role in the development of coordination chemistry (Gibson *et al.*, 2007). Aryl-substituted iminopyridine complexes have emerged as a powerful class of catalysts for a host of important bond-forming reactions including olefin polymerization (Ittel *et al.*, 2000; Gibson & Spitzmesser, 2003), hydrogenation and hydrosilation (Bart *et al.*, 2004). It is also now well established that iminopyridines are both redox (Sugiyama *et al.*, 2004) and chemically active ligands (Kooistra *et al.*, 2004) participating in electron transfer, addition reactions and deprotonation chemistry (Bouwkamp *et al.*, 2006). We report herein the crystal structure of the title compound.

In the molecule of the title compound (Fig. 1), the Zn atom is four-coordinated in distorted tetrahedral geometry (Table 1) by two N atoms of the Schiff base ligand, and two I atoms. The bond angles around the Zn atom deviate from the ideal tetrahedral geometry (Dehghanpour *et al.*, 2007). Rings A (N1/C1–C5) and B (C7/C8/C10/C11/C12/C14) are, of course, planar, and they are oriented at a dihedral angle of 70.75 (3)°. Ring C (Zn1/N1/N2/C5/C6) has envelope conformation, with N2 atom displaced by 0.104 (3) Å from the plane of the other ring atoms. The weak π – π interaction between B rings $\text{CgB} \cdots \text{CgB}^i$ [symmetry code: (iv) 1 - x, 1 - y, 1 - z] may stabilize the structure, with a centroid–centroid distance of 3.975 (4) Å.

Experimental

For the preparation of the title compound, (2,5-dimethyl-N-phenyl)(pyridine-2-yl)methanimine (21.0 mg, 0.1 mmol), and ZnI_2 (31.9 mg, 0.1 mmol) were dissolved in acetonitrile (50 ml). The mixture was stirred for 10 min at room temperature. The resulting solution was left in air for a few days, giving yellow crystals of the title compound (yield; 79%). Calc.: C 31.75 H 2.67, N 5.29%, found: C 31.79, H 2.49, N 5.31%.

Refinement

H atoms were positioned geometrically, with C–H = 0.93 and 0.96 Å for aromatic and methyl H, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

{2-[(2,5-Dimethylphenyl)iminomethyl]pyridine- $\kappa^2\text{N},\text{N}'$ }diiodidozinc(II)

Crystal data

[ZnI ₂ (C ₁₄ H ₁₄ N ₂)]	$F_{000} = 992$
$M_r = 529.46$	$D_x = 2.068 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
Hall symbol: -P 2ybc	$\lambda = 0.71073 \text{ \AA}$
$a = 11.467 (5) \text{ \AA}$	Cell parameters from 2112 reflections
$b = 9.627 (4) \text{ \AA}$	$\theta = 2.5\text{--}29.4^\circ$
$c = 15.868 (6) \text{ \AA}$	$\mu = 5.06 \text{ mm}^{-1}$
$\beta = 103.88 (3)^\circ$	$T = 298 (2) \text{ K}$
$V = 1700.6 (12) \text{ \AA}^3$	Block, yellow
$Z = 4$	$0.5 \times 0.4 \times 0.25 \text{ mm}$

Data collection

Stoe IPDSII diffractometer	$R_{\text{int}} = 0.098$
rotation method scans	$\theta_{\text{max}} = 29.4^\circ$
Absorption correction: numerical [X-RED and X-SHAPE (Stoe & Cie, 2005)]	$\theta_{\text{min}} = 2.5^\circ$
$T_{\text{min}} = 0.100$, $T_{\text{max}} = 0.280$	$h = -15 \rightarrow 11$
10947 measured reflections	$k = -11 \rightarrow 13$
4498 independent reflections	$l = -21 \rightarrow 21$
4035 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.1439P)^2 + 2.419P]$
$R[F^2 > 2\sigma(F^2)] = 0.071$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.230$	$(\Delta/\sigma)_{\text{max}} = 0.007$
$S = 1.12$	$\Delta\rho_{\text{max}} = 1.97 \text{ e \AA}^{-3}$
4498 reflections	$\Delta\rho_{\text{min}} = -1.66 \text{ e \AA}^{-3}$
172 parameters	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.83553 (6)	0.25679 (7)	0.02157 (4)	0.0406 (2)
I1	0.69958 (5)	0.45620 (5)	0.03661 (4)	0.0622 (2)
I2	0.87622 (5)	0.19974 (6)	-0.12580 (3)	0.0617 (2)
N1	0.9989 (4)	0.2665 (5)	0.1106 (3)	0.0400 (9)
N2	0.8103 (4)	0.1007 (5)	0.1083 (3)	0.0366 (8)
C1	1.0919 (6)	0.3467 (9)	0.1080 (5)	0.0524 (14)
H1	1.086	0.4083	0.062	0.063*
C2	1.1962 (6)	0.3412 (9)	0.1712 (5)	0.0590 (18)
H2	1.2614	0.3967	0.1681	0.071*
C3	1.2026 (6)	0.2495 (9)	0.2411 (5)	0.0596 (19)
H3	1.2721	0.2453	0.2856	0.072*
C4	1.1071 (5)	0.1665 (8)	0.2440 (4)	0.0498 (13)
H4	1.1104	0.1042	0.2893	0.06*
C5	1.0055 (5)	0.1786 (6)	0.1771 (4)	0.0411 (11)
C6	0.8989 (5)	0.0951 (6)	0.1752 (4)	0.0412 (11)
H6	0.8952	0.0384	0.2219	0.049*
C7	0.7036 (5)	0.0265 (5)	0.1120 (3)	0.0356 (9)
C8	0.6663 (5)	-0.0814 (6)	0.0540 (3)	0.0377 (10)
C9	0.7345 (7)	-0.1256 (8)	-0.0114 (5)	0.0531 (14)
H9A	0.8136	-0.1554	0.0183	0.064*
H9B	0.7404	-0.0488	-0.0487	0.064*
H9C	0.693	-0.201	-0.0455	0.064*
C10	0.5634 (5)	-0.1522 (6)	0.0595 (4)	0.0440 (12)
H10	0.5352	-0.2247	0.0213	0.053*
C11	0.5023 (5)	-0.1160 (7)	0.1214 (5)	0.0495 (13)
H11	0.4335	-0.1652	0.1241	0.059*
C12	0.5402 (5)	-0.0092 (8)	0.1791 (4)	0.0491 (13)
C13	0.4725 (8)	0.0300 (13)	0.2483 (7)	0.076 (3)
H13A	0.4453	0.1244	0.2396	0.091*
H13B	0.5253	0.0207	0.305	0.091*
H13C	0.4047	-0.0304	0.2435	0.091*
C14	0.6415 (5)	0.0638 (6)	0.1732 (4)	0.0416 (11)
H14	0.668	0.138	0.2104	0.05*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0414 (4)	0.0406 (4)	0.0392 (4)	-0.0024 (2)	0.0082 (3)	0.0058 (2)
I1	0.0607 (3)	0.0455 (3)	0.0819 (4)	0.00774 (18)	0.0200 (3)	0.0016 (2)
I2	0.0745 (4)	0.0713 (4)	0.0442 (3)	0.0013 (2)	0.0239 (2)	0.00481 (18)
N1	0.036 (2)	0.038 (2)	0.047 (2)	-0.0030 (17)	0.0124 (18)	-0.0020 (18)
N2	0.041 (2)	0.033 (2)	0.0357 (19)	0.0000 (16)	0.0107 (16)	0.0014 (16)
C1	0.048 (3)	0.060 (4)	0.052 (3)	-0.015 (3)	0.019 (3)	-0.006 (3)
C2	0.039 (3)	0.076 (5)	0.064 (4)	-0.011 (3)	0.016 (3)	-0.030 (4)

supplementary materials

C3	0.039 (3)	0.079 (5)	0.055 (4)	0.001 (3)	0.001 (3)	-0.032 (4)
C4	0.040 (3)	0.054 (3)	0.050 (3)	0.003 (2)	0.001 (2)	-0.006 (3)
C5	0.041 (2)	0.040 (3)	0.041 (2)	0.000 (2)	0.006 (2)	-0.007 (2)
C6	0.040 (2)	0.037 (2)	0.044 (3)	0.0001 (19)	0.005 (2)	0.006 (2)
C7	0.036 (2)	0.031 (2)	0.037 (2)	0.0000 (17)	0.0042 (17)	0.0039 (17)
C8	0.045 (2)	0.033 (2)	0.034 (2)	0.0013 (19)	0.0083 (19)	0.0027 (18)
C9	0.066 (4)	0.048 (3)	0.050 (3)	-0.003 (3)	0.023 (3)	-0.011 (3)
C10	0.040 (2)	0.036 (2)	0.051 (3)	-0.005 (2)	0.002 (2)	0.001 (2)
C11	0.037 (2)	0.050 (3)	0.060 (3)	-0.002 (2)	0.008 (2)	0.006 (3)
C12	0.033 (2)	0.064 (4)	0.051 (3)	0.002 (2)	0.010 (2)	0.001 (3)
C13	0.054 (4)	0.105 (7)	0.077 (5)	0.010 (4)	0.031 (4)	-0.010 (5)
C14	0.042 (3)	0.042 (3)	0.040 (2)	-0.002 (2)	0.008 (2)	-0.006 (2)

Geometric parameters (Å, °)

Zn1—I1	2.5201 (11)	C7—C8	1.387 (7)
Zn1—I2	2.5517 (12)	C7—N2	1.430 (7)
Zn1—N1	2.059 (5)	C8—C10	1.384 (8)
Zn1—N2	2.104 (5)	C8—C9	1.502 (9)
C1—N1	1.326 (8)	C9—H9A	0.96
C1—C2	1.365 (10)	C9—H9B	0.96
C1—H1	0.93	C9—H9C	0.96
C2—C3	1.406 (13)	C10—C11	1.381 (10)
C2—H2	0.93	C10—H10	0.93
C3—C4	1.365 (11)	C11—C12	1.377 (10)
C3—H3	0.93	C11—H11	0.93
C4—C5	1.380 (8)	C12—C14	1.380 (9)
C4—H4	0.93	C12—C13	1.535 (11)
C5—N1	1.341 (8)	C13—H13A	0.96
C5—C6	1.457 (8)	C13—H13B	0.96
C6—N2	1.283 (7)	C13—H13C	0.96
C6—H6	0.93	C14—H14	0.93
C7—C14	1.382 (8)		
I1—Zn1—I2	120.26 (4)	C14—C7—C8	122.2 (5)
N1—Zn1—N2	80.16 (19)	C14—C7—N2	119.3 (5)
N1—Zn1—I1	112.45 (14)	C8—C7—N2	118.5 (5)
N2—Zn1—I1	107.27 (13)	C10—C8—C7	117.3 (5)
N1—Zn1—I2	107.58 (15)	C10—C8—C9	119.8 (6)
N2—Zn1—I2	121.99 (13)	C7—C8—C9	122.9 (5)
C1—N1—C5	119.8 (6)	C8—C9—H9A	109.5
C1—N1—Zn1	127.7 (5)	C8—C9—H9B	109.5
C5—N1—Zn1	112.4 (4)	H9A—C9—H9B	109.5
C6—N2—C7	117.4 (5)	C8—C9—H9C	109.5
C6—N2—Zn1	111.5 (4)	H9A—C9—H9C	109.5
C7—N2—Zn1	129.6 (3)	H9B—C9—H9C	109.5
N1—C1—C2	121.7 (7)	C11—C10—C8	120.4 (6)
N1—C1—H1	119.2	C11—C10—H10	119.8
C2—C1—H1	119.2	C8—C10—H10	119.8
C1—C2—C3	118.3 (7)	C12—C11—C10	121.9 (6)

C1—C2—H2	120.8	C12—C11—H11	119
C3—C2—H2	120.8	C10—C11—H11	119
C4—C3—C2	120.2 (6)	C11—C12—C14	118.1 (6)
C4—C3—H3	119.9	C11—C12—C13	121.8 (7)
C2—C3—H3	119.9	C14—C12—C13	120.1 (7)
C3—C4—C5	117.4 (7)	C12—C13—H13A	109.5
C3—C4—H4	121.3	C12—C13—H13B	109.5
C5—C4—H4	121.3	H13A—C13—H13B	109.5
N1—C5—C4	122.5 (6)	C12—C13—H13C	109.5
N1—C5—C6	116.3 (5)	H13A—C13—H13C	109.5
C4—C5—C6	121.2 (6)	H13B—C13—H13C	109.5
N2—C6—C5	119.2 (5)	C12—C14—C7	120.0 (6)
N2—C6—H6	120.4	C12—C14—H14	120
C5—C6—H6	120.4	C7—C14—H14	120
N1—C1—C2—C3	-1.3 (11)	C4—C5—N1—C1	-0.7 (9)
C1—C2—C3—C4	1.4 (11)	C6—C5—N1—C1	179.5 (6)
C2—C3—C4—C5	-1.1 (10)	C4—C5—N1—Zn1	179.3 (5)
C3—C4—C5—N1	0.8 (10)	C6—C5—N1—Zn1	-0.5 (6)
C3—C4—C5—C6	-179.5 (6)	N2—Zn1—N1—C1	177.4 (6)
N1—C5—C6—N2	5.8 (8)	I1—Zn1—N1—C1	-77.9 (6)
C4—C5—C6—N2	-174.0 (6)	I2—Zn1—N1—C1	56.8 (6)
C14—C7—C8—C10	-0.2 (8)	N2—Zn1—N1—C5	-2.6 (4)
N2—C7—C8—C10	179.5 (5)	I1—Zn1—N1—C5	102.1 (4)
C14—C7—C8—C9	-178.4 (6)	I2—Zn1—N1—C5	-123.2 (4)
N2—C7—C8—C9	1.3 (8)	C5—C6—N2—C7	-174.8 (5)
C7—C8—C10—C11	-0.6 (8)	C5—C6—N2—Zn1	-7.7 (7)
C9—C8—C10—C11	177.6 (6)	C14—C7—N2—C6	62.5 (7)
C8—C10—C11—C12	0.3 (10)	C8—C7—N2—C6	-117.1 (6)
C10—C11—C12—C14	0.9 (10)	C14—C7—N2—Zn1	-101.8 (6)
C10—C11—C12—C13	-179.5 (7)	C8—C7—N2—Zn1	78.6 (6)
C11—C12—C14—C7	-1.7 (9)	N1—Zn1—N2—C6	5.6 (4)
C13—C12—C14—C7	178.7 (7)	I1—Zn1—N2—C6	-105.0 (4)
C8—C7—C14—C12	1.4 (9)	I2—Zn1—N2—C6	110.3 (4)
N2—C7—C14—C12	-178.3 (5)	N1—Zn1—N2—C7	170.6 (5)
C2—C1—N1—C5	0.9 (10)	I1—Zn1—N2—C7	60.0 (5)
C2—C1—N1—Zn1	-179.0 (5)	I2—Zn1—N2—C7	-84.7 (5)

Fig. 1

