

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-{[2,5-Dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]aminocarbonyl}-2,6-difluorobenzamide

Yin-hong Liu, Fang-shi Li* and Yi Li

Department of Applied Chemistry, College of Science, Nanjing University of Technology, Xinmofan Road No. 5, Nanjing 210009, People's Republic of China
Correspondence e-mail: fangshi.li@njut.edu.cn


Received 6 August 2008; accepted 8 August 2008

Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.086; wR factor = 0.248; data-to-parameter ratio = 12.8.

In the molecule of the title compound, $\text{C}_{17}\text{H}_8\text{Cl}_2\text{F}_8\text{N}_2\text{O}_3$, the two aromatic rings are oriented at a dihedral angle of $50.12(3)^\circ$. Intramolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds result in the formation of two six- and one five-membered rings. The six-membered rings have flattened-boat conformations, while the five-membered ring adopts an envelope conformation. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into centrosymmetric dimers.

Related literature

For related literature, see: Drabek & Boeger (1986). For bond-length data, see: Allen *et al.* (1987). For ring conformation puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{17}\text{H}_8\text{Cl}_2\text{F}_8\text{N}_2\text{O}_3$
 $M_r = 511.15$

Monoclinic, $P2_1/n$
 $a = 9.2300(18)$ Å

$b = 16.404(3)$ Å
 $c = 14.074(3)$ Å
 $\beta = 108.77(3)^\circ$
 $V = 2017.6(8)$ Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.42$ mm⁻¹
 $T = 294(2)$ K
 $0.40 \times 0.30 \times 0.20$ mm

Data collection

Enraf-Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.851$, $T_{\max} = 0.921$
3609 measured reflections

3609 independent reflections
1922 reflections with $I > 2\sigma(I)$
3 standard reflections every 200 reflections
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.086$
 $wR(F^2) = 0.247$
 $S = 1.09$
3609 reflections
283 parameters

96 restraints
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.42$ e Å⁻³
 $\Delta\rho_{\min} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H2A}\cdots\text{Cl2}$	0.86	2.43	2.892 (4)	115
$\text{N2}-\text{H2A}\cdots\text{O1}$	0.86	1.98	2.664 (6)	135
$\text{N1}-\text{H1A}\cdots\text{O2}^i$	0.86	1.98	2.814 (6)	163
$\text{C10}-\text{H10A}\cdots\text{O2}$	0.93	2.28	2.851 (6)	119

Symmetry code: (i) $-x + 1, -y + 2, -z + 1$.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2510).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
Drabek, J. & Boeger, M. (1986). European Patent No. 179022.
Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o1756 [doi:10.1107/S1600536808025506]

N-{[2,5-Dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]aminocarbonyl}-2,6-difluorobenzamide

Y. Liu, F. Li and Y. Li

Comment

The title compound is considered to belong to the fourth generation of insecticides with properties such as high selectivity, low acute toxicity for mammals and high biological activity. It is generally recognized as a chitin-synthesis inhibitor that interrupts chitin-synthesis during the development and reproduction of the insecticide. As part of our studies in this area, we report herein the crystal structure of the title compound.

In the molecule of the title compound, (Fig. 1) the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (C1-C6) and B (C9-C14) are, of course, planar, and the dihedral angle between them is A/B = 50.12 (3)°. The intramolecular N-H···O, C-H···O and N-H···Cl hydrogen bonds (Table 1) result in the formation of two six- and one five-membered non-planar rings: C (O1/N1/N2/C7/C8/H2A), D (O2/N2/C8-C10/H10A) and E (Cl2/N2/C9/C14/H2A). Rings C and D adopt twisted [$\varphi = -169.19$ (2)°, $\theta = 21.09$ (3)° (for ring C) and $\varphi = 178.48$ (3)°, $\theta = 127.74$ (3)° (for ring D)] conformations, having total puckering amplitudes, Q_T , of 0.113 (3) and 0.201 (3) Å, respectively (Cremer & Pople, 1975). Ring E adopts envelope conformation, with H2A atom displaced by 0.190 (3) Å from the plane of the other ring atoms.

In the crystal structure, intermolecular N-H···O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers (Fig. 2), in which they may be effective in the stabilization of the structure.

Experimental

The title compound was prepared according to the literature method (Drabek & Boeger, 1986). The crystals suitable for X-ray analysis were obtained by dissolving the title compound (0.3 g) in acetonitrile (25 ml) and evaporating the solvent slowly at room temperature for about 8 d.

Refinement

H atoms were positioned geometrically, with N-H = 0.86 Å (for NH) and C-H = 0.93 and 0.98 Å for aromatic and methine H, respectively, and constrained to ride on their parent atoms with $U_{iso}(H) = 1.2U_{eq}(C,N)$.

Figures




Fig. 1. The molecular structure of the title compound, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines.




Fig. 2. A packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

***N*-{[2,5-Dichloro-4-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]aminocarbonyl}-2,6-difluorobenzamide**

Crystal data

$C_{17}H_8Cl_2F_8N_2O_3$

$M_r = 511.15$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2yn$

$a = 9.2300$ (18) Å

$b = 16.404$ (3) Å

$c = 14.074$ (3) Å

$\beta = 108.77$ (3)°

$V = 2017.6$ (8) Å³

$Z = 4$

$F_{000} = 1016$

$D_x = 1.683$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$

$\mu = 0.42$ mm⁻¹

$T = 294$ (2) K

Block, colorless

$0.40 \times 0.30 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 294$ (2) K

$\omega/2\theta$ scans

Absorption correction: ψ scan
(North *et al.*, 1968)

$T_{\min} = 0.851$, $T_{\max} = 0.921$

3609 measured reflections

3609 independent reflections

1922 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.0000$

$\theta_{\max} = 25.2^\circ$

$\theta_{\min} = 2.0^\circ$

$h = -11 \rightarrow 10$

$k = 0 \rightarrow 19$

$l = 0 \rightarrow 16$

3 standard reflections

every 200 reflections

intensity decay: none

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.086$

$wR(F^2) = 0.248$

$S = 1.09$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1258P)^2 + 1.1819P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

3609 reflections $\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$
 283 parameters $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$
 96 restraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	1.21022 (15)	1.15534 (9)	0.61828 (12)	0.0527 (5)
C12	1.06547 (19)	0.83402 (11)	0.80001 (17)	0.0914 (8)
F1	0.4001 (5)	0.7889 (3)	0.7184 (3)	0.0975 (16)
F2	0.4224 (4)	0.8023 (3)	0.3906 (3)	0.0810 (13)
F3	1.5260 (5)	1.0359 (3)	0.6887 (5)	0.1116 (18)
F4	1.5722 (4)	0.9618 (3)	0.8001 (4)	0.1039 (17)
F5	1.8146 (6)	1.0579 (3)	0.7942 (5)	0.145 (3)
F6	1.8127 (6)	1.0156 (3)	0.9703 (4)	0.1121 (18)
F7	1.9281 (6)	1.1152 (3)	0.9350 (4)	0.119 (2)
F8	1.6616 (6)	1.1347 (3)	0.9601 (4)	0.1029 (15)
O1	0.6850 (4)	0.7921 (2)	0.6546 (4)	0.0703 (15)
O2	0.6981 (4)	1.0216 (3)	0.5376 (3)	0.0505 (11)
O3	1.4248 (4)	1.0694 (3)	0.7916 (4)	0.0758 (16)
N1	0.5870 (4)	0.9045 (2)	0.5577 (3)	0.0359 (11)
H1A	0.5033	0.9242	0.5175	0.043*
N2	0.8483 (4)	0.9219 (2)	0.6377 (3)	0.0327 (10)
H2A	0.8468	0.8724	0.6576	0.039*
C1	0.1954 (8)	0.7296 (5)	0.5816 (6)	0.074 (2)
H1B	0.1414	0.7140	0.6242	0.089*
C2	0.1405 (7)	0.7183 (4)	0.4832 (5)	0.0588 (17)
H2B	0.0444	0.6945	0.4569	0.071*
C3	0.2142 (7)	0.7389 (4)	0.4199 (5)	0.0607 (17)
H3A	0.1704	0.7275	0.3519	0.073*
C4	0.3505 (6)	0.7758 (4)	0.4526 (4)	0.0450 (14)
C5	0.4228 (6)	0.7893 (3)	0.5551 (4)	0.0435 (14)
C6	0.3397 (7)	0.7666 (4)	0.6175 (5)	0.0568 (16)
C7	0.5790 (6)	0.8247 (3)	0.5960 (4)	0.0437 (14)

supplementary materials

C8	0.7151 (6)	0.9534 (3)	0.5785 (4)	0.0407 (14)
C9	0.9852 (5)	0.9614 (3)	0.6689 (4)	0.0330 (12)
C10	1.0214 (5)	1.0332 (3)	0.6291 (4)	0.0368 (12)
H10A	0.9477	1.0595	0.5770	0.044*
C11	1.1661 (5)	1.0655 (3)	0.6667 (4)	0.0355 (12)
C12	1.2807 (6)	1.0310 (4)	0.7494 (5)	0.0518 (16)
C13	1.2445 (7)	0.9591 (4)	0.7901 (5)	0.0572 (17)
H13A	1.3159	0.9350	0.8453	0.069*
C14	1.1065 (5)	0.9252 (3)	0.7492 (4)	0.0391 (13)
C15	1.5460 (7)	1.0386 (4)	0.8024 (7)	0.079 (3)
C16	1.6854 (8)	1.0845 (5)	0.8271 (9)	0.107 (4)
H16A	1.6576	1.1394	0.7999	0.129*
C17	1.7772 (11)	1.0946 (6)	0.9319 (7)	0.089

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0401 (8)	0.0436 (9)	0.0782 (11)	-0.0152 (6)	0.0242 (7)	-0.0007 (7)
C12	0.0513 (10)	0.0599 (12)	0.1330 (18)	-0.0100 (8)	-0.0123 (10)	0.0497 (12)
F1	0.097 (3)	0.135 (4)	0.051 (3)	-0.054 (3)	0.010 (2)	0.026 (2)
F2	0.086 (3)	0.103 (3)	0.059 (3)	-0.045 (3)	0.030 (2)	-0.034 (2)
F3	0.099 (4)	0.106 (4)	0.133 (5)	-0.034 (3)	0.041 (3)	-0.032 (3)
F4	0.054 (2)	0.072 (3)	0.172 (5)	-0.010 (2)	0.019 (3)	-0.015 (3)
F5	0.092 (3)	0.141 (5)	0.225 (7)	-0.023 (3)	0.082 (4)	-0.111 (5)
F6	0.127 (4)	0.104 (4)	0.095 (4)	-0.017 (3)	0.021 (3)	0.052 (3)
F7	0.099 (3)	0.083 (3)	0.116 (4)	-0.024 (3)	-0.048 (3)	0.006 (3)
F8	0.097 (3)	0.111 (4)	0.103 (4)	-0.001 (3)	0.034 (3)	-0.012 (3)
O1	0.029 (2)	0.047 (3)	0.107 (4)	0.0006 (18)	-0.017 (2)	0.037 (2)
O2	0.046 (2)	0.047 (3)	0.057 (3)	-0.0033 (19)	0.013 (2)	-0.005 (2)
O3	0.035 (2)	0.043 (3)	0.129 (5)	-0.009 (2)	-0.001 (2)	0.009 (3)
N1	0.028 (2)	0.019 (2)	0.058 (3)	0.0007 (17)	0.009 (2)	0.0093 (19)
N2	0.028 (2)	0.025 (2)	0.043 (3)	-0.0005 (17)	0.0085 (19)	-0.0044 (19)
C1	0.070 (4)	0.082 (5)	0.076 (4)	-0.028 (4)	0.032 (4)	0.017 (4)
C2	0.046 (3)	0.051 (4)	0.071 (4)	-0.016 (3)	0.007 (3)	0.003 (3)
C3	0.059 (4)	0.061 (4)	0.051 (4)	-0.023 (3)	0.002 (3)	-0.014 (3)
C4	0.045 (3)	0.046 (3)	0.042 (3)	-0.003 (3)	0.011 (3)	-0.004 (3)
C5	0.039 (3)	0.042 (3)	0.047 (3)	-0.006 (2)	0.011 (2)	0.010 (3)
C6	0.044 (3)	0.067 (4)	0.053 (4)	-0.009 (3)	0.007 (3)	0.014 (3)
C7	0.045 (3)	0.039 (3)	0.047 (4)	-0.010 (3)	0.016 (3)	0.004 (3)
C8	0.037 (3)	0.027 (3)	0.051 (4)	0.000 (2)	0.005 (3)	0.009 (3)
C9	0.026 (2)	0.036 (3)	0.036 (3)	0.000 (2)	0.009 (2)	-0.005 (2)
C10	0.032 (2)	0.040 (3)	0.042 (3)	-0.007 (2)	0.016 (2)	-0.003 (2)
C11	0.033 (3)	0.035 (3)	0.044 (3)	-0.009 (2)	0.020 (2)	0.008 (2)
C12	0.030 (3)	0.053 (4)	0.065 (4)	-0.003 (2)	0.006 (3)	0.004 (3)
C13	0.042 (3)	0.051 (4)	0.071 (4)	0.013 (3)	0.007 (3)	0.017 (3)
C14	0.022 (2)	0.042 (3)	0.049 (3)	0.012 (2)	0.005 (2)	0.009 (2)
C15	0.029 (3)	0.050 (5)	0.142 (8)	0.006 (3)	0.007 (4)	-0.014 (4)
C16	0.041 (4)	0.046 (4)	0.219 (11)	-0.009 (3)	0.021 (5)	0.007 (6)

C17 0.080 0.089 0.096 -0.006 0.026 0.004

Geometric parameters (Å, °)

C11—C11	1.727 (5)	C2—C3	1.326 (9)
C12—C14	1.751 (6)	C2—H2B	0.9300
F1—C6	1.398 (7)	O3—C15	1.191 (7)
F2—C4	1.327 (7)	O3—C12	1.418 (6)
F3—C15	1.552 (10)	C3—C4	1.338 (8)
F4—C15	1.284 (8)	C3—H3A	0.9300
F5—C16	1.477 (9)	C4—C5	1.399 (8)
F6—C17	1.403 (10)	C5—C6	1.389 (8)
F7—C17	1.420 (10)	C5—C7	1.488 (7)
F8—C17	1.415 (10)	C9—C10	1.390 (7)
O1—C7	1.186 (6)	C9—C14	1.439 (7)
N1—C8	1.380 (6)	C10—C11	1.374 (7)
N1—C7	1.427 (7)	C10—H10A	0.9300
N1—H1A	0.8600	C11—C12	1.414 (7)
C1—C2	1.327 (10)	C12—C13	1.398 (8)
C1—C6	1.401 (9)	C13—C14	1.338 (8)
C1—H1B	0.9300	C13—H13A	0.9300
N2—C8	1.347 (6)	C15—C16	1.434 (9)
N2—C9	1.361 (6)	C16—C17	1.455 (13)
N2—H2A	0.8600	C16—H16A	0.9800
O2—C8	1.245 (6)		
C8—N1—C7	126.7 (4)	C11—C10—C9	120.1 (5)
C8—N1—H1A	116.6	C11—C10—H10A	120.0
C7—N1—H1A	116.6	C9—C10—H10A	120.0
C2—C1—C6	115.8 (6)	C10—C11—C12	122.7 (5)
C2—C1—H1B	122.1	C10—C11—C11	120.0 (4)
C6—C1—H1B	122.1	C12—C11—C11	117.2 (4)
C8—N2—C9	125.8 (4)	C13—C12—C11	117.4 (5)
C8—N2—H2A	117.1	C13—C12—O3	121.2 (5)
C9—N2—H2A	117.1	C11—C12—O3	121.4 (5)
C1—C2—C3	124.0 (6)	C14—C13—C12	119.7 (5)
C1—C2—H2B	118.0	C14—C13—H13A	120.2
C3—C2—H2B	118.0	C12—C13—H13A	120.2
C15—O3—C12	125.5 (6)	C13—C14—C9	123.9 (5)
C2—C3—C4	120.9 (6)	C13—C14—C12	118.8 (4)
C2—C3—H3A	119.6	C9—C14—C12	117.2 (4)
C4—C3—H3A	119.6	O3—C15—F4	126.1 (6)
F2—C4—C3	122.4 (6)	O3—C15—C16	122.7 (7)
F2—C4—C5	117.0 (5)	F4—C15—C16	111.1 (6)
C3—C4—C5	120.7 (6)	O3—C15—F3	94.9 (6)
C6—C5—C4	115.6 (5)	F4—C15—F3	84.7 (6)
C6—C5—C7	121.6 (5)	C16—C15—F3	93.7 (7)
C4—C5—C7	122.8 (5)	C15—C16—C17	119.4 (9)
C5—C6—F1	116.8 (5)	C15—C16—F5	121.4 (7)
C5—C6—C1	122.9 (6)	C17—C16—F5	94.9 (6)

supplementary materials

F1—C6—C1	120.1 (6)	C15—C16—H16A	106.6
O1—C7—N1	123.1 (5)	C17—C16—H16A	106.6
O1—C7—C5	125.0 (5)	F5—C16—H16A	106.6
N1—C7—C5	111.9 (5)	F6—C17—F8	115.2 (7)
O2—C8—N2	125.8 (5)	F6—C17—F7	95.9 (6)
O2—C8—N1	116.9 (4)	F8—C17—F7	134.6 (8)
N2—C8—N1	117.2 (5)	F6—C17—C16	105.8 (7)
N2—C9—C10	126.3 (5)	F8—C17—C16	95.3 (7)
N2—C9—C14	117.6 (5)	F7—C17—C16	107.7 (7)
C10—C9—C14	116.0 (4)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N2—H2A \cdots C12	0.86	2.43	2.892 (4)	115
N2—H2A \cdots O1	0.86	1.98	2.664 (6)	135
N1—H1A \cdots O2 ⁱ	0.86	1.98	2.814 (6)	163
C10—H10A \cdots O2	0.93	2.28	2.851 (6)	119

Symmetry codes: (i) $-x+1, -y+2, -z+1$.

Fig. 1




Fig. 2

