

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(2,2'-bipyridine- κ^2N,N')(4-methylbenzoato- κ^2O,O')copper(II) iodide hemihydrate

Chuan-Mei Tang and Guo-Hua Deng*

College of Chemistry, South China University of Technology, Guangzhou 510640, People's Republic of China

Correspondence e-mail: ghdeng@126.com


Received 19 July 2008; accepted 30 July 2008

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.010$ Å; R factor = 0.047; wR factor = 0.149; data-to-parameter ratio = 14.5.

The title compound, $[Cu(C_8H_7O_2)(C_{10}H_8N_2)_2]I \cdot 0.5H_2O$, was obtained by the hydrothermal reaction of copper(I) iodide, 4-methylbenzoic acid and 2,2'-bipyridine. The initial reactant of Cu^I was oxidized to Cu^{II} . The asymmetric unit contains two independent complex molecules, two I^- ions and one water molecule. Each Cu^{II} atom is coordinated by two O atoms from a 4-methylbenzoate ligand and four N atoms from two 2,2'-bipyridine ligands, displaying a distorted octahedral geometry. The structure involves $O-H \cdots I$ hydrogen bonds between the water molecule and iodide ions and $\pi-\pi$ stacking interactions between the benzene and pyridyl rings [centroid-centroid distance = 3.79 (1) Å] and between the pyridyl rings [centroid-centroid distance = 3.87 (1) Å].

Related literature

For related literature, see: Ma & Deng (2008); Mao *et al.* (2001); Song *et al.* (2008*a,b,c,d*).



Experimental

Crystal data

$[Cu(C_8H_7O_2)(C_{10}H_8N_2)_2]I \cdot 0.5H_2O$
 $M_r = 646.96$
 Triclinic, $P\bar{1}$
 $a = 14.6698$ (4) Å
 $b = 15.3588$ (4) Å
 $c = 15.4224$ (7) Å
 $\alpha = 100.943$ (2)°
 $\beta = 114.345$ (2)°

$\gamma = 111.996$ (2)°
 $V = 2680.64$ (19) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 2.00$ mm⁻¹
 $T = 296$ (2) K
 $0.37 \times 0.30 \times 0.26$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{min} = 0.525$, $T_{max} = 0.624$

29909 measured reflections
 9564 independent reflections
 6740 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.149$
 $S = 1.03$
 9564 reflections
 660 parameters

3 restraints
 H-atom parameters constrained
 $\Delta\rho_{max} = 1.55$ e Å⁻³
 $\Delta\rho_{min} = -1.53$ e Å⁻³

Table 1

Selected bond lengths (Å).

Cu1—O1	1.976 (4)	Cu2—O3	1.974 (3)
Cu1—O2	2.769 (4)	Cu2—O4	2.832 (3)
Cu1—N6	1.987 (4)	Cu2—N8	1.997 (4)
Cu1—N3	2.000 (4)	Cu2—N2	2.001 (4)
Cu1—N5	2.060 (4)	Cu2—N7	2.038 (4)
Cu1—N4	2.192 (4)	Cu2—N1	2.181 (4)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O1W—H1W \cdots N1 ⁱ	0.82	3.15	3.935 (8)	161
O1W—H2W \cdots N1 ⁱⁱ	0.82	2.76	3.568 (8)	170

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x + 1, y + 1, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors acknowledge South China University of Technology for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2148).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc, Madison, Wisconsin, USA.
- Ma, D.-Y. & Deng, G.-H. (2008). *Acta Cryst.* **C64**, m271–m273.
- Mao, Z.-W., Heinemann, F. W., Liehr, G. & van Eldik, R. (2001). *J. Chem. Soc. Dalton Trans.* pp. 3652–3662.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Song, W.-D., Gu, C.-S., Hao, X.-M. & Yan, J.-B. (2008a). *Acta Cryst.* **E64**, m522.
- Song, W.-D., Wang, H. & Ji, L.-L. (2008b). *Acta Cryst.* **E64**, m548.
- Song, W.-D., Wang, H. & Miao, Y.-L. (2008c). *Acta Cryst.* **E64**, m716.
- Song, W.-D., Yan, J.-B. & Hao, X.-M. (2008d). *Acta Cryst.* **E64**, m919–m920.

supplementary materials

Acta Cryst. (2008). E64, m1111-m1112 [doi:10.1107/S1600536808024252]

Bis(2,2'-bipyridine- κ^2N,N')(4-methylbenzoato- κ^2O,O')copper(II) iodide hemihydrate

C.-M. Tang and G.-H. Deng

Comment

As a building block, 4-methylbenzoate ligand is an excellent candidate for the construction of supramolecular complexes (Ma & Deng, 2008; Song *et al.*, 2008a,b,c,d). Recently, we obtained the title mononuclear complex by the hydrothermal reaction of CuI, 4-methylbenzoic acid and 2,2'-bipyridine.

As illustrated in Fig. 1, the asymmetric unit of the title compound contains two independent complex molecules, two I ions and one lattice water molecule. Each Cu^{II} atom has a distorted octahedral coordination geometry, involving two carboxylate O atoms from one 4-methylbenzoate ligand, and four N atoms from two 2,2'-bipyridine ligands. One Cu—O distance is distinctly longer than the others for each Cu^{II} atom (Table 1), but still within the range of a significant interaction (Mao *et al.* 2001). Judged from the blue crystals, the initial reactant of Cu^I was thus oxidated to Cu^{II} in the hydrothermal reaction. The structure involves O—H...I hydrogen bonds between the water molecule and I ions (Table 2) and π – π stacking interactions (Fig. 2). The centroid–centroid distances are 3.79 (1) Å between the adjacent phenyl and pyridyl rings, and 3.87 (1) Å between the adjacent pyridyl rings.

Experimental

A mixture of CuI (0.1 g, 0.5 mmol), 4-methylbenzoic acid (0.068 g, 0.5 mmol), 2,2'-bipyridine (0.078 g, 0.5 mmol) and H₂O (10 ml) was placed in a 23 ml Teflon-lined reactor, which was heated to 433 K for 3 d and then cooled to room temperature at a rate of 10 K h⁻¹. Block colorless crystals were obtained.

Refinement

C-bound H atoms were positioned geometrically and refined as riding, with C—H = 0.93 (CH) and 0.96 (CH₃) Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH group or $1.5U_{\text{eq}}(\text{C})$ for CH₃ group. H atoms of water molecule were tentatively located in difference Fourier maps and refined with distance restraints of O—H = 0.84 (1) and H...H = 1.35 (1) Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. The highest residual electron density was 1.14 Å from atom I1 and the deepest hole 0.81 Å from atom I1.

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity.




Fig. 2. A packing view of the title compound. Hydrogen bonds are depicted as dashed lines.

Bis(2,2'-bipyridine- κ^2N,N')(4-methylbenzoato- κ^2O,O')copper(II) iodide hemihydrate

Crystal data

[Cu(C₈H₇O₂)(C₁₀H₈N₂)₂]*I*·0.5H₂O

M_r = 646.96

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

a = 14.6698 (4) Å

b = 15.3588 (4) Å

c = 15.4224 (7) Å

α = 100.943 (2)°

β = 114.345 (2)°

γ = 111.996 (2)°

V = 2680.64 (19) Å³

Z = 4

*F*₀₀₀ = 1288

D_x = 1.603 Mg m⁻³

Mo *K*α radiation

λ = 0.71073 Å

Cell parameters from 5300 reflections

θ = 1.3–28.0°

μ = 2.00 mm⁻¹

T = 296 (2) K

Block, blue

0.37 × 0.30 × 0.26 mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

T = 296(2) K

ϕ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

T_{min} = 0.525, *T_{max}* = 0.624

29909 measured reflections

9564 independent reflections

6740 reflections with *I* > 2σ(*I*)

R_{int} = 0.037

θ_{\max} = 25.2°

θ_{\min} = 1.6°

h = -17→17

k = -18→18

l = -17→18

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2σ(*F*²)] = 0.047

wR(*F*²) = 0.149

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0774P)^2 + 2.4876P]$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.03$ $(\Delta/\sigma)_{\max} < 0.001$
 9564 reflections $\Delta\rho_{\max} = 1.56 \text{ e } \text{\AA}^{-3}$
 660 parameters $\Delta\rho_{\min} = -1.53 \text{ e } \text{\AA}^{-3}$
 3 restraints Extinction correction: none
 Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5583 (5)	-0.0992 (4)	0.8879 (4)	0.0588 (14)
H1	0.6232	-0.0884	0.8844	0.071*
C2	0.5284 (6)	-0.1583 (5)	0.9396 (5)	0.0685 (16)
H2	0.5726	-0.1862	0.9707	0.082*
C3	0.4332 (6)	-0.1746 (5)	0.9441 (5)	0.0787 (19)
H3	0.4115	-0.2137	0.9787	0.094*
C4	0.3693 (5)	-0.1328 (5)	0.8973 (5)	0.0698 (17)
H4	0.3035	-0.1439	0.8992	0.084*
C5	0.4036 (4)	-0.0738 (4)	0.8470 (4)	0.0515 (12)
C6	0.3422 (4)	-0.0237 (4)	0.7958 (4)	0.0543 (13)
C7	0.2355 (6)	-0.0438 (5)	0.7799 (6)	0.083 (2)
H7	0.1976	-0.0916	0.7998	0.099*
C8	0.1861 (6)	0.0083 (7)	0.7338 (7)	0.100 (3)
H8	0.1139	-0.0047	0.7221	0.121*
C9	0.2416 (7)	0.0782 (7)	0.7056 (6)	0.091 (2)
H9	0.2095	0.1151	0.6767	0.110*
C10	0.3481 (5)	0.0941 (5)	0.7206 (5)	0.0719 (17)
H10	0.3862	0.1408	0.6997	0.086*
C11	0.3340 (4)	-0.0838 (4)	0.5409 (4)	0.0536 (13)
H11	0.2900	-0.0645	0.5605	0.064*
C12	0.2824 (5)	-0.1491 (4)	0.4398 (4)	0.0617 (15)
H12	0.2052	-0.1726	0.3913	0.074*
C13	0.3469 (5)	-0.1793 (4)	0.4111 (4)	0.0632 (15)
H13	0.3130	-0.2250	0.3433	0.076*
C14	0.4609 (5)	-0.1414 (4)	0.4830 (4)	0.0573 (13)
H14	0.5056	-0.1600	0.4640	0.069*
C15	0.5095 (4)	-0.0754 (3)	0.5840 (4)	0.0441 (11)
C16	0.6312 (4)	-0.0306 (3)	0.6684 (4)	0.0443 (11)
C17	0.7116 (5)	-0.0489 (4)	0.6564 (5)	0.0617 (14)
H17	0.6910	-0.0899	0.5912	0.074*
C18	0.8221 (5)	-0.0065 (5)	0.7407 (5)	0.0733 (17)
H18	0.8765	-0.0189	0.7334	0.088*
C19	0.8507 (5)	0.0547 (5)	0.8365 (5)	0.0703 (16)
H19	0.9242	0.0830	0.8950	0.084*
C20	0.7693 (4)	0.0734 (4)	0.8442 (4)	0.0595 (14)
H20	0.7894	0.1163	0.9083	0.071*
C21	0.6656 (4)	0.2547 (4)	0.8568 (4)	0.0494 (12)
C22	0.7626 (4)	0.3594 (4)	0.9370 (4)	0.0472 (12)

supplementary materials

C23	0.8244 (5)	0.3793 (5)	1.0407 (4)	0.0666 (16)
H23	0.8066	0.3261	1.0623	0.080*
C24	0.9126 (5)	0.4775 (5)	1.1130 (5)	0.0778 (18)
H24	0.9520	0.4891	1.1829	0.093*
C25	0.9441 (5)	0.5588 (4)	1.0854 (5)	0.0642 (16)
C26	0.8836 (5)	0.5382 (4)	0.9817 (5)	0.0664 (16)
H26	0.9037	0.5911	0.9600	0.080*
C27	0.7936 (5)	0.4407 (4)	0.9084 (4)	0.0577 (13)
H27	0.7531	0.4297	0.8387	0.069*
C28	1.0395 (6)	0.6647 (5)	1.1648 (6)	0.099 (2)
H28A	1.0074	0.7023	1.1859	0.148*
H28B	1.0801	0.6992	1.1353	0.148*
H28C	1.0924	0.6605	1.2243	0.148*
C36	1.0037 (6)	0.3445 (7)	0.4202 (7)	0.105 (3)
H36A	0.9998	0.3090	0.3592	0.158*
H36B	1.0105	0.3080	0.4644	0.158*
H36C	1.0701	0.4127	0.4573	0.158*
C30	0.8952 (5)	0.3506 (5)	0.3882 (5)	0.0709 (17)
C31	0.8731 (6)	0.4128 (7)	0.3403 (6)	0.098 (3)
H31	0.9283	0.4540	0.3288	0.118*
C32	0.7725 (5)	0.4176 (5)	0.3080 (5)	0.0744 (18)
H32	0.7607	0.4606	0.2751	0.089*
C33	0.6900 (4)	0.3575 (4)	0.3255 (4)	0.0432 (11)
C34	0.7122 (4)	0.2956 (4)	0.3758 (4)	0.0521 (12)
H34	0.6580	0.2551	0.3886	0.063*
C35	0.5819 (4)	0.3622 (4)	0.2935 (4)	0.0461 (11)
C37	0.6852 (4)	0.6225 (4)	0.4203 (5)	0.0648 (15)
H37	0.7121	0.6057	0.3783	0.078*
C38	0.7636 (5)	0.6975 (4)	0.5194 (5)	0.0746 (18)
H38	0.8424	0.7318	0.5438	0.090*
C39	0.7249 (5)	0.7213 (4)	0.5815 (5)	0.0669 (16)
H39	0.7775	0.7697	0.6501	0.080*
C40	0.6076 (5)	0.6734 (4)	0.5428 (4)	0.0562 (13)
H40	0.5798	0.6901	0.5840	0.067*
C41	0.5320 (4)	0.6000 (3)	0.4412 (4)	0.0441 (11)
C42	0.4036 (4)	0.5452 (3)	0.3899 (3)	0.0417 (11)
C43	0.3480 (5)	0.5727 (4)	0.4316 (4)	0.0569 (13)
H43	0.3898	0.6275	0.4960	0.068*
C44	0.2272 (5)	0.5173 (5)	0.3759 (5)	0.0681 (16)
H44	0.1867	0.5358	0.4015	0.082*
C45	0.1691 (5)	0.4352 (4)	0.2830 (5)	0.0627 (15)
H45	0.0886	0.3955	0.2457	0.075*
C46	0.2306 (4)	0.4124 (4)	0.2458 (4)	0.0506 (12)
H46	0.1902	0.3567	0.1824	0.061*
C47	0.2668 (5)	0.2356 (4)	0.1058 (5)	0.0598 (14)
H47	0.2849	0.2271	0.1678	0.072*
C48	0.1841 (5)	0.1508 (4)	0.0136 (6)	0.0715 (17)
H48	0.1474	0.0861	0.0126	0.086*
C49	0.1574 (5)	0.1646 (5)	-0.0771 (6)	0.082 (2)

H49	0.1013	0.1087	-0.1408	0.098*
C50	0.2138 (5)	0.2621 (5)	-0.0742 (4)	0.0706 (17)
H50	0.1966	0.2717	-0.1356	0.085*
C51	0.2964 (4)	0.3448 (4)	0.0223 (4)	0.0499 (12)
C52	0.3624 (4)	0.4512 (4)	0.0340 (4)	0.0481 (12)
C29	0.8130 (5)	0.2930 (4)	0.4069 (5)	0.0606 (14)
H29	0.8261	0.2514	0.4414	0.073*
C53	0.3479 (5)	0.4781 (5)	-0.0497 (4)	0.0609 (14)
H53	0.2921	0.4294	-0.1177	0.073*
C54	0.4170 (6)	0.5774 (5)	-0.0305 (5)	0.0700 (17)
H54	0.4096	0.5968	-0.0856	0.084*
C55	0.4974 (5)	0.6486 (5)	0.0704 (5)	0.0643 (15)
H55	0.5449	0.7165	0.0847	0.077*
C56	0.5062 (5)	0.6173 (4)	0.1499 (4)	0.0533 (13)
H56	0.5604	0.6655	0.2183	0.064*
Cu2	0.44994 (5)	0.45165 (4)	0.24535 (4)	0.04330 (16)
Cu1	0.53604 (5)	0.04754 (4)	0.76873 (5)	0.04622 (17)
I1	0.04839 (6)	0.20792 (5)	0.62512 (4)	0.1112 (2)
I2	0.98058 (3)	0.14574 (3)	0.13668 (3)	0.07583 (16)
N1	0.4398 (3)	0.5204 (3)	0.1327 (3)	0.0449 (9)
N2	0.3228 (3)	0.3298 (3)	0.1110 (3)	0.0483 (10)
N3	0.3964 (4)	0.0430 (3)	0.7649 (3)	0.0544 (11)
N4	0.4981 (3)	-0.0577 (3)	0.8433 (3)	0.0498 (10)
N5	0.4459 (3)	-0.0467 (3)	0.6130 (3)	0.0440 (9)
N6	0.6613 (3)	0.0313 (3)	0.7616 (3)	0.0479 (10)
N7	0.3464 (3)	0.4665 (3)	0.2964 (3)	0.0416 (9)
N8	0.5707 (3)	0.5723 (3)	0.3818 (3)	0.0478 (10)
O1	0.6478 (3)	0.1823 (3)	0.8889 (3)	0.0568 (9)
O2	0.6079 (3)	0.2405 (3)	0.7651 (3)	0.0674 (10)
O3	0.5675 (3)	0.4192 (3)	0.2437 (3)	0.0522 (8)
O4	0.5122 (3)	0.3137 (3)	0.3162 (3)	0.0597 (9)
O1W	0.8745 (9)	0.9386 (6)	0.5468 (7)	0.201 (4)
H2W	0.9082	1.0013	0.5663	0.301*
H1W	0.8900	0.9191	0.5042	0.301*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.063 (3)	0.062 (3)	0.066 (4)	0.034 (3)	0.041 (3)	0.030 (3)
C2	0.075 (4)	0.066 (4)	0.066 (4)	0.034 (3)	0.039 (3)	0.031 (3)
C3	0.093 (5)	0.070 (4)	0.078 (4)	0.029 (4)	0.062 (4)	0.028 (3)
C4	0.069 (4)	0.064 (4)	0.067 (4)	0.018 (3)	0.050 (3)	0.011 (3)
C5	0.049 (3)	0.045 (3)	0.048 (3)	0.015 (2)	0.031 (3)	0.004 (2)
C6	0.042 (3)	0.056 (3)	0.049 (3)	0.017 (2)	0.027 (3)	0.001 (2)
C7	0.058 (4)	0.090 (5)	0.097 (5)	0.035 (4)	0.049 (4)	0.020 (4)
C8	0.060 (4)	0.124 (7)	0.101 (6)	0.050 (5)	0.041 (4)	0.011 (5)
C9	0.079 (5)	0.106 (6)	0.091 (5)	0.071 (5)	0.034 (4)	0.023 (4)
C10	0.067 (4)	0.069 (4)	0.076 (4)	0.046 (3)	0.032 (3)	0.013 (3)

supplementary materials

C11	0.049 (3)	0.054 (3)	0.048 (3)	0.023 (2)	0.022 (3)	0.018 (2)
C12	0.051 (3)	0.065 (4)	0.045 (3)	0.021 (3)	0.016 (3)	0.017 (3)
C13	0.068 (4)	0.068 (4)	0.043 (3)	0.028 (3)	0.029 (3)	0.015 (3)
C14	0.065 (4)	0.055 (3)	0.056 (3)	0.030 (3)	0.037 (3)	0.017 (3)
C15	0.047 (3)	0.040 (2)	0.050 (3)	0.021 (2)	0.029 (2)	0.022 (2)
C16	0.048 (3)	0.043 (3)	0.053 (3)	0.027 (2)	0.032 (2)	0.022 (2)
C17	0.059 (3)	0.062 (3)	0.070 (4)	0.034 (3)	0.038 (3)	0.019 (3)
C18	0.059 (4)	0.083 (4)	0.087 (5)	0.043 (3)	0.042 (4)	0.029 (4)
C19	0.042 (3)	0.085 (4)	0.070 (4)	0.030 (3)	0.025 (3)	0.021 (3)
C20	0.041 (3)	0.065 (3)	0.055 (3)	0.022 (3)	0.022 (3)	0.015 (3)
C21	0.044 (3)	0.054 (3)	0.048 (3)	0.022 (2)	0.031 (3)	0.009 (2)
C22	0.039 (3)	0.043 (3)	0.049 (3)	0.016 (2)	0.023 (2)	0.008 (2)
C23	0.060 (4)	0.069 (4)	0.052 (3)	0.019 (3)	0.027 (3)	0.023 (3)
C24	0.056 (4)	0.084 (5)	0.047 (3)	0.017 (3)	0.014 (3)	0.010 (3)
C25	0.041 (3)	0.056 (3)	0.067 (4)	0.015 (3)	0.026 (3)	0.000 (3)
C26	0.054 (3)	0.049 (3)	0.083 (5)	0.021 (3)	0.034 (3)	0.019 (3)
C27	0.055 (3)	0.054 (3)	0.056 (3)	0.027 (3)	0.026 (3)	0.016 (3)
C28	0.057 (4)	0.075 (4)	0.098 (5)	0.013 (3)	0.026 (4)	-0.008 (4)
C36	0.064 (4)	0.188 (8)	0.124 (6)	0.083 (5)	0.063 (5)	0.107 (7)
C30	0.055 (3)	0.109 (5)	0.072 (4)	0.048 (3)	0.039 (3)	0.053 (4)
C31	0.059 (4)	0.168 (7)	0.121 (6)	0.064 (5)	0.062 (4)	0.109 (6)
C32	0.057 (4)	0.097 (5)	0.088 (5)	0.041 (3)	0.041 (3)	0.061 (4)
C33	0.038 (3)	0.052 (3)	0.039 (3)	0.022 (2)	0.021 (2)	0.016 (2)
C34	0.050 (3)	0.056 (3)	0.060 (3)	0.027 (2)	0.037 (3)	0.025 (3)
C35	0.042 (3)	0.055 (3)	0.031 (2)	0.025 (2)	0.015 (2)	0.006 (2)
C37	0.041 (3)	0.061 (3)	0.071 (4)	0.022 (3)	0.025 (3)	0.007 (3)
C38	0.036 (3)	0.057 (3)	0.088 (5)	0.020 (3)	0.015 (3)	0.000 (3)
C39	0.051 (3)	0.053 (3)	0.058 (3)	0.023 (3)	0.009 (3)	0.003 (3)
C40	0.053 (3)	0.055 (3)	0.044 (3)	0.024 (3)	0.019 (3)	0.010 (2)
C41	0.042 (3)	0.039 (2)	0.039 (3)	0.017 (2)	0.015 (2)	0.015 (2)
C42	0.042 (3)	0.039 (2)	0.039 (3)	0.018 (2)	0.020 (2)	0.017 (2)
C43	0.056 (3)	0.053 (3)	0.054 (3)	0.021 (3)	0.033 (3)	0.014 (2)
C44	0.070 (4)	0.066 (4)	0.080 (4)	0.032 (3)	0.054 (4)	0.023 (3)
C45	0.048 (3)	0.060 (3)	0.075 (4)	0.021 (3)	0.038 (3)	0.018 (3)
C46	0.042 (3)	0.045 (3)	0.053 (3)	0.015 (2)	0.023 (2)	0.014 (2)
C47	0.051 (3)	0.051 (3)	0.067 (4)	0.026 (3)	0.029 (3)	0.013 (3)
C48	0.051 (3)	0.050 (3)	0.084 (5)	0.015 (3)	0.032 (3)	0.005 (3)
C49	0.044 (3)	0.067 (4)	0.077 (5)	0.011 (3)	0.020 (3)	-0.015 (3)
C50	0.049 (3)	0.087 (5)	0.045 (3)	0.026 (3)	0.019 (3)	0.003 (3)
C51	0.040 (3)	0.061 (3)	0.041 (3)	0.027 (2)	0.020 (2)	0.007 (2)
C52	0.044 (3)	0.064 (3)	0.043 (3)	0.035 (2)	0.025 (2)	0.016 (2)
C29	0.062 (3)	0.076 (4)	0.070 (4)	0.045 (3)	0.041 (3)	0.043 (3)
C53	0.067 (4)	0.084 (4)	0.044 (3)	0.049 (3)	0.029 (3)	0.026 (3)
C54	0.092 (5)	0.094 (5)	0.062 (4)	0.066 (4)	0.048 (4)	0.046 (4)
C55	0.081 (4)	0.067 (4)	0.075 (4)	0.046 (3)	0.052 (4)	0.041 (3)
C56	0.062 (3)	0.053 (3)	0.053 (3)	0.034 (3)	0.032 (3)	0.025 (3)
Cu2	0.0401 (3)	0.0434 (3)	0.0389 (3)	0.0195 (3)	0.0192 (3)	0.0103 (2)
Cu1	0.0409 (3)	0.0460 (3)	0.0491 (4)	0.0218 (3)	0.0244 (3)	0.0136 (3)
II	0.1320 (5)	0.1664 (6)	0.0966 (4)	0.1121 (5)	0.0670 (4)	0.0762 (4)

I2	0.0619 (3)	0.0608 (3)	0.0672 (3)	0.01033 (19)	0.0296 (2)	0.0105 (2)
N1	0.045 (2)	0.053 (2)	0.040 (2)	0.029 (2)	0.0223 (19)	0.0163 (18)
N2	0.042 (2)	0.043 (2)	0.046 (2)	0.0190 (18)	0.0195 (19)	0.0059 (18)
N3	0.046 (2)	0.055 (3)	0.054 (3)	0.028 (2)	0.024 (2)	0.009 (2)
N4	0.046 (2)	0.051 (2)	0.051 (2)	0.021 (2)	0.031 (2)	0.015 (2)
N5	0.044 (2)	0.039 (2)	0.045 (2)	0.0182 (18)	0.023 (2)	0.0153 (17)
N6	0.042 (2)	0.046 (2)	0.053 (3)	0.0201 (19)	0.027 (2)	0.017 (2)
N7	0.040 (2)	0.043 (2)	0.039 (2)	0.0192 (18)	0.0207 (18)	0.0167 (17)
N8	0.040 (2)	0.049 (2)	0.049 (2)	0.0215 (19)	0.023 (2)	0.0126 (19)
O1	0.058 (2)	0.046 (2)	0.059 (2)	0.0213 (17)	0.0337 (19)	0.0136 (17)
O2	0.068 (2)	0.065 (2)	0.048 (2)	0.022 (2)	0.029 (2)	0.0106 (18)
O3	0.053 (2)	0.060 (2)	0.051 (2)	0.0343 (17)	0.0282 (17)	0.0239 (17)
O4	0.048 (2)	0.084 (3)	0.058 (2)	0.0359 (19)	0.0337 (19)	0.031 (2)
O1W	0.325 (12)	0.176 (7)	0.233 (9)	0.142 (8)	0.225 (10)	0.116 (7)

Geometric parameters (Å, °)

C1—N4	1.322 (6)	C31—H31	0.9300
C1—C2	1.384 (8)	C32—C33	1.385 (7)
C1—H1	0.9300	C32—H32	0.9300
C2—C3	1.357 (9)	C33—C34	1.384 (7)
C2—H2	0.9300	C33—C35	1.489 (6)
C3—C4	1.372 (9)	C34—C29	1.372 (7)
C3—H3	0.9300	C34—H34	0.9300
C4—C5	1.387 (8)	C35—O4	1.243 (6)
C4—H4	0.9300	C35—O3	1.289 (6)
C5—N4	1.341 (6)	C37—N8	1.342 (6)
C5—C6	1.480 (8)	C37—C38	1.370 (8)
C6—N3	1.340 (7)	C37—H37	0.9300
C6—C7	1.377 (8)	C38—C39	1.358 (9)
C7—C8	1.377 (11)	C38—H38	0.9300
C7—H7	0.9300	C39—C40	1.377 (8)
C8—C9	1.348 (11)	C39—H39	0.9300
C8—H8	0.9300	C40—C41	1.384 (7)
C9—C10	1.395 (9)	C40—H40	0.9300
C9—H9	0.9300	C41—N8	1.344 (6)
C10—N3	1.344 (7)	C41—C42	1.487 (6)
C10—H10	0.9300	C42—N7	1.351 (6)
C11—N5	1.346 (6)	C42—C43	1.358 (7)
C11—C12	1.369 (7)	C43—C44	1.393 (8)
C11—H11	0.9300	C43—H43	0.9300
C12—C13	1.380 (8)	C44—C45	1.368 (8)
C12—H12	0.9300	C44—H44	0.9300
C13—C14	1.364 (8)	C45—C46	1.360 (7)
C13—H13	0.9300	C45—H45	0.9300
C14—C15	1.380 (7)	C46—N7	1.338 (6)
C14—H14	0.9300	C46—H46	0.9300
C15—N5	1.357 (6)	C47—N2	1.336 (7)
C15—C16	1.476 (7)	C47—C48	1.371 (8)

supplementary materials

C16—N6	1.345 (6)	C47—H47	0.9300
C16—C17	1.384 (7)	C48—C49	1.370 (10)
C17—C18	1.375 (8)	C48—H48	0.9300
C17—H17	0.9300	C49—C50	1.394 (9)
C18—C19	1.382 (9)	C49—H49	0.9300
C18—H18	0.9300	C50—C51	1.393 (7)
C19—C20	1.372 (8)	C50—H50	0.9300
C19—H19	0.9300	C51—N2	1.349 (6)
C20—N6	1.344 (6)	C51—C52	1.484 (7)
C20—H20	0.9300	C52—N1	1.339 (6)
C21—O2	1.233 (6)	C52—C53	1.387 (7)
C21—O1	1.284 (6)	C29—H29	0.9300
C21—C22	1.490 (7)	C53—C54	1.366 (8)
C22—C23	1.373 (7)	C53—H53	0.9300
C22—C27	1.381 (7)	C54—C55	1.372 (8)
C23—C24	1.380 (8)	C54—H54	0.9300
C23—H23	0.9300	C55—C56	1.375 (7)
C24—C25	1.373 (9)	C55—H55	0.9300
C24—H24	0.9300	C56—N1	1.336 (6)
C25—C26	1.369 (8)	C56—H56	0.9300
C25—C28	1.492 (8)	Cu1—O1	1.976 (4)
C26—C27	1.381 (8)	Cu1—O2	2.769 (4)
C26—H26	0.9300	Cu1—N6	1.987 (4)
C27—H27	0.9300	Cu1—N3	2.000 (4)
C28—H28A	0.9600	Cu1—N5	2.060 (4)
C28—H28B	0.9600	Cu1—N4	2.192 (4)
C28—H28C	0.9600	Cu2—O3	1.974 (3)
C36—C30	1.506 (8)	Cu2—O4	2.832 (3)
C36—H36A	0.9600	Cu2—N8	1.997 (4)
C36—H36B	0.9600	Cu2—N2	2.001 (4)
C36—H36C	0.9600	Cu2—N7	2.038 (4)
C30—C31	1.367 (8)	Cu2—N1	2.181 (4)
C30—C29	1.379 (8)	O1W—H2W	0.8200
C31—C32	1.385 (8)	O1W—H1W	0.8200
N4—C1—C2	122.5 (5)	C37—C38—H38	120.4
N4—C1—H1	118.7	C38—C39—C40	119.8 (5)
C2—C1—H1	118.7	C38—C39—H39	120.1
C3—C2—C1	118.7 (6)	C40—C39—H39	120.1
C3—C2—H2	120.6	C39—C40—C41	118.5 (5)
C1—C2—H2	120.6	C39—C40—H40	120.7
C2—C3—C4	119.3 (6)	C41—C40—H40	120.7
C2—C3—H3	120.3	N8—C41—C40	121.6 (5)
C4—C3—H3	120.3	N8—C41—C42	115.0 (4)
C3—C4—C5	119.5 (6)	C40—C41—C42	123.5 (5)
C3—C4—H4	120.2	N7—C42—C43	122.7 (4)
C5—C4—H4	120.2	N7—C42—C41	113.9 (4)
N4—C5—C4	120.7 (5)	C43—C42—C41	123.4 (4)
N4—C5—C6	115.5 (4)	C42—C43—C44	118.7 (5)
C4—C5—C6	123.8 (5)	C42—C43—H43	120.7

N3—C6—C7	121.3 (6)	C44—C43—H43	120.7
N3—C6—C5	116.0 (4)	C45—C44—C43	118.8 (5)
C7—C6—C5	122.8 (6)	C45—C44—H44	120.6
C6—C7—C8	118.7 (7)	C43—C44—H44	120.6
C6—C7—H7	120.7	C46—C45—C44	119.2 (5)
C8—C7—H7	120.7	C46—C45—H45	120.4
C9—C8—C7	120.6 (7)	C44—C45—H45	120.4
C9—C8—H8	119.7	N7—C46—C45	123.0 (5)
C7—C8—H8	119.7	N7—C46—H46	118.5
C8—C9—C10	118.8 (7)	C45—C46—H46	118.5
C8—C9—H9	120.6	N2—C47—C48	123.2 (6)
C10—C9—H9	120.6	N2—C47—H47	118.4
N3—C10—C9	120.8 (7)	C48—C47—H47	118.4
N3—C10—H10	119.6	C49—C48—C47	117.8 (6)
C9—C10—H10	119.6	C49—C48—H48	121.1
N5—C11—C12	122.4 (5)	C47—C48—H48	121.1
N5—C11—H11	118.8	C48—C49—C50	120.3 (6)
C12—C11—H11	118.8	C48—C49—H49	119.8
C11—C12—C13	118.8 (5)	C50—C49—H49	119.8
C11—C12—H12	120.6	C51—C50—C49	118.7 (6)
C13—C12—H12	120.6	C51—C50—H50	120.6
C14—C13—C12	119.5 (5)	C49—C50—H50	120.6
C14—C13—H13	120.3	N2—C51—C50	120.2 (5)
C12—C13—H13	120.3	N2—C51—C52	116.7 (4)
C13—C14—C15	119.7 (5)	C50—C51—C52	123.0 (5)
C13—C14—H14	120.1	N1—C52—C53	121.9 (5)
C15—C14—H14	120.1	N1—C52—C51	114.6 (4)
N5—C15—C14	121.0 (5)	C53—C52—C51	123.5 (5)
N5—C15—C16	114.4 (4)	C34—C29—C30	121.6 (5)
C14—C15—C16	124.6 (5)	C34—C29—H29	119.2
N6—C16—C17	120.4 (5)	C30—C29—H29	119.2
N6—C16—C15	115.3 (4)	C54—C53—C52	118.8 (5)
C17—C16—C15	124.3 (5)	C54—C53—H53	120.6
C18—C17—C16	120.1 (6)	C52—C53—H53	120.6
C18—C17—H17	119.9	C53—C54—C55	119.7 (5)
C16—C17—H17	119.9	C53—C54—H54	120.1
C17—C18—C19	118.7 (6)	C55—C54—H54	120.1
C17—C18—H18	120.7	C54—C55—C56	118.6 (6)
C19—C18—H18	120.7	C54—C55—H55	120.7
C20—C19—C18	119.3 (6)	C56—C55—H55	120.7
C20—C19—H19	120.4	N1—C56—C55	122.6 (5)
C18—C19—H19	120.4	N1—C56—H56	118.7
N6—C20—C19	121.7 (5)	C55—C56—H56	118.7
N6—C20—H20	119.1	O1—Cu1—O2	52.46 (16)
C19—C20—H20	119.1	O1—Cu1—N6	92.58 (16)
O2—C21—O1	123.2 (5)	O1—Cu1—N3	94.78 (16)
O2—C21—C22	120.5 (5)	O2—Cu1—N6	96.13 (16)
O1—C21—C22	116.3 (5)	O2—Cu1—N3	90.52 (16)
C23—C22—C27	117.5 (5)	N6—Cu1—N3	172.15 (16)

supplementary materials

C23—C22—C21	122.2 (5)	O1—Cu1—N5	152.67 (15)
C27—C22—C21	120.3 (5)	O2—Cu1—N5	101.75 (16)
C22—C23—C24	120.7 (6)	N6—Cu1—N5	80.38 (16)
C22—C23—H23	119.7	N3—Cu1—N5	94.23 (16)
C24—C23—H23	119.7	O1—Cu1—N4	102.51 (15)
C25—C24—C23	122.3 (6)	O2—Cu1—N4	152.05 (16)
C25—C24—H24	118.9	N6—Cu1—N4	97.40 (16)
C23—C24—H24	118.9	N3—Cu1—N4	78.34 (17)
C26—C25—C24	116.8 (5)	N5—Cu1—N4	104.56 (15)
C26—C25—C28	121.7 (6)	O3—Cu2—O4	51.67 (15)
C24—C25—C28	121.6 (6)	O3—Cu2—N8	90.84 (15)
C25—C26—C27	121.7 (6)	O3—Cu2—N2	91.78 (15)
C25—C26—H26	119.1	O4—Cu2—N8	93.70 (15)
C27—C26—H26	119.1	O4—Cu2—N2	86.63 (15)
C26—C27—C22	121.0 (5)	N8—Cu2—N2	176.92 (16)
C26—C27—H27	119.5	O3—Cu2—N7	156.24 (15)
C22—C27—H27	119.5	O4—Cu2—N7	106.46 (15)
C25—C28—H28A	109.5	N8—Cu2—N7	80.59 (16)
C25—C28—H28B	109.5	N2—Cu2—N7	96.37 (16)
H28A—C28—H28B	109.5	O3—Cu2—N1	94.97 (14)
C25—C28—H28C	109.5	O4—Cu2—N1	143.21 (15)
H28A—C28—H28C	109.5	N8—Cu2—N1	102.89 (16)
H28B—C28—H28C	109.5	N2—Cu2—N1	78.49 (16)
C30—C36—H36A	109.5	N7—Cu2—N1	108.50 (14)
C30—C36—H36B	109.5	C56—N1—C52	118.4 (4)
H36A—C36—H36B	109.5	C56—N1—Cu2	128.6 (3)
C30—C36—H36C	109.5	C52—N1—Cu2	112.8 (3)
H36A—C36—H36C	109.5	C47—N2—C51	119.7 (4)
H36B—C36—H36C	109.5	C47—N2—Cu2	123.0 (4)
C31—C30—C29	116.7 (5)	C51—N2—Cu2	117.3 (3)
C31—C30—C36	122.4 (6)	C6—N3—C10	119.8 (5)
C29—C30—C36	120.9 (6)	C6—N3—Cu1	117.7 (3)
C30—C31—C32	123.3 (6)	C10—N3—Cu1	122.0 (4)
C30—C31—H31	118.4	C1—N4—C5	119.2 (5)
C32—C31—H31	118.4	C1—N4—Cu1	129.0 (3)
C33—C32—C31	119.1 (5)	C5—N4—Cu1	111.7 (3)
C33—C32—H32	120.5	C11—N5—C15	118.6 (4)
C31—C32—H32	120.5	C11—N5—Cu1	127.7 (3)
C34—C33—C32	118.3 (5)	C15—N5—Cu1	113.7 (3)
C34—C33—C35	121.2 (4)	C20—N6—C16	119.8 (4)
C32—C33—C35	120.5 (5)	C20—N6—Cu1	124.0 (4)
C29—C34—C33	121.1 (5)	C16—N6—Cu1	116.1 (3)
C29—C34—H34	119.5	C46—N7—C42	117.5 (4)
C33—C34—H34	119.5	C46—N7—Cu2	127.8 (3)
O4—C35—O3	123.7 (4)	C42—N7—Cu2	114.5 (3)
O4—C35—C33	120.0 (4)	C37—N8—C41	118.6 (4)
O3—C35—C33	116.3 (4)	C37—N8—Cu2	125.8 (4)
N8—C37—C38	122.2 (6)	C41—N8—Cu2	115.5 (3)
N8—C37—H37	118.9	C21—O1—Cu1	110.0 (3)

C38—C37—H37	118.9	C35—O3—Cu2	111.0 (3)
C39—C38—C37	119.2 (5)	H2W—O1W—H1W	103.5
C39—C38—H38	120.4		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H1W \cdots I1 ⁱ	0.82	3.15	3.935 (8)	161
O1W—H2W \cdots I1 ⁱⁱ	0.82	2.76	3.568 (8)	170

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x+1, y+1, z$.

Fig. 1




Fig. 2

