

3,5-Dinitropyridin-4(1*H*)-one monohydrate

Ying Li,^a Ping Li,^a Qiu-Ping Zhou,^b Guo-Fang Zhang^b and Seik Weng Ng^{c*}

^aChemistry Department, Jingning Normal College, Wulanchabu, Inner Mongolia 012000, People's Republic of China, ^bSchool of Chemistry and Materials Science, Shaanxi Normal University, Xi'an 710062, People's Republic of China, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 25 July 2008; accepted 31 July 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.041; wR factor = 0.124; data-to-parameter ratio = 12.7.

The three independent organic molecules of 3,5-dinitropyridin-4(1*H*)-one monohydrate, $\text{C}_5\text{H}_3\text{N}_3\text{O}_5 \cdot \text{H}_2\text{O}$, each feature an $\text{N}-\text{H} \cdots \text{O}_{\text{water}}$ hydrogen bond. Each water molecule serves as hydrogen-bond donor to two carbonyl O atoms; these hydrogen bonds give rise to a layer motif. Two of the three formula units lie on special positions of site symmetry 2.

Related literature

The parent pyridin-4-one homolog crystallizes with five pyridone and six water molecules in the asymmetric unit; see: Jones (2001).



Experimental

Crystal data

$\text{C}_5\text{H}_3\text{N}_3\text{O}_5 \cdot \text{H}_2\text{O}$
 $M_r = 203.12$
Orthorhombic, $Pbcn$
 $a = 21.728$ (2) Å
 $b = 21.654$ (2) Å
 $c = 6.5713$ (5) Å

$V = 3091.7$ (4) Å³
 $Z = 16$
Mo $K\alpha$ radiation
 $\mu = 0.16$ mm⁻¹
 $T = 293$ (2) K
0.45 × 0.45 × 0.20 mm

Data collection

Bruker APEXII diffractometer
Absorption correction: none
21800 measured reflections

3555 independent reflections
2852 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.123$
 $S = 1.08$
3555 reflections
281 parameters
10 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N3}-\text{H3} \cdots \text{O1w}$	0.85 (1)	1.86 (1)	2.703 (2)	172 (2)
$\text{N5}-\text{H5} \cdots \text{O2w}$	0.85 (1)	1.84 (1)	2.692 (3)	180
$\text{N6}-\text{H6} \cdots \text{O3w}$	0.85 (1)	1.87 (1)	2.723 (3)	180
$\text{O1w}-\text{H11} \cdots \text{O8}^i$	0.85 (1)	2.04 (1)	2.878 (2)	168 (2)
$\text{O1w}-\text{H12} \cdots \text{O11}^{\text{ii}}$	0.86 (1)	2.02 (1)	2.866 (2)	168 (2)
$\text{O2w}-\text{H21} \cdots \text{O3}^{\text{iii}}$	0.84 (1)	2.05 (1)	2.888 (2)	173 (1)
$\text{O3w}-\text{H31} \cdots \text{O3}$	0.84 (1)	2.05 (1)	2.890 (2)	172 (2)

Symmetry codes: (i) $-x + \frac{3}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$; (ii) $x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2008).

We thank Jingning Normal College, Shaanxi Normal University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2110).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Jones, P. G. (2001). *Acta Cryst.* **C57**, 880–882.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2008). *pubCIF*. In preparation.

supplementary materials

Acta Cryst. (2008). E64, o1701 [doi:10.1107/S1600536808024604]

3,5-Dinitropyridin-4(1*H*)-one monohydrate

Y. Li, P. Li, Q.-P. Zhou, G.-F. Zhang and S. W. Ng

Comment

3,5-Dinitro-4-pyridinol, a specialty chemical, is assumed in chemical catalogs to exist in the enol form. The homolog, 4-pyridinol, is in fact 4-pyridinone·6/5hydrate. It has five independent pyridone and six water molecules that are hydrogen bonded to form layers (Jones, 2001). The presence of two electron-withdrawing groups in the title compound should enhance its propensity to form hydrogen bonds, and this is borne out in the present study. The three independent molecules of 3,5-dinitro-1*H*-pyridin-4-one hydrate (Fig. 1) each feature an N–H···O_{water} hydrogen bond; each water molecule serves as hydrogen-bond donor to two carbonyl oxygen atoms, and these hydrogen bonds give rise to a layered structure.

Experimental

4-Hydroxy-3-pyridine (19 g, 0.02 mol) was dissolved in fuming sulfuric acid (50% by SO₃ content) (60 ml), and to the solution was added an oleum-fuming nitric acid (1/3) mixture (50 ml). The temperature was kept at 0°C for an hour. The temperature was raised to 413 K over a period of one hour, and then held at 403 K for another 16 h. The mixture was poured into ice (200 g) to quench the reaction. Some 27 g of material was isolated. Crystals were obtained by recrystallization from water.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93 Å) and were included in the refinement using a riding model approximation, with $U(\text{H}) 1.2U_{\text{eq}}(\text{C})$. The amino and water H-atoms were refined with distance restraints of O–H = N–H 0.85 (1) and H···H 1.39 (1) Å; their temperature factors U_{iso} were freely refined.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) plot of the three independent molecules of C₅H₃N₃O₅·H₂O at the 50% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

3,5-Dinitropyridin-4(1*H*)-one monohydrate

Crystal data

C₅H₃N₃O₅·H₂O

$F_{000} = 1664$

supplementary materials

$M_r = 203.12$

Orthorhombic, *Pbcn*

Hall symbol: -P 2n 2ab

$a = 21.728$ (2) Å

$b = 21.654$ (2) Å

$c = 6.5713$ (5) Å

$V = 3091.7$ (4) Å³

$Z = 16$

$D_x = 1.746$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 6494 reflections

$\theta = 2.7$ – 27.9°

$\mu = 0.16$ mm⁻¹

$T = 293$ (2) K

Block, colorless

$0.45 \times 0.45 \times 0.20$ mm

Data collection

Bruker APEXII
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 9 pixels mm⁻¹

$T = 293$ (2) K

φ and ω scans

Absorption correction: none

21800 measured reflections

3555 independent reflections

2852 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.022$

$\theta_{\text{max}} = 27.5^\circ$

$\theta_{\text{min}} = 2.7^\circ$

$h = -28 \rightarrow 27$

$k = -28 \rightarrow 27$

$l = -8 \rightarrow 8$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.123$

$S = 1.08$

3555 reflections

281 parameters

10 restraints

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H atoms treated by a mixture of
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0528P)^2 + 1.6904P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.35$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.31$ e Å⁻³

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.69961 (7)	0.46193 (6)	0.2892 (3)	0.0552 (4)
O2	0.62136 (6)	0.50982 (6)	0.4163 (3)	0.0464 (4)
O3	0.60319 (6)	0.62125 (6)	0.2292 (2)	0.0431 (4)
O4	0.61633 (6)	0.73545 (6)	0.0695 (3)	0.0486 (4)
O5	0.70310 (8)	0.77792 (8)	0.1081 (6)	0.1176 (12)
O6	0.60858 (6)	0.75858 (6)	0.5467 (3)	0.0496 (4)
O7	0.65774 (6)	0.67898 (7)	0.6579 (3)	0.0562 (4)

O8	0.5000	0.77618 (7)	0.7500	0.0389 (4)
O9	0.65904 (7)	0.92189 (7)	0.1991 (3)	0.0659 (5)
O10	0.61401 (6)	1.00203 (6)	0.0759 (3)	0.0482 (4)
O11	0.5000	1.02039 (7)	0.2500	0.0357 (4)
O1W	0.91543 (6)	0.62174 (6)	0.2293 (3)	0.0507 (4)
O2W	0.5000	0.46255 (9)	0.7500	0.0627 (7)
O3W	0.5000	0.70522 (8)	0.2500	0.0416 (5)
N1	0.67114 (7)	0.50953 (7)	0.3278 (3)	0.0360 (3)
N2	0.67066 (7)	0.73271 (7)	0.1089 (3)	0.0435 (4)
N3	0.79137 (7)	0.61956 (8)	0.1992 (3)	0.0434 (4)
N4	0.61008 (6)	0.70807 (7)	0.6311 (3)	0.0357 (3)
N5	0.5000	0.58686 (9)	0.7500	0.0376 (5)
N6	0.5000	0.83100 (9)	0.2500	0.0384 (5)
N7	0.61264 (7)	0.95139 (7)	0.1582 (3)	0.0377 (4)
C1	0.76100 (8)	0.56854 (9)	0.2549 (3)	0.0390 (4)
H1A	0.7829	0.5329	0.2857	0.047*
C2	0.69872 (8)	0.56805 (8)	0.2672 (3)	0.0315 (4)
C3	0.65995 (8)	0.62114 (7)	0.2208 (3)	0.0302 (4)
C4	0.69804 (8)	0.67347 (8)	0.1610 (3)	0.0337 (4)
C5	0.76066 (8)	0.67135 (9)	0.1539 (3)	0.0405 (4)
H5A	0.7825	0.7065	0.1169	0.049*
C6	0.55166 (8)	0.61756 (8)	0.7027 (3)	0.0345 (4)
H6A	0.5874	0.5957	0.6733	0.041*
C7	0.55244 (7)	0.68038 (7)	0.6972 (3)	0.0290 (4)
C8	0.5000	0.71922 (10)	0.7500	0.0269 (5)
C9	0.55223 (9)	0.86172 (8)	0.2128 (3)	0.0360 (4)
H9	0.5884	0.8398	0.1893	0.043*
C10	0.55324 (8)	0.92451 (7)	0.2090 (3)	0.0301 (4)
C11	0.5000	0.96327 (10)	0.2500	0.0278 (5)
H11	0.9376 (8)	0.6540 (6)	0.220 (4)	0.059 (7)*
H12	0.9385 (8)	0.5898 (6)	0.218 (4)	0.053 (7)*
H21	0.4680 (2)	0.4409 (7)	0.757 (4)	0.055 (7)*
H31	0.5318 (3)	0.6832 (7)	0.236 (4)	0.058 (7)*
H3	0.8306 (5)	0.6181 (10)	0.199 (4)	0.057 (7)*
H5	0.5000	0.5477 (5)	0.7500	0.051 (9)*
H6	0.5000	0.7915 (5)	0.2500	0.051 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0543 (9)	0.0322 (7)	0.0790 (12)	0.0112 (6)	0.0045 (8)	0.0018 (7)
O2	0.0369 (7)	0.0417 (7)	0.0607 (9)	-0.0020 (6)	0.0068 (7)	0.0077 (7)
O3	0.0218 (6)	0.0344 (7)	0.0730 (10)	0.0014 (5)	0.0029 (6)	0.0061 (6)
O4	0.0381 (7)	0.0401 (7)	0.0676 (10)	0.0086 (6)	-0.0039 (7)	0.0016 (7)
O5	0.0504 (10)	0.0435 (10)	0.259 (4)	-0.0121 (8)	-0.0082 (15)	0.0441 (15)
O6	0.0415 (7)	0.0392 (7)	0.0681 (10)	-0.0090 (6)	0.0128 (7)	0.0059 (7)
O7	0.0273 (7)	0.0575 (9)	0.0837 (12)	0.0076 (6)	0.0053 (7)	-0.0044 (8)
O8	0.0326 (9)	0.0211 (8)	0.0631 (13)	0.000	0.0069 (8)	0.000

supplementary materials

O9	0.0304 (7)	0.0501 (9)	0.1171 (16)	0.0104 (6)	0.0009 (8)	0.0031 (9)
O10	0.0399 (7)	0.0372 (7)	0.0673 (10)	-0.0056 (6)	0.0073 (7)	0.0088 (7)
O11	0.0305 (9)	0.0199 (8)	0.0567 (12)	0.000	0.0020 (8)	0.000
O1W	0.0263 (7)	0.0294 (7)	0.0962 (13)	0.0008 (5)	-0.0054 (7)	0.0013 (7)
O2W	0.0296 (10)	0.0253 (9)	0.133 (2)	0.000	-0.0014 (12)	0.000
O3W	0.0286 (9)	0.0244 (8)	0.0719 (14)	0.000	0.0004 (9)	0.000
N1	0.0345 (8)	0.0322 (8)	0.0414 (9)	0.0028 (6)	-0.0042 (7)	0.0014 (6)
N2	0.0356 (8)	0.0335 (8)	0.0614 (11)	-0.0008 (6)	0.0060 (8)	0.0067 (7)
N3	0.0206 (7)	0.0488 (10)	0.0608 (11)	0.0008 (6)	-0.0006 (7)	0.0012 (8)
N4	0.0275 (7)	0.0353 (8)	0.0442 (9)	-0.0025 (6)	0.0039 (6)	-0.0084 (7)
N5	0.0434 (12)	0.0192 (9)	0.0503 (14)	0.000	0.0058 (10)	0.000
N6	0.0460 (13)	0.0188 (9)	0.0504 (13)	0.000	0.0006 (10)	0.000
N7	0.0293 (7)	0.0333 (8)	0.0505 (10)	0.0011 (6)	0.0040 (7)	-0.0052 (7)
C1	0.0282 (9)	0.0412 (10)	0.0476 (11)	0.0067 (7)	-0.0031 (8)	-0.0008 (8)
C2	0.0270 (8)	0.0305 (8)	0.0371 (9)	0.0010 (6)	-0.0011 (7)	-0.0017 (7)
C3	0.0237 (8)	0.0296 (8)	0.0374 (9)	0.0006 (6)	0.0012 (7)	-0.0023 (7)
C4	0.0264 (8)	0.0324 (9)	0.0424 (10)	0.0001 (7)	0.0008 (7)	0.0007 (7)
C5	0.0288 (9)	0.0410 (10)	0.0517 (12)	-0.0062 (7)	0.0027 (8)	0.0011 (9)
C6	0.0339 (9)	0.0289 (8)	0.0406 (10)	0.0063 (7)	0.0034 (7)	-0.0019 (7)
C7	0.0261 (8)	0.0269 (8)	0.0339 (9)	-0.0002 (6)	0.0009 (7)	-0.0015 (6)
C8	0.0258 (11)	0.0233 (10)	0.0316 (12)	0.000	-0.0020 (9)	0.000
C9	0.0387 (10)	0.0270 (8)	0.0421 (10)	0.0055 (7)	0.0003 (8)	-0.0018 (7)
C10	0.0296 (8)	0.0239 (8)	0.0369 (9)	0.0001 (6)	-0.0006 (7)	-0.0009 (6)
C11	0.0281 (11)	0.0228 (10)	0.0323 (12)	0.000	-0.0021 (9)	0.000

Geometric parameters (Å, °)

O1—N1	1.2285 (19)	N5—C6	1.341 (2)
O2—N1	1.228 (2)	N5—C6 ⁱ	1.341 (2)
O3—C3	1.235 (2)	N5—H5	0.848 (10)
O4—N2	1.210 (2)	N6—C9	1.338 (2)
O5—N2	1.206 (2)	N6—C9 ⁱⁱ	1.338 (2)
O6—N4	1.227 (2)	N6—H6	0.854 (10)
O7—N4	1.225 (2)	N7—C10	1.455 (2)
O8—C8	1.233 (3)	C1—C2	1.356 (2)
O9—N7	1.223 (2)	C1—H1A	0.9300
O10—N7	1.223 (2)	C2—C3	1.457 (2)
O11—C11	1.237 (3)	C3—C4	1.457 (2)
O1W—H11	0.851 (9)	C4—C5	1.362 (2)
O1W—H12	0.858 (9)	C5—H5A	0.9300
O2W—H21	0.840 (8)	C6—C7	1.361 (2)
O3W—H31	0.844 (8)	C6—H6A	0.9300
N1—C2	1.457 (2)	C7—C8	1.458 (2)
N2—C4	1.455 (2)	C8—C7 ⁱ	1.458 (2)
N3—C1	1.338 (3)	C9—C10	1.360 (2)
N3—C5	1.338 (3)	C9—H9	0.9300
N3—H3	0.853 (10)	C10—C11	1.454 (2)
N4—C7	1.455 (2)	C11—C10 ⁱⁱ	1.454 (2)

H11—O1W—H12	109.1 (14)	O3—C3—C4	125.31 (15)
O2—N1—O1	123.08 (16)	O3—C3—C2	124.72 (15)
O2—N1—C2	119.15 (14)	C4—C3—C2	109.97 (14)
O1—N1—C2	117.76 (15)	C5—C4—N2	115.48 (16)
O5—N2—O4	121.96 (17)	C5—C4—C3	123.38 (16)
O5—N2—C4	118.53 (17)	N2—C4—C3	121.12 (15)
O4—N2—C4	119.51 (15)	N3—C5—C4	121.24 (17)
C1—N3—C5	120.47 (16)	N3—C5—H5A	119.4
C1—N3—H3	117.6 (16)	C4—C5—H5A	119.4
C5—N3—H3	121.8 (16)	N5—C6—C7	120.80 (16)
O7—N4—O6	123.11 (15)	N5—C6—H6A	119.6
O7—N4—C7	118.20 (15)	C7—C6—H6A	119.6
O6—N4—C7	118.66 (14)	C6—C7—N4	115.51 (15)
C6—N5—C6 ⁱ	120.6 (2)	C6—C7—C8	124.09 (16)
C6—N5—H5	119.71 (10)	N4—C7—C8	120.39 (14)
C6 ⁱ —N5—H5	119.71 (10)	O8—C8—C7 ⁱ	125.23 (9)
C9—N6—C9 ⁱⁱ	120.4 (2)	O8—C8—C7	125.23 (9)
C9—N6—H6	119.82 (11)	C7 ⁱ —C8—C7	109.54 (19)
C9 ⁱⁱ —N6—H6	119.82 (10)	N6—C9—C10	120.94 (17)
O10—N7—O9	123.05 (16)	N6—C9—H9	119.5
O10—N7—C10	118.79 (14)	C10—C9—H9	119.5
O9—N7—C10	118.16 (16)	C9—C10—C11	124.11 (16)
N3—C1—C2	120.99 (17)	C9—C10—N7	114.74 (15)
N3—C1—H1A	119.5	C11—C10—N7	121.15 (14)
C2—C1—H1A	119.5	O11—C11—C10 ⁱⁱ	125.25 (9)
C1—C2—N1	115.69 (15)	O11—C11—C10	125.25 (9)
C1—C2—C3	123.94 (16)	C10 ⁱⁱ —C11—C10	109.50 (19)
N1—C2—C3	120.36 (14)		
C5—N3—C1—C2	-0.1 (3)	C6 ⁱ —N5—C6—C7	-1.47 (13)
N3—C1—C2—N1	179.48 (18)	N5—C6—C7—N4	-176.15 (14)
N3—C1—C2—C3	0.5 (3)	N5—C6—C7—C8	3.0 (3)
O2—N1—C2—C1	152.19 (18)	O7—N4—C7—C6	-26.7 (2)
O1—N1—C2—C1	-27.0 (3)	O6—N4—C7—C6	151.55 (18)
O2—N1—C2—C3	-28.8 (3)	O7—N4—C7—C8	154.12 (15)
O1—N1—C2—C3	151.95 (18)	O6—N4—C7—C8	-27.7 (2)
C1—C2—C3—O3	179.23 (19)	C6—C7—C8—O8	178.50 (13)
N1—C2—C3—O3	0.3 (3)	N4—C7—C8—O8	-2.34 (18)
C1—C2—C3—C4	-0.1 (3)	C6—C7—C8—C7 ⁱ	-1.50 (13)
N1—C2—C3—C4	-179.00 (16)	N4—C7—C8—C7 ⁱ	177.66 (18)
O5—N2—C4—C5	-16.1 (3)	C9 ⁱⁱ —N6—C9—C10	-0.98 (13)
O4—N2—C4—C5	164.46 (19)	N6—C9—C10—C11	2.0 (3)
O5—N2—C4—C3	162.5 (3)	N6—C9—C10—N7	-177.52 (15)
O4—N2—C4—C3	-16.9 (3)	O10—N7—C10—C9	152.08 (18)
O3—C3—C4—C5	179.9 (2)	O9—N7—C10—C9	-27.5 (3)
C2—C3—C4—C5	-0.7 (3)	O10—N7—C10—C11	-27.5 (2)
O3—C3—C4—N2	1.4 (3)	O9—N7—C10—C11	152.96 (17)

supplementary materials

C2—C3—C4—N2	-179.23 (17)	C9—C10—C11—O11	179.01 (13)
C1—N3—C5—C4	-0.7 (3)	N7—C10—C11—O11	-1.48 (19)
N2—C4—C5—N3	179.75 (19)	C9—C10—C11—C10 ⁱⁱ	-0.99 (13)
C3—C4—C5—N3	1.2 (3)	N7—C10—C11—C10 ⁱⁱ	178.52 (19)

Symmetry codes: (i) $-x+1, y, -z+3/2$; (ii) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N3—H3 \cdots O1w	0.85 (1)	1.86 (1)	2.703 (2)	172 (2)
N5—H5 \cdots O2w	0.85 (1)	1.84 (1)	2.692 (3)	180
N6—H6 \cdots O3w	0.85 (1)	1.87 (1)	2.723 (3)	180
O1w—H11 \cdots O8 ⁱⁱⁱ	0.85 (1)	2.04 (1)	2.878 (2)	168 (2)
O1w—H12 \cdots O11 ^{iv}	0.86 (1)	2.02 (1)	2.866 (2)	168 (2)
O2w—H21 \cdots O3 ^v	0.84 (1)	2.05 (1)	2.888 (2)	173 (1)
O3w—H31 \cdots O3	0.84 (1)	2.05 (1)	2.890 (2)	172 (2)

Symmetry codes: (iii) $-x+3/2, -y+3/2, z-1/2$; (iv) $x+1/2, y-1/2, -z+1/2$; (v) $-x+1, -y+1, -z+1$.

Fig. 1

