

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# Bis[2-(2-pyridylmethyleamino)-benzenesulfonato- $\kappa^3N,N',O$ ]zinc(II) dihydrate

 Cheng-Xiang Cai,<sup>a,b</sup> Miao Ou-Yang,<sup>a</sup> Zhi-Yuan Zhao<sup>a</sup> and Yi-Min Jiang<sup>a\*</sup>

<sup>a</sup>College of Chemistry and Chemical Engineering, Guangxi Normal University, Guilin, Guangxi 541004, People's Republic of China, and <sup>b</sup>Department of Chemistry and Life Science, Baie University, Baie 533000, People's Republic of China  
Correspondence e-mail: ouyangmiao123456@126.com


Received 26 July 2008; accepted 14 August 2008

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(C-C) = 0.003$  Å; R factor = 0.027; wR factor = 0.075; data-to-parameter ratio = 13.6.

In the title complex,  $[Zn(C_{12}H_9N_2O_3S)_2] \cdot 2H_2O$ , the Zn<sup>II</sup> ion lies on a crystallographic inversion center and is coordinated by four N atoms and two O atoms from two tridentate 2-(2-pyridylmethyleamino)benzenesulfonate ligands in a slightly distorted octahedral environment. In the crystal structure, the complex forms a two-dimensional network through intermolecular O—H...O and C—H...O hydrogen bonds.

## Related literature

For related literature, see: Casella & Gullotti (1981, 1986); Jiang *et al.* (2006); Li *et al.* (2006, 2007); Wang *et al.* (1994); Zhang *et al.* (2004, 2007, 2008); Correia *et al.* (2003); Zheng *et al.* (2001); Zhou *et al.* (2004).



## Experimental

### Crystal data

$[Zn(C_{12}H_9N_2O_3S)_2] \cdot 2H_2O$   
 $M_r = 623.95$   
 Orthorhombic, *Pbcn*  
 $a = 19.7090$  (15) Å

$b = 8.0722$  (6) Å  
 $c = 16.3390$  (13) Å  
 $V = 2599.5$  (3) Å<sup>3</sup>  
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 1.16$  mm<sup>-1</sup>

$T = 295$  (2) K  
 $0.49 \times 0.45 \times 0.37$  mm

### Data collection

Bruker SMART CCD area-detector diffractometer  
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $T_{min} = 0.600$ ,  $T_{max} = 0.673$

17894 measured reflections  
 2412 independent reflections  
 2100 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.024$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$   
 $wR(F^2) = 0.074$   
 $S = 1.04$   
 2412 reflections

177 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{max} = 0.46$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.42$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

D—H...A	D—H	H...A	D...A	D—H...A
O4—H1W...O3 <sup>i</sup>	0.83	2.21	3.014 (3)	162
O4—H2W...O2	0.83	2.06	2.877 (3)	166
C4—H4...O3 <sup>iii</sup>	0.93	2.48	3.407 (3)	175
C6—H6...O4 <sup>iii</sup>	0.93	2.57	3.436 (3)	155

Symmetry codes: (i)  $-x + \frac{1}{2}, y + \frac{1}{2}, z$ ; (ii)  $-x + 1, y + 1, -z + \frac{1}{2}$ ; (iii)  $x, -y + 2, z - \frac{1}{2}$ .

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was funded by Guangxi Science Foundation, Guangxi Zhuang Autonomous Region of the People's Republic of China (grant No. 0731053).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PK2112).

## References

- Bruker (2004). *SMART* and *SAINT*. Bruker AXS inc., Madison, WI, USA.
- Casella, L. & Gullotti, M. (1981). *J. Am. Chem. Soc.* **103**, 6338–6347.
- Casella, L. & Gullotti, M. (1986). *Inorg. Chem.* **25**, 1293–1303.
- Correia, V. R., Bortoluzzi, A. J., Neves, A., Joussef, A. C., Vieira, M. G. M. & Batista, S. C. (2003). *Acta Cryst.* **E59**, m464–m466.
- Jiang, Y.-M., Li, J.-M., Xie, F.-Q. & Wang, Y.-F. (2006). *Chin. J. Struct. Chem.* **25**, 767–770.
- Li, J.-X., Jiang, Y.-M. & Li, H.-Y. (2006). *Acta Cryst.* **E62**, m2984–m2986.
- Li, H.-Y., Liao, B.-L., Jiang, Y.-M., Zhang, S.-H. & Li, J.-X. (2007). *Chin. J. Struct. Chem.* **26**, 907–910.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, Z., Wu, Z., Yen, Z., Le, Z., Zhu, X. & Huang, Q. (1994). *Synth. React. Inorg. Metal Org. Chem.* **24**, 1453–1460.
- Zhang, S. H., Jiang, Y.-M. & Liu, Z. M. (2008). *J. Coord. Chem.* **61**, 1927–1934.
- Zhang, S.-H., Jiang, Y.-M., Liu, Z. & Zhou, Z.-Y. (2004). *Chin. J. Struct. Chem.* **23**, 882–885.
- Zhang, S. H., Li, G.-Z., Zhong, F. & Feng, X.-Z. (2007). *Chin. J. Struct. Chem.* **26**, 1491–1494.
- Zheng, S.-L., Tong, M.-L., Yu, X.-L. & Chen, X.-M. (2001). *J. Chem. Soc. Dalton Trans.* pp. 586–592.
- Zhou, J.-S., Cai, J.-W., Wang, L. & Ng, S.-W. (2004). *Dalton Trans.* pp. 1493–1497.

**supplementary materials**

*Acta Cryst.* (2008). E64, m1195 [ doi:10.1107/S1600536808026342 ]

## Bis[2-(2-pyridylmethyleneamino)benzenesulfonato- $\kappa^3$ N,N',O]zinc(II) dihydrate

C.-X. Cai, M. Ou-Yang, Z.-Y. Zhao and Y.-M. Jiang

### Comment

The design and control of supermolecular coordination complex networks in which both coordination bonds and hydrogen bonds take part in the self-assembly chemistry (Zheng, *et al.*, 2001; Zhou, *et al.*, 2004) have recently garnered increasing interest. Schiff base complexes that contain both sulfur and amino acid functionalities have received much attention owing to their potential applications in pharmacy. (Casella & Gullotti, 1981, 1986; Wang *et al.*, 1994; Li *et al.*, 2006; Zhang *et al.*, 2007, 2008).

Our group has focused on the exploration of the coordination chemistry of the sulfonate ligands for years (Zhang *et al.* 2004; Jiang *et al.* 2006; Li *et al.* 2007). We report here the synthesis and the structure of the mononuclear Zn<sup>II</sup> Paba complex (Fig. 1). The structure is composed of one Zn<sup>II</sup>, two deprotonated Paba<sup>-</sup> ligands and two guest water molecules. The six-coordinated Zn<sup>II</sup> atom has a distorted octahedral geometry, being coordinated by pyridine N, imine N and sulfonate O atoms from two deprotonated Paba<sup>-</sup> ligands in a tridentate facial arrangement. This structure is similar to those reported for complexes with N,N',O-tridentate donor ligands (Li *et al.*, 2006; Correia *et al.*, 2003).

There are extensive hydrogen bonds (O4-H2W $\cdots$ O2 and O4-H1W $\cdots$ O3), in which the donor is O-H of the guest water and S=O acts as acceptor, which forms a two-dimension sheet structure (Fig. 2).

### Experimental

The potassium salt of 2-(pyridylmethyl)imine-2-benzenesulfonic acid (PabaK) was synthesized according to the literature method (Casella & Gullotti, 1986).

To prepare the title complex, the ligand PabaK (1 mmol, 0.30 g) was dissolved in methanol (10 mL) at 333 K and an aqueous solution (10 mL) containing ZnCl<sub>2</sub>(0.5 mmol, 0.068 g) was added. The resulting solution was stirred at 333 K for 4 h, then cooled to room temperature and filtered. Yellow crystals suitable for X-ray diffraction were obtained by slow evaporation over several days, with a yield of 55%. Elemental analysis, found (%): C, 46.05; H, 3.55; N, 8.95; S, 10.42; calc (%): C, 46.16; H, 3.53; N, 8.98; S, 10.26.

### Refinement

H atoms bonded to C atoms were positioned geometrically with the C-H distance of 0.93 Å, and treated as riding atoms, with U<sub>iso</sub>(H) = 1.2U<sub>eq</sub>(C). Water hydrogens were placed in fixed positions and assigned U<sub>iso</sub> values of 1.5U<sub>eq</sub> of the water oxygen atom.

## Figures




Fig. 1. An ellipsoid plot (30% probability) showing the numbering scheme. Dashed lines indicate hydrogen bonds. Symmetry code: (a)  $-x+1, y, -z+1/2$ .




Fig. 2. 2-D structure, as viewed down the a axis. Dashed lines indicate hydrogen bonds.

## Bis[2-(2-pyridylmethyleneamino)benzenesulfonato- $\kappa^3N,N',O$ ]zinc(II) dihydrate

### Crystal data

$[\text{Zn}(\text{C}_{12}\text{H}_9\text{N}_2\text{O}_3\text{S}_1)_2] \cdot 2\text{H}_2\text{O}$

$M_r = 623.95$

Orthorhombic, *Pbcn*

Hall symbol:  $-P\ 2n\ 2ab$

$a = 19.7090\ (15)\ \text{\AA}$

$b = 8.0722\ (6)\ \text{\AA}$

$c = 16.3390\ (13)\ \text{\AA}$

$V = 2599.5\ (3)\ \text{\AA}^3$

$Z = 4$

$F_{000} = 1280$

$D_x = 1.594\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 7323 reflections

$\theta = 2.5\text{--}28.2^\circ$

$\mu = 1.16\ \text{mm}^{-1}$

$T = 295\ (2)\ \text{K}$

Block, yellow

$0.49 \times 0.45 \times 0.37\ \text{mm}$

### Data collection

Bruker SMART CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295\ (2)\ \text{K}$

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.600, T_{\max} = 0.673$

17894 measured reflections

2412 independent reflections

2100 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 25.5^\circ$

$\theta_{\min} = 2.5^\circ$

$h = -23 \rightarrow 23$

$k = -9 \rightarrow 9$

$l = -19 \rightarrow 19$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.026$	H-atom parameters constrained
$wR(F^2) = 0.074$	$w = 1/[\sigma^2(F_o^2) + (0.0373P)^2 + 1.7627P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
2412 reflections	$(\Delta/\sigma)_{\max} < 0.001$
177 parameters	$\Delta\rho_{\max} = 0.46 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.42 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

*Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.82446 (4)	0.2500	0.02769 (11)
S1	0.37616 (2)	0.67719 (6)	0.34320 (3)	0.03070 (14)
O1	0.45098 (7)	0.67650 (18)	0.33822 (8)	0.0344 (3)
O2	0.34865 (8)	0.84229 (19)	0.33446 (10)	0.0456 (4)
O3	0.35256 (7)	0.5876 (2)	0.41443 (8)	0.0431 (4)
N1	0.49613 (8)	1.0216 (2)	0.16032 (10)	0.0341 (4)
N2	0.40820 (8)	0.7687 (2)	0.17245 (9)	0.0297 (4)
C1	0.44459 (10)	1.0145 (3)	0.10605 (12)	0.0332 (4)
C2	0.43673 (13)	1.1301 (3)	0.04460 (14)	0.0452 (6)
H2	0.4007	1.1223	0.0081	0.054*
C3	0.48322 (13)	1.2579 (3)	0.03809 (15)	0.0506 (6)
H3	0.4786	1.3380	-0.0025	0.061*
C4	0.53638 (13)	1.2645 (3)	0.09261 (15)	0.0483 (6)
H4	0.5688	1.3480	0.0890	0.058*
C5	0.54084 (12)	1.1447 (3)	0.15295 (14)	0.0420 (5)
H5	0.5766	1.1503	0.1900	0.050*
C6	0.39854 (10)	0.8728 (3)	0.11481 (12)	0.0354 (5)
H6	0.3626	0.8590	0.0786	0.042*

## supplementary materials

---

C7	0.36755 (9)	0.6238 (3)	0.17777 (12)	0.0313 (4)
C8	0.34696 (11)	0.5353 (3)	0.10925 (13)	0.0437 (5)
H8	0.3579	0.5736	0.0572	0.052*
C9	0.31015 (12)	0.3900 (3)	0.11817 (15)	0.0524 (6)
H9	0.2962	0.3320	0.0720	0.063*
C10	0.29391 (13)	0.3306 (3)	0.19512 (16)	0.0516 (6)
H10	0.2698	0.2322	0.2005	0.062*
C11	0.31364 (11)	0.4180 (3)	0.26416 (13)	0.0403 (5)
H11	0.3022	0.3792	0.3160	0.048*
C12	0.35039 (10)	0.5632 (3)	0.25581 (11)	0.0300 (4)
O4	0.29845 (11)	1.0939 (4)	0.44343 (14)	0.1079 (10)
H1W	0.2572	1.1144	0.4394	0.162*
H2W	0.3083	1.0105	0.4158	0.162*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.02732 (18)	0.03263 (19)	0.02311 (18)	0.000	-0.00389 (11)	0.000
S1	0.0291 (3)	0.0396 (3)	0.0234 (2)	-0.0001 (2)	0.00018 (18)	-0.0016 (2)
O1	0.0287 (7)	0.0462 (9)	0.0285 (7)	-0.0028 (6)	-0.0034 (6)	0.0038 (6)
O2	0.0482 (9)	0.0452 (9)	0.0436 (9)	0.0102 (7)	-0.0018 (7)	-0.0079 (7)
O3	0.0416 (8)	0.0620 (10)	0.0257 (7)	-0.0062 (8)	0.0039 (6)	0.0035 (7)
N1	0.0400 (9)	0.0346 (9)	0.0278 (9)	-0.0007 (7)	-0.0057 (7)	0.0011 (7)
N2	0.0278 (8)	0.0375 (9)	0.0238 (8)	-0.0006 (7)	0.0004 (6)	0.0004 (7)
C1	0.0343 (10)	0.0377 (11)	0.0275 (10)	0.0035 (9)	-0.0036 (8)	0.0023 (8)
C2	0.0519 (14)	0.0471 (13)	0.0367 (12)	0.0017 (11)	-0.0113 (10)	0.0096 (10)
C3	0.0728 (17)	0.0386 (13)	0.0403 (13)	-0.0023 (12)	-0.0064 (12)	0.0105 (11)
C4	0.0628 (15)	0.0363 (12)	0.0458 (13)	-0.0117 (11)	-0.0019 (11)	0.0022 (10)
C5	0.0490 (13)	0.0391 (12)	0.0380 (12)	-0.0084 (10)	-0.0102 (10)	0.0012 (9)
C6	0.0310 (11)	0.0475 (12)	0.0276 (10)	0.0001 (9)	-0.0057 (8)	0.0034 (9)
C7	0.0245 (10)	0.0410 (11)	0.0284 (10)	-0.0009 (8)	-0.0004 (8)	-0.0015 (9)
C8	0.0417 (12)	0.0609 (15)	0.0286 (11)	-0.0085 (11)	0.0014 (9)	-0.0050 (10)
C9	0.0505 (14)	0.0679 (16)	0.0389 (13)	-0.0180 (13)	-0.0001 (11)	-0.0169 (12)
C10	0.0464 (14)	0.0552 (15)	0.0530 (15)	-0.0211 (12)	0.0021 (11)	-0.0085 (12)
C11	0.0363 (12)	0.0495 (13)	0.0351 (11)	-0.0088 (10)	0.0038 (9)	0.0017 (10)
C12	0.0232 (9)	0.0399 (11)	0.0270 (10)	-0.0007 (8)	-0.0001 (7)	-0.0023 (8)
O4	0.0592 (13)	0.168 (3)	0.0964 (18)	0.0323 (15)	-0.0209 (12)	-0.0725 (19)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Zn1—O1	2.1065 (14)	C3—C4	1.376 (3)
Zn1—O1 <sup>i</sup>	2.1065 (14)	C3—H3	0.9300
Zn1—N1 <sup>i</sup>	2.1643 (18)	C4—C5	1.384 (3)
Zn1—N1	2.1643 (18)	C4—H4	0.9300
Zn1—N2	2.2544 (16)	C5—H5	0.9300
Zn1—N2 <sup>i</sup>	2.2544 (16)	C6—H6	0.9300
S1—O2	1.4459 (16)	C7—C8	1.389 (3)
S1—O3	1.4470 (15)	C7—C12	1.407 (3)

S1—O1	1.4769 (14)	C8—C9	1.387 (3)
S1—C12	1.7730 (19)	C8—H8	0.9300
N1—C5	1.334 (3)	C9—C10	1.383 (3)
N1—C1	1.350 (2)	C9—H9	0.9300
N2—C6	1.277 (3)	C10—C11	1.386 (3)
N2—C7	1.421 (3)	C10—H10	0.9300
C1—C2	1.380 (3)	C11—C12	1.384 (3)
C1—C6	1.467 (3)	C11—H11	0.9300
C2—C3	1.384 (3)	O4—H1W	0.8332
C2—H2	0.9300	O4—H2W	0.8339
O1—Zn1—O1 <sup>i</sup>	110.92 (8)	C3—C2—H2	120.5
O1—Zn1—N1 <sup>i</sup>	88.27 (6)	C4—C3—C2	118.9 (2)
O1 <sup>i</sup> —Zn1—N1 <sup>i</sup>	149.76 (6)	C4—C3—H3	120.6
O1—Zn1—N1	149.76 (6)	C2—C3—H3	120.6
O1 <sup>i</sup> —Zn1—N1	88.27 (6)	C3—C4—C5	118.9 (2)
N1 <sup>i</sup> —Zn1—N1	85.36 (9)	C3—C4—H4	120.6
O1—Zn1—N2	84.45 (5)	C5—C4—H4	120.6
O1 <sup>i</sup> —Zn1—N2	82.55 (5)	N1—C5—C4	122.9 (2)
N1 <sup>i</sup> —Zn1—N2	123.75 (6)	N1—C5—H5	118.5
N1—Zn1—N2	74.81 (6)	C4—C5—H5	118.5
O1—Zn1—N2 <sup>i</sup>	82.55 (5)	N2—C6—C1	119.53 (18)
O1 <sup>i</sup> —Zn1—N2 <sup>i</sup>	84.45 (5)	N2—C6—H6	120.2
N1 <sup>i</sup> —Zn1—N2 <sup>i</sup>	74.81 (6)	C1—C6—H6	120.2
N1—Zn1—N2 <sup>i</sup>	123.75 (6)	C8—C7—C12	118.80 (19)
N2—Zn1—N2 <sup>i</sup>	156.97 (9)	C8—C7—N2	122.62 (18)
O2—S1—O3	114.81 (10)	C12—C7—N2	118.51 (17)
O2—S1—O1	111.87 (9)	C9—C8—C7	120.2 (2)
O3—S1—O1	111.31 (9)	C9—C8—H8	119.9
O2—S1—C12	106.94 (9)	C7—C8—H8	119.9
O3—S1—C12	107.23 (9)	C10—C9—C8	120.7 (2)
O1—S1—C12	103.86 (9)	C10—C9—H9	119.7
S1—O1—Zn1	119.57 (8)	C8—C9—H9	119.7
C5—N1—C1	118.00 (18)	C9—C10—C11	119.9 (2)
C5—N1—Zn1	125.86 (14)	C9—C10—H10	120.1
C1—N1—Zn1	116.11 (14)	C11—C10—H10	120.1
C6—N2—C7	120.22 (17)	C12—C11—C10	119.8 (2)
C6—N2—Zn1	113.76 (14)	C12—C11—H11	120.1
C7—N2—Zn1	125.63 (12)	C10—C11—H11	120.1
N1—C1—C2	122.3 (2)	C11—C12—C7	120.62 (18)
N1—C1—C6	115.76 (17)	C11—C12—S1	120.65 (15)
C2—C1—C6	121.93 (19)	C7—C12—S1	118.73 (15)
C1—C2—C3	119.1 (2)	H1W—O4—H2W	110.2
C1—C2—H2	120.5		

Symmetry codes: (i)  $-x+1, y, -z+1/2$ .

## supplementary materials

---

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O4—H1W $\cdots$ O3 <sup>ii</sup>	0.83	2.21	3.014 (3)	162
O4—H2W $\cdots$ O2	0.83	2.06	2.877 (3)	166
C4—H4 $\cdots$ O3 <sup>iii</sup>	0.93	2.48	3.407 (3)	175
C6—H6 $\cdots$ O4 <sup>iv</sup>	0.93	2.57	3.436 (3)	155

Symmetry codes: (ii)  $-x+1/2, y+1/2, z$ ; (iii)  $-x+1, y+1, -z+1/2$ ; (iv)  $x, -y+2, z-1/2$ .

Fig. 1




Fig. 2

