

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

A second monoclinic polymorph of 2-amino-4,6-dichloropyrimidine

 Hoong-Kun Fun,^{a*} Suchada Chantrapromma,^{b‡} Subrata Jana,^c Rinku Chakrabarty^c and Shyamaprosad Goswami^c

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, ^bCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand, and ^cDepartment of Chemistry, Bengal Engineering and Science University, Shibpur, Howrah, India 711 103

Correspondence e-mail: hkfun@usm.my


Received 25 July 2008; accepted 27 July 2008

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.098; data-to-parameter ratio = 15.4.

The title chloro-substituted 2-aminopyrimidine, $\text{C}_4\text{H}_3\text{Cl}_2\text{N}_3$, is a second monoclinic polymorph of this compound which crystallizes in the space group $C2/c$. The structure was previously reported [Clews & Cochran (1948). *Acta Cryst.* **1**, 4–11] in the space group $P21/a$. There are two crystallographically independent molecules in the asymmetric unit and each molecule is planar. The dihedral angle between the two pyrimidine rings is 30.71 (12)°. In the crystal structure, molecules are linked *via* $\text{N}-\text{H}\cdots\text{N}$ intermolecular hydrogen bonds, forming infinite one-dimensional chains along the a axis. These hydrogen bonds generate $R_2^2(8)$ ring motifs. The chains are stacked along the b axis.

Related literature

For bond-length data, see: Allen *et al.* (1987). For details of hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see: the polymorph reported by Clews & Cochran (1948); Low *et al.* (2002). For applications of pyrimidine compounds and their supramolecular chemistry, see, for example: Blackburn & Gait (1996); Brown (1988); Hurst (1980); Goswami *et al.* (2008a,b); Lighthart *et al.* (2005); Sherington & Taskinen (2001).



‡ Additional correspondence author, email: suchada.c@psu.ac.th.

Experimental

Crystal data

$\text{C}_4\text{H}_3\text{Cl}_2\text{N}_3$	$V = 2539.6$ (6) Å ³
$M_r = 163.99$	$Z = 16$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 32.060$ (4) Å	$\mu = 0.92$ mm ⁻¹
$b = 3.8045$ (6) Å	$T = 296$ (2) K
$c = 21.302$ (3) Å	$0.57 \times 0.14 \times 0.02$ mm
$\beta = 102.193$ (7)°	

Data collection

Bruker SMART APEX2 CCD area-detector diffractometer	12772 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2005)	2886 independent reflections
$T_{\min} = 0.620$, $T_{\max} = 0.985$	1875 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$	187 parameters
$wR(F^2) = 0.098$	All H-atom parameters refined
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.22$ e Å ⁻³
2886 reflections	$\Delta\rho_{\text{min}} = -0.24$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3A}-\text{H2NA}\cdots\text{N1A}^i$	0.75 (3)	2.43 (3)	3.172 (3)	176 (2)
$\text{N3A}-\text{H1NA}\cdots\text{N2B}^i$	0.87 (3)	2.33 (3)	3.201 (3)	172 (2)
$\text{N3B}-\text{H1NB}\cdots\text{N2A}^i$	0.87 (3)	2.39 (3)	3.253 (4)	174 (3)
$\text{N3B}-\text{H2NB}\cdots\text{N1B}^{ii}$	0.84 (3)	2.41 (3)	3.242 (3)	172 (3)

 Symmetry codes: (i) $-x + 1, -y + 2, -z + 1$; (ii) $-x + \frac{3}{2}, -y + \frac{3}{2}, -z + 1$.

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: SAINT (Bruker, 2005); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

SJ, RC and SG acknowledge the DST [SR/S1/OC-13/2005] and CSIR [01(1913)/04/EMR-II], Government of India for financial support. SJ and RC thank the CSIR, Government of India, for research fellowships. The authors also thank Universiti Sains Malaysia for the Research University Golden Goose Grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2524).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Blackburn, G. M. & Gait, M. J. (1996). *Nucleic Acids in Chemistry and Biology*. Editors. Oxford University Press.
- Brown, D. J. (1988). *Fused Pyrimidines The Chemistry of Heterocyclic Compounds*, Vol. 24, pt. 3. New York: John Wiley & Sons.
- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Clews, C. J. B. & Cochran, W. (1948). *Acta Cryst.* **1**, 4–11.
- Goswami, S., Jana, S., Das, N. K., Fun, H.-K. & Chantrapromma, S. (2008a). *J. Mol. Struct.* **876**, 313–321.

- Goswami, S., Jana, S., Hazra, A., Fun, H.-K. & Chantrapromma, S. (2008b). *Supramol. Chem.* **20**, 495–500.
- Hurst, D. T. (1980). *Chemistry and Biochemistry of Pyrimidines, Purines, Pteridines*. Chichester: Wiley.
- Ligthart, G. B. W. L., Ohkawa, H., Sijbesma, R. P. & Meijer, E. W. (2005). *J. Am. Chem. Soc.* **127**, 810–811.
- Low, J. N., Quesada, A., Marchal, A., Melguizo, M., Nogueras, M. & Glidewell, C. (2002). *Acta Cryst.* **C58**, o289–o294.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sherrington, D. C. & Taskinen, K. A. (2001). *Chem. Soc. Rev.* **30**, 83–93.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o1659-o1660 [doi:10.1107/S1600536808023714]

A second monoclinic polymorph of 2-amino-4,6-dichloropyrimidine

H.-K. Fun, S. Chantrapromma, S. Jana, R. Chakrabarty and S. Goswami

Comment

Functionalized pyrimidines play a major role in the synthesis of different drug molecules and of naturally occurring pyrimidine bases (Blackburn & Gait, 1996; Brown, 1988; Hurst, 1980). Substituted pyrimidines are also very important for studies on multiple hydrogen bonding interactions in molecular recognition and supramolecular chemistry (Sherrington & Taskinen, 2001; Goswami *et al.*, 2008*a,b*; Lighthart *et al.*, 20050). In this work we report the crystal structure of the title compound, Fig 1, which is a second monoclinic polymorph of 2-amino-4,6-dichloropyrimidine.

The crystal structure of the title compound (I) was previously reported by Clews & Cochran (1948) in the monoclinic space group $P21/a$, with $a = 16.447$, $b = 3.845$, $c = 10.283$ Å, $\beta = 107.58^\circ$ and $Z = 4$. In the present work, the compound crystallized out in the monoclinic space group $C2/c$ with $Z = 16$. There are two crystallographically independent molecules in the asymmetric unit, A and B , (see Fig. 1) with slightly different bond lengths and bond angles. Both molecules A and B are planar with maximum deviations of 0.005 (2) Å for atom N2A in A and 0.009 (2) Å for atom C2B in B . The dihedral angle between the two pyrimidine rings is 30.71 (12) $^\circ$. The amino group acts as a double donor in N—H \cdots N hydrogen bonds, while the two ring N atoms (N1 and N2) act as the acceptors. The molecules are linked *via* N—H \cdots N intermolecular hydrogen bonds to form infinite one-dimensional chains along the a axis, Table 1. These hydrogen bonds generate $R^2_2(8)$ ring motifs (Bernstein *et al.*, 1995) (Fig. 2). Interestingly, the Cl atoms do not form N—H \cdots Cl hydrogen bonds. The closest Cl \cdots Cl distance is 3.3635 (11) Å [3.37 Å in Clews & Cochran (1948)]. The bond lengths and angles in (I) are within normal ranges (Allen *et al.*, 1987) and comparable to those found in related structures (Clews & Cochran, 1948; Low *et al.*, 2002).

In the crystal packing shown in Fig. 2, the [1 0 0] molecular chains are stacked along the b axis.

Experimental

Phosphorus oxy-chloride (POCl₃) (25 ml) was added to anhydrous 2-amino-4,6-dioxypyrimidine (6 g) and the mixture refluxed at 383 K for 12 h. Excess POCl₃ was distilled off. The solid residue was neutralized using KOH solution in an ice bath and saturated NaHCO₃ solution was added. The solid residue was filtered off, extracted with CHCl₃ and the solution was dried over Na₂SO₄ and then concentrated under vacuum. The crude product was purified by column chromatography using 20% ethyl acetate in petroleum ether as eluent and the title compound (I) (4.29 g, 61%) was isolated. Single crystals were grown by slow evaporation of a CH₂Cl₂/ethanol (*v/v* 3:1) solution, Mp. 492–494 K.

Refinement

All H atoms were located in a difference map and freely refined isotropically. The highest residual electron density peak is located at 1.00 Å from N2A and the deepest hole is located at 0.81 Å from H2NA.

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atomic numbering.




Fig. 2. The crystal packing of (I), viewed approximately along the *b* axis showing one-dimensional chains along the *a* axis. Hydrogen bonds were shown as dashed lines.

A second monoclinic polymorph of 2-amino-4,6-dichloropyrimidine

Crystal data

$C_4H_3Cl_2N_3$

$M_r = 163.99$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 32.060\ (4)\ \text{\AA}$

$b = 3.8045\ (6)\ \text{\AA}$

$c = 21.302\ (3)\ \text{\AA}$

$\beta = 102.193\ (7)^\circ$

$V = 2539.6\ (6)\ \text{\AA}^3$

$Z = 16$

$F_{000} = 1312$

$D_x = 1.716\ \text{Mg m}^{-3}$

Melting point = 492–494 K

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2886 reflections

$\theta = 1.3\text{--}27.5^\circ$

$\mu = 0.92\ \text{mm}^{-1}$

$T = 296\ (2)\ \text{K}$

Block, colorless

$0.57 \times 0.14 \times 0.02\ \text{mm}$

Data collection

Bruker SMART APEX2 CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: $8.33\ \text{pixels mm}^{-1}$

$T = 296\ (2)\ \text{K}$

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.620$, $T_{\max} = 0.985$

12772 measured reflections

2886 independent reflections

1875 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.051$

$\theta_{\max} = 27.5^\circ$

$\theta_{\min} = 1.3^\circ$

$h = -40 \rightarrow 40$

$k = -4 \rightarrow 4$

$l = -27 \rightarrow 27$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	All H-atom parameters refined
$wR(F^2) = 0.098$	$w = 1/[\sigma^2(F_o^2) + (0.0403P)^2]$
$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
2886 reflections	$(\Delta/\sigma)_{\max} = 0.001$
187 parameters	$\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11A	0.330814 (19)	0.45738 (18)	0.35093 (3)	0.0475 (2)
C12A	0.49584 (2)	0.5271 (2)	0.33992 (4)	0.0572 (2)
N1A	0.46421 (6)	0.7708 (5)	0.43348 (9)	0.0371 (5)
N2A	0.38986 (6)	0.7401 (5)	0.43849 (9)	0.0349 (5)
N3A	0.43941 (8)	0.9927 (7)	0.51907 (11)	0.0490 (6)
H2NA	0.4619 (8)	1.056 (7)	0.5293 (12)	0.033 (8)*
H1NA	0.4174 (9)	1.026 (7)	0.5366 (14)	0.057 (9)*
C1A	0.38306 (7)	0.5786 (6)	0.38271 (11)	0.0338 (6)
C2A	0.41389 (8)	0.5008 (7)	0.34867 (12)	0.0377 (6)
H2A	0.4102 (7)	0.386 (6)	0.3120 (11)	0.034 (7)*
C3A	0.45423 (7)	0.6088 (7)	0.37766 (11)	0.0353 (6)
C4A	0.43110 (7)	0.8296 (7)	0.46279 (11)	0.0350 (6)
C11B	0.58263 (2)	1.02542 (19)	0.31034 (3)	0.0510 (2)
C12B	0.73894 (2)	0.50031 (18)	0.32094 (3)	0.0474 (2)
N1B	0.71199 (6)	0.7404 (6)	0.41918 (9)	0.0389 (5)
N2B	0.64127 (6)	0.9690 (6)	0.41463 (9)	0.0406 (5)
N3B	0.69119 (9)	0.9534 (8)	0.50908 (11)	0.0589 (7)
H1NB	0.6706 (10)	1.049 (8)	0.5237 (16)	0.080 (12)*

supplementary materials

H2NB	0.7156 (10)	0.881 (9)	0.5264 (16)	0.075 (11)*
C1B	0.63348 (7)	0.9103 (6)	0.35235 (12)	0.0364 (6)
C2B	0.66172 (7)	0.7709 (7)	0.31888 (11)	0.0374 (6)
H2B	0.6550 (7)	0.753 (7)	0.2743 (12)	0.047 (7)*
C3B	0.70066 (7)	0.6887 (7)	0.35702 (11)	0.0361 (6)
C4B	0.68134 (8)	0.8851 (7)	0.44634 (11)	0.0403 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11A	0.0320 (3)	0.0586 (5)	0.0491 (4)	-0.0065 (3)	0.0027 (3)	-0.0029 (3)
C12A	0.0442 (4)	0.0698 (5)	0.0658 (5)	-0.0003 (4)	0.0298 (3)	-0.0132 (4)
N1A	0.0293 (10)	0.0422 (13)	0.0402 (11)	0.0003 (10)	0.0086 (9)	-0.0014 (11)
N2A	0.0295 (10)	0.0416 (13)	0.0339 (10)	0.0028 (10)	0.0073 (8)	-0.0013 (10)
N3A	0.0351 (14)	0.0707 (19)	0.0418 (13)	-0.0073 (14)	0.0093 (11)	-0.0161 (13)
C1A	0.0293 (12)	0.0320 (14)	0.0385 (13)	-0.0009 (11)	0.0040 (10)	0.0039 (11)
C2A	0.0378 (14)	0.0402 (16)	0.0355 (13)	-0.0012 (13)	0.0087 (11)	-0.0057 (13)
C3A	0.0338 (13)	0.0354 (14)	0.0396 (13)	0.0033 (12)	0.0141 (10)	0.0001 (12)
C4A	0.0327 (13)	0.0404 (15)	0.0315 (12)	-0.0020 (12)	0.0060 (10)	0.0013 (12)
C11B	0.0344 (4)	0.0669 (5)	0.0514 (4)	0.0072 (4)	0.0083 (3)	0.0056 (4)
C12B	0.0358 (3)	0.0566 (4)	0.0546 (4)	-0.0001 (3)	0.0204 (3)	-0.0103 (3)
N1B	0.0355 (11)	0.0438 (13)	0.0386 (11)	0.0036 (11)	0.0109 (9)	0.0016 (10)
N2B	0.0388 (12)	0.0481 (14)	0.0377 (11)	0.0044 (11)	0.0143 (9)	0.0001 (11)
N3B	0.0534 (17)	0.087 (2)	0.0368 (13)	0.0187 (16)	0.0102 (12)	-0.0034 (13)
C1B	0.0315 (13)	0.0386 (15)	0.0407 (13)	0.0001 (12)	0.0112 (10)	0.0014 (12)
C2B	0.0352 (14)	0.0465 (17)	0.0328 (13)	-0.0025 (13)	0.0123 (11)	-0.0036 (13)
C3B	0.0329 (13)	0.0369 (15)	0.0416 (14)	-0.0031 (12)	0.0151 (11)	-0.0009 (12)
C4B	0.0398 (15)	0.0477 (16)	0.0352 (13)	0.0043 (13)	0.0122 (11)	0.0023 (12)

Geometric parameters (\AA , $^\circ$)

C11A—C1A	1.731 (2)	C11B—C1B	1.742 (2)
C12A—C3A	1.725 (2)	C12B—C3B	1.735 (2)
N1A—C3A	1.317 (3)	N1B—C3B	1.312 (3)
N1A—C4A	1.359 (3)	N1B—C4B	1.358 (3)
N2A—C1A	1.314 (3)	N2B—C1B	1.316 (3)
N2A—C4A	1.357 (3)	N2B—C4B	1.357 (3)
N3A—C4A	1.326 (3)	N3B—C4B	1.332 (3)
N3A—H2NA	0.75 (2)	N3B—H1NB	0.87 (3)
N3A—H1NA	0.87 (3)	N3B—H2NB	0.84 (3)
C1A—C2A	1.376 (3)	C1B—C2B	1.372 (3)
C2A—C3A	1.373 (3)	C2B—C3B	1.374 (3)
C2A—H2A	0.88 (2)	C2B—H2B	0.93 (2)
C3A—N1A—C4A	115.3 (2)	C3B—N1B—C4B	114.8 (2)
C1A—N2A—C4A	115.11 (19)	C1B—N2B—C4B	114.9 (2)
C4A—N3A—H2NA	114 (2)	C4B—N3B—H1NB	114 (2)
C4A—N3A—H1NA	115.4 (19)	C4B—N3B—H2NB	112 (2)
H2NA—N3A—H1NA	130 (3)	H1NB—N3B—H2NB	134 (3)

N2A—C1A—C2A	125.2 (2)	N2B—C1B—C2B	125.7 (2)
N2A—C1A—C11A	116.12 (18)	N2B—C1B—C11B	115.67 (18)
C2A—C1A—C11A	118.68 (19)	C2B—C1B—C11B	118.62 (19)
C3A—C2A—C1A	114.3 (2)	C1B—C2B—C3B	113.4 (2)
C3A—C2A—H2A	118.9 (14)	C1B—C2B—H2B	121.6 (15)
C1A—C2A—H2A	126.7 (15)	C3B—C2B—H2B	124.8 (14)
N1A—C3A—C2A	124.9 (2)	N1B—C3B—C2B	125.9 (2)
N1A—C3A—C12A	116.14 (18)	N1B—C3B—C12B	115.99 (17)
C2A—C3A—C12A	118.95 (19)	C2B—C3B—C12B	118.12 (18)
N3A—C4A—N2A	117.1 (2)	N3B—C4B—N2B	116.9 (2)
N3A—C4A—N1A	117.7 (2)	N3B—C4B—N1B	117.8 (2)
N2A—C4A—N1A	125.2 (2)	N2B—C4B—N1B	125.3 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N3A—H2NA \cdots N1A ⁱ	0.75 (3)	2.43 (3)	3.172 (3)	176 (2)
N3A—H1NA \cdots N2B ⁱ	0.87 (3)	2.33 (3)	3.201 (3)	172 (2)
N3B—H1NB \cdots N2A ⁱ	0.87 (3)	2.39 (3)	3.253 (4)	174 (3)
N3B—H2NB \cdots N1B ⁱⁱ	0.84 (3)	2.41 (3)	3.242 (3)	172 (3)

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $-x+3/2, -y+3/2, -z+1$.

Fig. 1




Fig. 2

