

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Diaqua- μ_3 -oxido-hexakis(μ_2 -trichloroacetato- κ^2 O:O')(trichloroacetato- κ O)-trichromium(III) acetonitrile trisolvate

B. B. Mougang D. Soume, Rosiyah Yahya, Seng Neon Gan and Seik Weng Ng*

 Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
 Correspondence e-mail: seikweng@um.edu.my


Received 8 August 2008; accepted 11 August 2008

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; disorder in main residue; R factor = 0.042; wR factor = 0.119; data-to-parameter ratio = 18.8.

In the crystal structure of the title compound, $[\text{Cr}_3(\text{C}_2\text{Cl}_3\text{O}_2)_7\text{O}(\text{H}_2\text{O})_2] \cdot 3\text{CH}_3\text{CN}$, the trinuclear $[\text{Cr}_3\text{O}(\text{H}_2\text{O})_2(\text{Cl}_3\text{CCO}_2)_7]$ molecule has an oxide O atom that is connected to one monodentate trichloroacetate-coordinated and two water-coordinated Cr^{III} atoms, the three metal atoms forming the points of an equilateral triangle. Each of the six remaining carboxylate groups bridges a $\text{Cr}-\text{O}-\text{Cr}$ fragment. The cluster interacts with the three solvent molecules through water-acetonitrile $\text{O}-\text{H} \cdots \text{N}$ hydrogen bonds. Adjacent clusters are linked by a water-carboxylate $\text{O}-\text{H} \cdots \text{O}$ hydrogen bond to give a helical chain. One of the CCl_3 groups was found to be disordered over two positions, with the major component having a site-occupancy factor of 0.64 (1).

Related literature

Oxo-centred chromium(III) chloroacetates form an efficient class of Ziegler-Natta catalysts for the polymerization of olefins; see: Gan *et al.* (2000).



Experimental

Crystal data

$[\text{Cr}_3(\text{C}_2\text{Cl}_3\text{O}_2)_7\text{O}(\text{H}_2\text{O})_2] \cdot 3\text{C}_2\text{H}_3\text{N}$
 $M_r = 1467.78$
 Monoclinic, $P2_1/n$
 $a = 11.6307$ (6) Å
 $b = 19.481$ (1) Å
 $c = 22.949$ (1) Å
 $\beta = 95.355$ (1)°
 $V = 5177.0$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.76$ mm⁻¹
 $T = 100$ (2) K
 $0.25 \times 0.20 \times 0.15$ mm

Data collection

Bruker SMART APEX diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.667$, $T_{\text{max}} = 0.778$
 29504 measured reflections
 11718 independent reflections
 9590 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.118$
 $S = 1.04$
 11718 reflections
 624 parameters
 72 restraints
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 1.42$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.74$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O1w}-\text{H1w1} \cdots \text{N1}$	0.84 (1)	1.97 (2)	2.791 (4)	166 (5)
$\text{O1w}-\text{H1w2} \cdots \text{N2}$	0.85 (1)	1.97 (1)	2.810 (5)	173 (4)
$\text{O2w}-\text{H2w1} \cdots \text{N3}$	0.85 (1)	1.88 (1)	2.719 (5)	171 (4)
$\text{O2w}-\text{H2w2} \cdots \text{O2}^i$	0.83 (1)	1.81 (1)	2.613 (3)	165 (4)

 Symmetry code: (i) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

The authors thank the University of Malaya for supporting this study (grant Nos. PS078-2007C and PS208-2008A).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2293).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Gan, S.-N., Jalan, A. M. & Ooi, C. P. (2000). Trinuclear oxo-centered chromium(III) carboxylate complexes as Ziegler-Natta catalysts for ethylene polymerization in *Progress and Development of Catalytic Olefin Polymerization*, edited by T. Sano, T. Uozumi, H. Nakatani & M. Terano, pp. 25–32. Tokyo: Technology and Education Publishers.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2008). publCIF. In preparation.

supplementary materials

Acta Cryst. (2008). E64, m1175 [doi:10.1107/S1600536808025798]

Diaqua- μ_3 -oxido-hexakis(μ_2 -trichloroacetato- $\kappa^2O:O'$)(trichloroacetato- κO)trichromium(III) acetonitrile trisolvate

B. B. M. D. Soume, R. Yahya, S. N. Gan and S. W. Ng

Comment

Oxo-centered chromium(III) chloroacetates form an efficient class of Ziegler–Natta catalysts for the polymerization of olefins (Gan *et al.*, 2000). The title trichloroacetate oxo-cluster crystallizes with acetonitrile (Scheme I, Fig. 1). In the crystal structure, the oxo-O atom is connected to one monodentate trichloroacetate-coordinated and two water-coordinated chromium(III) atoms, the three metal atoms forming the points of an equilateral triangle. Each of the six remaining carboxylate groups chelates a Cr–O–Cr fragment. The cluster interacts with the three solvate molecules through hydrogen bonds; hydrogen bonds involving the water molecule as a donor give rise to a helical chain that runs along the *b*-axis.

Experimental

Chromium(III) chloride hexahydrate (10 g) was refluxed with trichloroacetic acid in a molar ratio of 1:6 for 6 h. The solution was filtered hot; the cooled solution yielded a green product that was washed with chloroform (90% yield). Analysis found: Cr, 11.13, C, 11.98, H, 0.72; $\text{Cr}_3\text{Cl}_{21}\text{C}_{14}\text{O}_{20}\text{H}_{10}$ requires: Cr, 11.15, C, 12.02, H, 0.72. Dark-green crystals of the acetonitrile solvate were obtained by recrystallization from an acetonitrile solution.

Refinement

One of the seven trichloroacetate groups was found to be disordered over two sites. The six C—Cl distances were restrained to within 0.01 Å of each other, as were the Cl···Cl distances. The anisotropic displacement parameters of the disordered Cl atoms were restrained to be nearly isotropic. The disorder refined to a 0.636 (12):0.364 (12) site occupancy ratio. The water hydrogen atoms were located in a difference Fourier map, and were refined with a distance restraint of O—H 0.84 (1) Å; their temperature factors were freely refined. The methyl-H atoms were generated geometrically (C—H = 0.98 Å), and were included in the refinement in the riding model approximation with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. The final difference Fourier map had a large peak in the vicinity of the disordered Cl atoms, but was otherwise featureless.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) plot of $[\text{Cr}_3\text{O}(\text{H}_2\text{O})_2(\text{Cl}_3\text{CCO}_2)_7].3\text{CH}_3\text{CN}$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius. For clarity, the minor disorder component is not shown. Hydrogen bonds are denoted by dashed lines.

Diaqua- μ_3 -oxido-hexakis(μ_2 -trichloroacetato- κ^2 O:O')(trichloroacetato- κ O)trichromium(III) acetonitrile trisolvate

Crystal data

$[\text{Cr}_3(\text{C}_2\text{Cl}_3\text{O}_2)_7\text{O}(\text{H}_2\text{O})_2] \cdot 3\text{C}_2\text{H}_3\text{N}$

$M_r = 1467.78$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 11.6307$ (6) Å

$b = 19.481$ (1) Å

$c = 22.949$ (1) Å

$\beta = 95.355$ (1)°

$V = 5177.0$ (5) Å³

$Z = 4$

$F_{000} = 2876$

$D_x = 1.883$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 9535 reflections

$\theta = 2.2$ – 28.2 °

$\mu = 1.76$ mm⁻¹

$T = 100$ (2) K

Block, green

$0.25 \times 0.20 \times 0.15$ mm

Data collection

Bruker SMART APEX
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.667$, $T_{\max} = 0.778$

29504 measured reflections

11718 independent reflections

9590 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\text{max}} = 27.5$ °

$\theta_{\text{min}} = 1.4$ °

$h = -15$ → 11

$k = -25$ → 25

$l = -29$ → 29

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.118$

$S = 1.04$

11718 reflections

624 parameters

72 restraints

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring
sites

H atoms treated by a mixture of
independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0626P)^2 + 7.2035P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.001$

$\Delta\rho_{\text{max}} = 1.42$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.74$ e Å⁻³

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cr1	0.74255 (4)	0.76778 (3)	0.70072 (2)	0.01404 (12)	
Cr2	0.68641 (4)	0.73747 (3)	0.83719 (2)	0.01277 (11)	
Cr3	0.72390 (4)	0.60693 (3)	0.74830 (2)	0.01179 (11)	
Cl1	0.96872 (8)	0.85657 (6)	0.56393 (4)	0.0333 (2)	
Cl2	0.89615 (9)	0.99687 (5)	0.57400 (4)	0.0345 (2)	
Cl3	0.73830 (8)	0.89677 (4)	0.52068 (3)	0.02158 (18)	
Cl4	0.36046 (10)	0.88419 (6)	0.80350 (5)	0.0431 (3)	
Cl5	0.35756 (9)	0.81589 (5)	0.69151 (5)	0.0430 (3)	
Cl6	0.46821 (8)	0.94801 (4)	0.70989 (4)	0.02438 (19)	
Cl7	0.98178 (10)	0.88074 (7)	0.90381 (5)	0.0451 (3)	
Cl8	1.09295 (9)	0.78569 (5)	0.83228 (5)	0.0346 (2)	
Cl9	1.03479 (9)	0.92225 (5)	0.78947 (5)	0.0356 (2)	
Cl10	0.44418 (9)	0.70978 (5)	0.55064 (4)	0.0322 (2)	
Cl11	0.38086 (9)	0.61375 (7)	0.63683 (5)	0.0399 (3)	
Cl12	0.53908 (9)	0.57331 (5)	0.55303 (4)	0.0336 (2)	
Cl13	0.9998 (4)	0.6385 (3)	0.58065 (14)	0.0556 (10)	0.636 (12)
Cl14	1.1108 (2)	0.71821 (13)	0.67552 (16)	0.0352 (8)	0.636 (12)
Cl15	1.0981 (8)	0.5711 (3)	0.6842 (4)	0.0316 (13)	0.636 (12)
Cl3'	0.9729 (10)	0.6101 (9)	0.5847 (3)	0.105 (5)	0.364 (12)
Cl4'	1.1068 (6)	0.7141 (3)	0.6463 (8)	0.103 (3)	0.364 (12)
Cl5'	1.1046 (14)	0.5774 (5)	0.6926 (7)	0.0252 (17)	0.364 (12)
Cl16	0.31914 (8)	0.66992 (5)	0.78820 (4)	0.0287 (2)	
Cl17	0.35454 (8)	0.52405 (5)	0.79999 (4)	0.0292 (2)	
Cl18	0.37880 (7)	0.61005 (4)	0.90215 (3)	0.01931 (17)	
Cl19	0.93022 (8)	0.51554 (5)	0.92623 (4)	0.0272 (2)	
Cl20	0.99064 (8)	0.65247 (6)	0.96451 (4)	0.0344 (2)	
Cl21	1.07222 (7)	0.60054 (5)	0.85829 (4)	0.02607 (19)	
O1	0.7710 (2)	0.82999 (12)	0.63619 (10)	0.0187 (5)	
O2	0.8135 (3)	0.93428 (14)	0.67484 (11)	0.0281 (6)	
O3	0.6252 (2)	0.83313 (12)	0.72363 (10)	0.0179 (5)	
O4	0.55621 (19)	0.79526 (12)	0.80588 (10)	0.0163 (5)	
O5	0.8720 (2)	0.81086 (13)	0.74973 (10)	0.0197 (5)	
O6	0.8001 (2)	0.81248 (12)	0.83713 (10)	0.0184 (5)	
O7	0.6175 (2)	0.72533 (12)	0.64643 (10)	0.0178 (5)	
O8	0.6429 (2)	0.61373 (12)	0.66921 (9)	0.0160 (5)	
O9	0.8600 (2)	0.71060 (12)	0.66684 (10)	0.0184 (5)	
O10	0.8769 (2)	0.61126 (12)	0.71653 (10)	0.0168 (5)	
O11	0.57304 (19)	0.66600 (12)	0.85012 (10)	0.0155 (5)	
O12	0.56962 (19)	0.58956 (12)	0.77652 (9)	0.0147 (5)	
O13	0.81379 (19)	0.68469 (12)	0.87790 (10)	0.0169 (5)	
O14	0.81080 (19)	0.58767 (12)	0.82463 (9)	0.0151 (5)	
O15	0.71611 (19)	0.70351 (11)	0.76204 (9)	0.0136 (4)	
O1W	0.6544 (2)	0.77209 (13)	0.91690 (10)	0.0190 (5)	
H1W1	0.684 (3)	0.747 (2)	0.9440 (15)	0.059 (17)*	
H1W2	0.5872 (17)	0.783 (2)	0.9250 (17)	0.034 (13)*	

supplementary materials

O2W	0.7303 (2)	0.50636 (12)	0.73377 (10)	0.0171 (5)
H2W1	0.759 (4)	0.4871 (18)	0.7053 (11)	0.032 (12)*
H2W2	0.726 (4)	0.4790 (17)	0.7610 (12)	0.050 (15)*
N1	0.7157 (3)	0.67991 (19)	1.00764 (16)	0.0338 (8)
N2	0.4284 (4)	0.7968 (3)	0.9455 (2)	0.0556 (12)
N3	0.8447 (5)	0.4523 (3)	0.6464 (2)	0.0685 (15)
C1	0.8074 (3)	0.89117 (17)	0.63661 (14)	0.0157 (6)
C2	0.8515 (3)	0.91082 (17)	0.57647 (14)	0.0172 (7)
C3	0.5520 (3)	0.82935 (16)	0.75960 (14)	0.0158 (6)
C4	0.4390 (3)	0.86978 (19)	0.74313 (17)	0.0229 (7)
C5	0.8757 (3)	0.82309 (17)	0.80318 (15)	0.0173 (7)
C6	0.9912 (3)	0.85391 (19)	0.83110 (15)	0.0215 (7)
C7	0.5957 (3)	0.66336 (18)	0.64181 (13)	0.0156 (6)
C8	0.4943 (3)	0.64175 (18)	0.59620 (15)	0.0196 (7)
C9	0.9091 (3)	0.65627 (17)	0.68302 (14)	0.0161 (6)
C10	1.0241 (3)	0.64354 (16)	0.65623 (14)	0.0254 (8)
C11	0.5281 (3)	0.62049 (17)	0.81711 (14)	0.0142 (6)
C12	0.3999 (3)	0.60421 (18)	0.82725 (14)	0.0177 (7)
C13	0.8484 (3)	0.62681 (17)	0.86503 (14)	0.0144 (6)
C14	0.9565 (3)	0.59973 (18)	0.90309 (15)	0.0187 (7)
C15	0.7356 (3)	0.6309 (2)	1.03291 (16)	0.0255 (8)
C16	0.7599 (4)	0.5680 (2)	1.06581 (18)	0.0342 (9)
H16A	0.8074	0.5785	1.1023	0.051*
H16B	0.8016	0.5360	1.0424	0.051*
H16C	0.6872	0.5470	1.0750	0.051*
C17	0.3472 (4)	0.7913 (2)	0.9685 (2)	0.0423 (11)
C18	0.2451 (4)	0.7862 (3)	0.9985 (3)	0.0504 (13)
H18A	0.2630	0.7620	1.0356	0.076*
H18B	0.1858	0.7609	0.9742	0.076*
H18C	0.2167	0.8324	1.0062	0.076*
C19	0.8283 (8)	0.4449 (4)	0.5933 (4)	0.091 (2)
C20	0.8044 (9)	0.4336 (6)	0.5267 (3)	0.126 (4)
H20A	0.8704	0.4102	0.5119	0.189*
H20B	0.7925	0.4781	0.5071	0.189*
H20C	0.7350	0.4053	0.5187	0.189*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cr1	0.0128 (3)	0.0158 (3)	0.0139 (2)	0.0004 (2)	0.00299 (19)	0.00348 (19)
Cr2	0.0104 (2)	0.0147 (3)	0.0134 (2)	-0.00009 (19)	0.00208 (18)	0.00052 (19)
Cr3	0.0095 (2)	0.0149 (3)	0.0112 (2)	0.00076 (19)	0.00208 (18)	0.00201 (18)
Cl1	0.0189 (5)	0.0495 (6)	0.0329 (5)	0.0045 (4)	0.0097 (4)	-0.0016 (4)
Cl2	0.0441 (6)	0.0272 (5)	0.0306 (5)	-0.0214 (4)	-0.0047 (4)	0.0062 (4)
Cl3	0.0244 (4)	0.0247 (4)	0.0147 (4)	-0.0056 (3)	-0.0031 (3)	-0.0002 (3)
Cl4	0.0320 (6)	0.0450 (6)	0.0567 (7)	0.0210 (5)	0.0277 (5)	0.0231 (5)
Cl5	0.0313 (6)	0.0279 (5)	0.0645 (7)	-0.0076 (4)	-0.0240 (5)	0.0102 (5)
Cl6	0.0246 (4)	0.0168 (4)	0.0320 (5)	0.0038 (3)	0.0037 (3)	0.0075 (3)

Cl7	0.0338 (6)	0.0700 (8)	0.0317 (5)	-0.0181 (5)	0.0043 (4)	-0.0204 (5)
Cl8	0.0231 (5)	0.0291 (5)	0.0492 (6)	0.0034 (4)	-0.0098 (4)	0.0042 (4)
Cl9	0.0253 (5)	0.0303 (5)	0.0495 (6)	-0.0113 (4)	-0.0057 (4)	0.0172 (4)
Cl10	0.0359 (5)	0.0282 (5)	0.0291 (5)	0.0015 (4)	-0.0152 (4)	0.0076 (4)
Cl11	0.0192 (5)	0.0669 (8)	0.0334 (5)	-0.0113 (5)	0.0016 (4)	0.0109 (5)
Cl12	0.0399 (6)	0.0328 (5)	0.0258 (5)	0.0067 (4)	-0.0081 (4)	-0.0095 (4)
Cl13	0.0395 (16)	0.112 (3)	0.0170 (11)	0.0297 (15)	0.0124 (9)	0.0041 (11)
Cl14	0.0124 (10)	0.0279 (10)	0.0660 (18)	-0.0058 (7)	0.0066 (10)	0.0156 (10)
Cl15	0.0214 (19)	0.0290 (15)	0.047 (3)	0.0084 (12)	0.0165 (17)	0.0106 (15)
Cl3'	0.076 (6)	0.221 (13)	0.020 (2)	0.086 (7)	0.014 (3)	-0.001 (5)
Cl4'	0.050 (3)	0.058 (3)	0.214 (9)	0.025 (2)	0.084 (5)	0.081 (4)
Cl5'	0.017 (2)	0.030 (3)	0.029 (3)	0.012 (2)	0.0058 (17)	0.008 (2)
Cl16	0.0158 (4)	0.0420 (6)	0.0283 (5)	0.0072 (4)	0.0009 (3)	0.0121 (4)
Cl17	0.0231 (5)	0.0326 (5)	0.0336 (5)	-0.0133 (4)	0.0119 (4)	-0.0121 (4)
Cl18	0.0165 (4)	0.0251 (4)	0.0174 (4)	-0.0001 (3)	0.0070 (3)	0.0011 (3)
Cl19	0.0242 (5)	0.0303 (5)	0.0259 (4)	0.0024 (4)	-0.0043 (3)	0.0115 (4)
Cl20	0.0232 (5)	0.0452 (6)	0.0318 (5)	0.0122 (4)	-0.0136 (4)	-0.0189 (4)
Cl21	0.0117 (4)	0.0285 (5)	0.0387 (5)	0.0009 (3)	0.0060 (3)	-0.0024 (4)
O1	0.0216 (13)	0.0186 (12)	0.0162 (11)	-0.0034 (10)	0.0031 (9)	0.0034 (9)
O2	0.0416 (17)	0.0240 (14)	0.0190 (13)	0.0025 (12)	0.0051 (11)	-0.0055 (10)
O3	0.0166 (12)	0.0177 (12)	0.0201 (12)	0.0016 (9)	0.0046 (9)	0.0038 (9)
O4	0.0140 (12)	0.0170 (12)	0.0182 (11)	0.0018 (9)	0.0027 (9)	0.0030 (9)
O5	0.0162 (12)	0.0244 (13)	0.0186 (12)	-0.0050 (10)	0.0022 (9)	0.0053 (10)
O6	0.0168 (12)	0.0189 (12)	0.0201 (12)	-0.0048 (9)	0.0053 (9)	-0.0020 (9)
O7	0.0180 (12)	0.0196 (12)	0.0156 (11)	0.0002 (9)	0.0007 (9)	0.0041 (9)
O8	0.0174 (12)	0.0169 (12)	0.0135 (11)	0.0015 (9)	0.0003 (9)	0.0011 (9)
O9	0.0163 (12)	0.0211 (12)	0.0186 (12)	0.0029 (10)	0.0064 (9)	0.0052 (9)
O10	0.0132 (11)	0.0190 (12)	0.0188 (11)	0.0022 (9)	0.0049 (9)	0.0041 (9)
O11	0.0138 (11)	0.0172 (12)	0.0158 (11)	-0.0017 (9)	0.0039 (9)	0.0000 (9)
O12	0.0114 (11)	0.0182 (12)	0.0148 (11)	-0.0002 (9)	0.0032 (8)	0.0013 (9)
O13	0.0129 (11)	0.0203 (12)	0.0172 (11)	0.0032 (9)	-0.0001 (9)	-0.0012 (9)
O14	0.0132 (11)	0.0164 (11)	0.0154 (11)	0.0001 (9)	-0.0001 (9)	0.0023 (9)
O15	0.0124 (11)	0.0150 (11)	0.0138 (11)	0.0027 (9)	0.0035 (8)	0.0023 (8)
O1W	0.0174 (13)	0.0230 (13)	0.0173 (12)	0.0018 (10)	0.0050 (10)	-0.0007 (10)
O2W	0.0216 (13)	0.0163 (12)	0.0140 (11)	0.0016 (10)	0.0046 (9)	0.0022 (9)
N1	0.0257 (18)	0.038 (2)	0.038 (2)	0.0025 (15)	0.0060 (15)	0.0135 (16)
N2	0.047 (3)	0.076 (3)	0.046 (3)	0.011 (2)	0.013 (2)	0.000 (2)
N3	0.089 (4)	0.057 (3)	0.067 (3)	-0.005 (3)	0.046 (3)	-0.022 (3)
C1	0.0133 (16)	0.0204 (17)	0.0135 (15)	0.0014 (13)	0.0016 (12)	0.0020 (12)
C2	0.0181 (17)	0.0179 (16)	0.0154 (15)	-0.0046 (13)	0.0001 (13)	0.0015 (12)
C3	0.0132 (16)	0.0124 (15)	0.0217 (16)	0.0001 (12)	0.0014 (12)	0.0016 (12)
C4	0.0183 (18)	0.0192 (18)	0.031 (2)	0.0029 (14)	0.0032 (14)	0.0066 (14)
C5	0.0150 (16)	0.0137 (15)	0.0234 (17)	-0.0021 (12)	0.0023 (13)	0.0028 (13)
C6	0.0162 (17)	0.0268 (19)	0.0212 (17)	-0.0045 (14)	0.0004 (13)	0.0014 (14)
C7	0.0144 (16)	0.0225 (17)	0.0106 (14)	0.0022 (13)	0.0040 (12)	0.0026 (12)
C8	0.0195 (17)	0.0212 (17)	0.0177 (16)	0.0015 (14)	0.0005 (13)	0.0033 (13)
C9	0.0112 (15)	0.0218 (17)	0.0152 (15)	0.0011 (13)	0.0013 (12)	-0.0005 (12)
C10	0.0178 (18)	0.031 (2)	0.0286 (19)	0.0065 (15)	0.0093 (14)	0.0119 (15)
C11	0.0100 (15)	0.0182 (16)	0.0145 (15)	0.0012 (12)	0.0022 (11)	0.0065 (12)

supplementary materials

C12	0.0139 (16)	0.0223 (17)	0.0175 (16)	0.0006 (13)	0.0049 (12)	0.0013 (13)
C13	0.0083 (15)	0.0206 (17)	0.0146 (15)	-0.0007 (12)	0.0022 (11)	0.0023 (12)
C14	0.0129 (16)	0.0236 (18)	0.0191 (16)	0.0021 (13)	-0.0012 (13)	-0.0023 (13)
C15	0.0207 (19)	0.032 (2)	0.0237 (18)	-0.0009 (16)	0.0027 (14)	0.0001 (16)
C16	0.046 (3)	0.028 (2)	0.028 (2)	0.0033 (18)	-0.0004 (18)	0.0012 (16)
C17	0.036 (3)	0.039 (3)	0.052 (3)	0.008 (2)	0.005 (2)	-0.004 (2)
C18	0.035 (3)	0.036 (3)	0.082 (4)	0.002 (2)	0.018 (3)	0.008 (2)
C19	0.110 (6)	0.059 (4)	0.112 (7)	0.002 (4)	0.052 (5)	-0.005 (4)
C20	0.133 (9)	0.177 (10)	0.073 (6)	0.024 (7)	0.036 (5)	-0.032 (6)

Geometric parameters (Å, °)

Cr1—O15	1.930 (2)	Cl21—C14	1.768 (4)
Cr1—O1	1.965 (2)	O1—C1	1.264 (4)
Cr1—O3	1.973 (2)	O2—C1	1.212 (4)
Cr1—O9	1.978 (2)	O3—C3	1.242 (4)
Cr1—O5	1.980 (2)	O4—C3	1.250 (4)
Cr1—O7	2.003 (2)	O5—C5	1.246 (4)
Cr2—O15	1.909 (2)	O6—C5	1.246 (4)
Cr2—O11	1.959 (2)	O7—C7	1.236 (4)
Cr2—O13	1.966 (2)	O8—C7	1.252 (4)
Cr2—O4	1.968 (2)	O9—C9	1.243 (4)
Cr2—O6	1.971 (2)	O10—C9	1.247 (4)
Cr2—O1W	2.017 (2)	O11—C11	1.249 (4)
Cr3—O15	1.911 (2)	O12—C11	1.244 (4)
Cr3—O8	1.970 (2)	O13—C13	1.242 (4)
Cr3—O14	1.975 (2)	O14—C13	1.248 (4)
Cr3—O10	1.986 (2)	O1W—H1W1	0.839 (10)
Cr3—O2W	1.990 (2)	O1W—H1W2	0.849 (10)
Cr3—O12	1.993 (2)	O2W—H2W1	0.850 (10)
Cl1—C2	1.769 (4)	O2W—H2W2	0.826 (10)
Cl2—C2	1.757 (3)	N1—C15	1.131 (5)
Cl3—C2	1.770 (3)	N2—C17	1.130 (6)
Cl4—C4	1.752 (4)	N3—C19	1.225 (10)
Cl5—C4	1.787 (4)	C1—C2	1.564 (4)
Cl6—C4	1.752 (4)	C3—C4	1.549 (5)
Cl7—C6	1.762 (4)	C5—C6	1.554 (5)
Cl8—C6	1.778 (4)	C7—C8	1.560 (5)
Cl9—C6	1.741 (4)	C9—C10	1.543 (5)
Cl10—C8	1.754 (3)	C11—C12	1.562 (4)
Cl11—C8	1.772 (4)	C13—C14	1.555 (4)
Cl12—C8	1.768 (4)	C15—C16	1.453 (5)
Cl13—C10	1.734 (4)	C16—H16A	0.9800
Cl14—C10	1.802 (4)	C16—H16B	0.9800
Cl15—C10	1.744 (5)	C16—H16C	0.9800
Cl3'—C10	1.813 (6)	C17—C18	1.429 (7)
Cl4'—C10	1.705 (5)	C18—H18A	0.9800
Cl5'—C10	1.757 (6)	C18—H18B	0.9800
Cl16—C12	1.780 (4)	C18—H18C	0.9800

Cl17—C12	1.745 (4)	C19—C20	1.544 (11)
Cl18—C12	1.762 (3)	C20—H20A	0.9800
Cl19—C14	1.759 (4)	C20—H20B	0.9800
Cl20—C14	1.759 (3)	C20—H20C	0.9800
O15—Cr1—O1	177.60 (10)	C3—C4—Cl6	110.7 (2)
O15—Cr1—O3	93.62 (9)	Cl4—C4—Cl6	110.0 (2)
O1—Cr1—O3	88.40 (10)	C3—C4—Cl5	104.4 (2)
O15—Cr1—O9	94.83 (9)	Cl4—C4—Cl5	109.6 (2)
O1—Cr1—O9	83.11 (10)	Cl6—C4—Cl5	109.6 (2)
O3—Cr1—O9	171.35 (10)	O6—C5—O5	128.5 (3)
O15—Cr1—O5	91.45 (10)	O6—C5—C6	116.1 (3)
O1—Cr1—O5	89.66 (10)	O5—C5—C6	115.4 (3)
O3—Cr1—O5	94.78 (10)	C5—C6—C19	110.7 (2)
O9—Cr1—O5	86.86 (10)	C5—C6—C17	112.2 (2)
O15—Cr1—O7	91.71 (9)	Cl9—C6—Cl7	109.9 (2)
O1—Cr1—O7	87.10 (10)	C5—C6—Cl8	105.4 (2)
O3—Cr1—O7	87.27 (10)	Cl9—C6—Cl8	110.6 (2)
O9—Cr1—O7	90.63 (10)	Cl7—C6—Cl8	107.91 (19)
O5—Cr1—O7	176.12 (10)	O7—C7—O8	129.4 (3)
O15—Cr2—O11	94.11 (10)	O7—C7—C8	117.2 (3)
O15—Cr2—O13	93.23 (9)	O8—C7—C8	113.4 (3)
O11—Cr2—O13	92.54 (10)	C7—C8—Cl10	112.6 (2)
O15—Cr2—O4	93.81 (9)	C7—C8—Cl12	109.6 (2)
O11—Cr2—O4	87.48 (10)	Cl10—C8—Cl12	109.56 (19)
O13—Cr2—O4	172.95 (10)	C7—C8—Cl11	106.5 (2)
O15—Cr2—O6	94.50 (10)	Cl10—C8—Cl11	108.88 (19)
O11—Cr2—O6	171.16 (10)	Cl12—C8—Cl11	109.6 (2)
O13—Cr2—O6	84.97 (10)	O9—C9—O10	128.8 (3)
O4—Cr2—O6	93.95 (10)	O9—C9—C10	114.1 (3)
O15—Cr2—O1W	179.24 (10)	C9—C10—Cl4'	116.4 (4)
O11—Cr2—O1W	85.21 (10)	C9—C10—Cl13	110.0 (3)
O13—Cr2—O1W	86.47 (10)	C9—C10—Cl15	113.5 (4)
O4—Cr2—O1W	86.50 (10)	Cl4'—C10—Cl15	115.9 (5)
O6—Cr2—O1W	86.17 (10)	Cl13—C10—Cl15	110.6 (4)
O15—Cr3—O8	93.43 (9)	C9—C10—Cl5'	111.8 (7)
O15—Cr3—O14	93.95 (9)	Cl4'—C10—Cl5'	111.8 (5)
O8—Cr3—O14	172.40 (10)	Cl13—C10—Cl5'	117.6 (7)
O15—Cr3—O10	94.39 (10)	C9—C10—Cl14	105.0 (2)
O8—Cr3—O10	91.54 (10)	Cl13—C10—Cl14	109.1 (2)
O14—Cr3—O10	86.09 (10)	Cl15—C10—Cl14	108.5 (3)
O15—Cr3—O2W	179.43 (10)	Cl5'—C10—Cl14	102.3 (5)
O8—Cr3—O2W	86.22 (10)	C9—C10—Cl3'	101.2 (4)
O14—Cr3—O2W	86.41 (10)	Cl4'—C10—Cl3'	108.1 (4)
O10—Cr3—O2W	86.07 (10)	Cl5'—C10—Cl3'	106.4 (5)
O15—Cr3—O12	93.18 (9)	Cl14—C10—Cl3'	129.9 (5)
O8—Cr3—O12	86.73 (9)	O12—C11—O11	128.9 (3)
O14—Cr3—O12	94.66 (9)	O12—C11—C12	117.1 (3)
O10—Cr3—O12	172.32 (10)	O11—C11—C12	113.9 (3)
O2W—Cr3—O12	86.35 (10)	C11—C12—Cl17	112.9 (2)

supplementary materials

C1—O1—Cr1	130.9 (2)	C11—C12—C118	110.7 (2)
C3—O3—Cr1	132.3 (2)	C117—C12—C118	110.02 (18)
C3—O4—Cr2	125.6 (2)	C11—C12—C116	104.2 (2)
C5—O5—Cr1	126.5 (2)	C117—C12—C116	109.59 (19)
C5—O6—Cr2	129.5 (2)	C118—C12—C116	109.25 (18)
C7—O7—Cr1	126.0 (2)	O13—C13—O14	129.3 (3)
C7—O8—Cr3	131.6 (2)	O13—C13—C14	115.9 (3)
C9—O9—Cr1	132.6 (2)	O14—C13—C14	114.8 (3)
C9—O10—Cr3	126.0 (2)	C13—C14—C119	109.4 (2)
C11—O11—Cr2	131.5 (2)	C13—C14—C120	111.2 (2)
C11—O12—Cr3	125.8 (2)	C119—C14—C120	109.58 (19)
C13—O13—Cr2	127.4 (2)	C13—C14—C121	106.9 (2)
C13—O14—Cr3	131.1 (2)	C119—C14—C121	110.30 (19)
Cr2—O15—Cr3	120.38 (11)	C120—C14—C121	109.49 (19)
Cr2—O15—Cr1	119.25 (12)	N1—C15—C16	179.3 (5)
Cr3—O15—Cr1	120.35 (11)	C15—C16—H16A	109.5
Cr2—O1W—H1W1	112 (3)	C15—C16—H16B	109.5
Cr2—O1W—H1W2	122 (3)	H16A—C16—H16B	109.5
H1W1—O1W—H1W2	108.1 (17)	C15—C16—H16C	109.5
Cr3—O2W—H2W1	126 (3)	H16A—C16—H16C	109.5
Cr3—O2W—H2W2	120 (3)	H16B—C16—H16C	109.5
H2W1—O2W—H2W2	110.6 (18)	N2—C17—C18	178.2 (6)
O2—C1—O1	131.0 (3)	C17—C18—H18A	109.5
O2—C1—C2	117.9 (3)	C17—C18—H18B	109.5
O1—C1—C2	111.2 (3)	H18A—C18—H18B	109.5
C1—C2—C12	112.6 (2)	C17—C18—H18C	109.5
C1—C2—C13	108.5 (2)	H18A—C18—H18C	109.5
C12—C2—C13	109.03 (18)	H18B—C18—H18C	109.5
C1—C2—C11	108.6 (2)	N3—C19—C20	178.0 (9)
C12—C2—C11	109.29 (19)	C19—C20—H20A	109.5
C13—C2—C11	108.81 (18)	C19—C20—H20B	109.5
O3—C3—O4	128.4 (3)	H20A—C20—H20B	109.5
O3—C3—C4	115.3 (3)	C19—C20—H20C	109.5
O4—C3—C4	116.3 (3)	H20A—C20—H20C	109.5
C3—C4—C14	112.4 (2)	H20B—C20—H20C	109.5
O3—Cr1—O1—C1	62.2 (3)	O7—Cr1—O15—Cr3	56.08 (14)
O9—Cr1—O1—C1	-119.5 (3)	Cr1—O1—C1—O2	-17.1 (6)
O5—Cr1—O1—C1	-32.6 (3)	Cr1—O1—C1—C2	162.3 (2)
O7—Cr1—O1—C1	149.5 (3)	O2—C1—C2—C12	-3.5 (4)
O15—Cr1—O3—C3	-7.9 (3)	O1—C1—C2—C12	177.0 (2)
O1—Cr1—O3—C3	170.8 (3)	O2—C1—C2—C13	-124.3 (3)
O5—Cr1—O3—C3	-99.7 (3)	O1—C1—C2—C13	56.2 (3)
O7—Cr1—O3—C3	83.6 (3)	O2—C1—C2—C11	117.6 (3)
O15—Cr2—O4—C3	-39.5 (3)	O1—C1—C2—C11	-61.9 (3)
O11—Cr2—O4—C3	-133.4 (3)	Cr1—O3—C3—O4	28.9 (5)
O6—Cr2—O4—C3	55.3 (3)	Cr1—O3—C3—C4	-148.2 (2)
O1W—Cr2—O4—C3	141.2 (3)	Cr2—O4—C3—O3	1.6 (5)
O15—Cr1—O5—C5	-40.0 (3)	Cr2—O4—C3—C4	178.7 (2)
O1—Cr1—O5—C5	142.1 (3)	O3—C3—C4—C14	-161.7 (3)

O3—Cr1—O5—C5	53.7 (3)	O4—C3—C4—C14	20.9 (4)
O9—Cr1—O5—C5	-134.8 (3)	O3—C3—C4—C16	-38.2 (4)
O15—Cr2—O6—C5	-10.7 (3)	O4—C3—C4—C16	144.4 (3)
O13—Cr2—O6—C5	82.2 (3)	O3—C3—C4—C15	79.6 (3)
O4—Cr2—O6—C5	-104.8 (3)	O4—C3—C4—C15	-97.8 (3)
O1W—Cr2—O6—C5	168.9 (3)	Cr2—O6—C5—O5	34.3 (5)
O15—Cr1—O7—C7	-38.4 (3)	Cr2—O6—C5—C6	-144.0 (3)
O1—Cr1—O7—C7	139.6 (3)	Cr1—O5—C5—O6	-1.4 (5)
O3—Cr1—O7—C7	-131.9 (3)	Cr1—O5—C5—C6	176.9 (2)
O9—Cr1—O7—C7	56.5 (3)	O6—C5—C6—C19	-133.0 (3)
O15—Cr3—O8—C7	-5.6 (3)	O5—C5—C6—C19	48.5 (4)
O10—Cr3—O8—C7	-100.0 (3)	O6—C5—C6—C17	-9.9 (4)
O2W—Cr3—O8—C7	174.0 (3)	O5—C5—C6—C17	171.6 (3)
O12—Cr3—O8—C7	87.4 (3)	O6—C5—C6—C18	107.3 (3)
O15—Cr1—O9—C9	-7.1 (3)	O5—C5—C6—C18	-71.2 (3)
O1—Cr1—O9—C9	174.1 (3)	Cr1—O7—C7—O8	1.2 (5)
O5—Cr1—O9—C9	84.1 (3)	Cr1—O7—C7—C8	-179.7 (2)
O7—Cr1—O9—C9	-98.9 (3)	Cr3—O8—C7—O7	27.7 (5)
O15—Cr3—O10—C9	-40.5 (3)	Cr3—O8—C7—C8	-151.5 (2)
O8—Cr3—O10—C9	53.1 (3)	O7—C7—C8—C110	9.9 (4)
O14—Cr3—O10—C9	-134.1 (3)	O8—C7—C8—C110	-170.8 (2)
O2W—Cr3—O10—C9	139.2 (3)	O7—C7—C8—C112	132.2 (3)
O15—Cr2—O11—C11	-7.5 (3)	O8—C7—C8—C112	-48.5 (3)
O13—Cr2—O11—C11	-100.9 (3)	O7—C7—C8—C111	-109.4 (3)
O4—Cr2—O11—C11	86.1 (3)	O8—C7—C8—C111	69.9 (3)
O1W—Cr2—O11—C11	172.8 (3)	Cr1—O9—C9—O10	22.5 (5)
O15—Cr3—O12—C11	-35.3 (3)	Cr1—O9—C9—C10	-158.9 (2)
O8—Cr3—O12—C11	-128.6 (3)	Cr3—O10—C9—O9	7.7 (5)
O14—Cr3—O12—C11	58.9 (3)	Cr3—O10—C9—C10	-170.8 (2)
O2W—Cr3—O12—C11	145.0 (3)	O9—C9—C10—C14'	36.3 (7)
O15—Cr2—O13—C13	-34.4 (3)	O10—C9—C10—C14'	-145.0 (7)
O11—Cr2—O13—C13	59.8 (3)	O9—C9—C10—C113	-60.9 (4)
O6—Cr2—O13—C13	-128.7 (3)	O10—C9—C10—C113	117.9 (3)
O1W—Cr2—O13—C13	144.9 (3)	O9—C9—C10—C115	174.7 (4)
O15—Cr3—O14—C13	-2.0 (3)	O10—C9—C10—C115	-6.6 (5)
O10—Cr3—O14—C13	92.1 (3)	O9—C9—C10—C15'	166.5 (5)
O2W—Cr3—O14—C13	178.4 (3)	O10—C9—C10—C15'	-14.7 (6)
O12—Cr3—O14—C13	-95.5 (3)	O9—C9—C10—C114	56.3 (3)
O11—Cr2—O15—Cr3	-39.09 (14)	O10—C9—C10—C114	-124.9 (3)
O13—Cr2—O15—Cr3	53.69 (14)	O9—C9—C10—C13'	-80.5 (6)
O4—Cr2—O15—Cr3	-126.83 (13)	O10—C9—C10—C13'	98.2 (6)
O6—Cr2—O15—Cr3	138.90 (13)	Cr3—O12—C11—O11	-3.2 (5)
O11—Cr2—O15—Cr1	142.66 (13)	Cr3—O12—C11—C12	173.2 (2)
O13—Cr2—O15—Cr1	-124.55 (13)	Cr2—O11—C11—O12	31.1 (5)
O4—Cr2—O15—Cr1	54.92 (13)	Cr2—O11—C11—C12	-145.3 (2)
O6—Cr2—O15—Cr1	-39.35 (14)	O12—C11—C12—C117	23.3 (4)
O8—Cr3—O15—Cr2	140.83 (13)	O11—C11—C12—C117	-159.8 (2)
O14—Cr3—O15—Cr2	-40.99 (14)	O12—C11—C12—C118	147.1 (2)
O10—Cr3—O15—Cr2	-127.36 (13)	O11—C11—C12—C118	-36.0 (3)

supplementary materials

O12—Cr3—O15—Cr2	53.92 (13)	O12—C11—C12—C116	-95.5 (3)
O8—Cr3—O15—Cr1	-40.94 (14)	O11—C11—C12—C116	81.4 (3)
O14—Cr3—O15—Cr1	137.24 (13)	Cr2—O13—C13—O14	-1.5 (5)
O10—Cr3—O15—Cr1	50.87 (14)	Cr2—O13—C13—C14	176.2 (2)
O12—Cr3—O15—Cr1	-127.86 (13)	Cr3—O14—C13—O13	25.1 (5)
O3—Cr1—O15—Cr2	-38.30 (14)	Cr3—O14—C13—C14	-152.6 (2)
O9—Cr1—O15—Cr2	143.55 (13)	O13—C13—C14—C119	131.3 (3)
O5—Cr1—O15—Cr2	56.58 (14)	O14—C13—C14—C119	-50.7 (3)
O7—Cr1—O15—Cr2	-125.67 (13)	O13—C13—C14—C120	10.2 (4)
O3—Cr1—O15—Cr3	143.45 (13)	O14—C13—C14—C120	-171.8 (2)
O9—Cr1—O15—Cr3	-34.69 (14)	O13—C13—C14—C121	-109.3 (3)
O5—Cr1—O15—Cr3	-121.66 (14)	O14—C13—C14—C121	68.7 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1W—H1W1 \cdots N1	0.84 (1)	1.97 (2)	2.791 (4)	166 (5)
O1W—H1W2 \cdots N2	0.85 (1)	1.97 (1)	2.810 (5)	173 (4)
O2W—H2W1 \cdots N3	0.85 (1)	1.88 (1)	2.719 (5)	171 (4)
O2W—H2W2 \cdots O2 ⁱ	0.83 (1)	1.81 (1)	2.613 (3)	165 (4)

Symmetry codes: (i) $-x+3/2, y-1/2, -z+3/2$.

Fig. 1

