

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(Z)-6-{2-[(E)-2,4-Dihydroxybenzylideneamino]phenylaminomethylene}-3-hydroxycyclohexa-2,4-dienone toluene solvate

Hoong-Kun Fun,^{a*} Reza Kia,^{a‡} Valiollah Mirkhani^{b§} and Hasan Zargoshi^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bChemistry Department, University of Isfahan, Isfahan 81746-73441, Iran

Correspondence e-mail: hkfun@usm.my

Received 7 August 2008; accepted 14 August 2008


Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.059; wR factor = 0.164; data-to-parameter ratio = 20.8.

The bis-Schiff base title compound, $\text{C}_{20}\text{H}_{16}\text{N}_2\text{O}_4 \cdot \text{C}_7\text{H}_8$, crystallized as a toluene solvate. In the solid state, it is present as its prototropic tautomer formed by transfer of one of the *ortho*-hydroxyl H atoms. The proton transfer is accompanied by a shift of electron pairs, as is evident from the observed C—O and C—N bond distances of 1.305 (2) and 1.315 (2) Å, which are largely consistent with C=O and C—N distances. The actual molecule present in the solid state is thus the charge-neutral β -keto amine, with a small contribution of its zwitterionic valence tautomer *via* partial delocalization of electron pairs along the N—C—C—O atom chain. The dihedral angles between the central benzene ring and the two outer benzene rings of the Schiff base are 51.99 (8) and 12.95 (9)°. Intramolecular O—H...N and N—H...O hydrogen bonds generate $S(6)$ ring motifs, whereas intramolecular N—H...N hydrogen bonds generate $S(5)$ ring motifs. In the crystal structure, O—H...O hydrogen bonds and weak C—H...O interactions link the molecules into one-dimensional zigzag chains along the b axis; these chains are further stacked by O—H...O and weak C—H...O interactions along the c axis, forming two-dimensional extended networks parallel to the bc plane. In addition, the crystal structure is further stabilized by weak C—H... π and π - π interactions.

Related literature

For bond-length data, see: Allen *et al.* (1987). For details of hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see, for example: Cakir *et al.* (2002); Eltayeb *et al.*

(2007*a,b*); Karabiyik *et al.* (2007); Fun, Kargar & Kia (2008); Fun, Kia & Kargar (2008); Fun, Mirkhani *et al.* (2008*a,b*). For background on applications of Schiff base ligands, see, for example: Hajioudis *et al.* (1987); Granovski *et al.* (1993); Dao *et al.* (2000); Shahrokhian *et al.* (2000); Eltayeb & Ahmed (2005*a,b*); Fakhari *et al.* (2005); Karthikeyan *et al.* (2006); Sriram *et al.* (2006). For related literature, see: Fun & Kia (2008).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{16}\text{N}_2\text{O}_4 \cdot \text{C}_7\text{H}_8$
 $M_r = 440.48$
 Monoclinic, $P2_1/c$
 $a = 11.9753$ (3) Å
 $b = 18.8539$ (5) Å
 $c = 9.9240$ (2) Å
 $\beta = 108.819$ (1)°

$V = 2120.87$ (9) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 100.0$ (1) K
 $0.25 \times 0.13 \times 0.02$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.954$, $T_{\max} = 0.994$

24830 measured reflections
 6233 independent reflections
 4023 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.163$
 $S = 1.11$
 6233 reflections

299 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.76$ e Å⁻³
 $\Delta\rho_{\min} = -0.32$ e Å⁻³

Table 1

Selected centroid...centroid distances (Å).

$\text{Cg1} \cdots \text{Cg1}^i$	3.7867 (1)	$\text{Cg2} \cdots \text{Cg3}^{ii}$	4.5626 (3)
----------------------------------	------------	-------------------------------------	------------

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x, -y - \frac{1}{2}, z - \frac{3}{2}$. Cg1, Cg2, and Cg3 are the centroids of the C1–C6, C8–C13 and C15–C20 benzene rings, respectively.

‡ Additional correspondence author, e-mail: zsrkk@yahoo.com.

§ Additional correspondence author, e-mail: mirkhani@sci.ui.ac.ir.

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1O1 \cdots N1	0.94	1.83	2.6568 (17)	145
O3—H1O3 \cdots O2 ⁱ	0.96	1.64	2.5919 (18)	171
O4—H1O4 \cdots O3 ⁱⁱⁱ	0.90	1.87	2.7403 (16)	162
N2—H1N2 \cdots O2	0.88	1.84	2.5954 (18)	143
N2—H1N2 \cdots N1	0.88	2.37	2.7245 (19)	104
C16—H16A \cdots O1 ^{iv}	0.95	2.55	3.439 (2)	157
C17—H17A \cdots O4 ^v	0.95	2.51	3.381 (2)	152
C11—H11A \cdots Cg4 ^{vi}	0.95	2.97	3.619 (2)	126

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (iii) $-x+1, y-\frac{1}{2}, -z+\frac{3}{2}$; (iv) $x, -y+\frac{1}{2}, z-\frac{1}{2}$; (v) $-x+1, -y, -z+1$; (vi) $-x, y-\frac{1}{2}, -z+\frac{1}{2}$. Cg4 is the centroid of the C21–C26 benzene ring.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2003).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. RK thanks Universiti Sains Malaysia for a post-doctoral research fellowship. VM and HZ thank the University of Isfahan for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2135).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cakir, O., Elerman, Y. & Elmali, A. (2002). *Anal. Sci.*, **18**, 377–377.
- Dao, V.-T., Gaspard, C., Mayer, M., Werner, G. H., Nguyen, S. N. & Michelot, R. J. (2000). *Eur. J. Med. Chem.* **35**, 805–813.
- Eltayeb, N. E. & Ahmed, T. A. (2005a). *J. Sci. Tech.* **6**, 51–59.
- Eltayeb, N. E. & Ahmed, T. A. (2005b). *Sudan J. Basic Sci.* **7**, 97–108.
- Eltayeb, N. E., Teoh, S. G., Chantrapromma, S., Fun, H.-K. & Ibrahim, K. (2007a). *Acta Cryst. E* **63**, o3094–o3095.
- Eltayeb, N. E., Teoh, S. G., Chantrapromma, S., Fun, H.-K. & Ibrahim, K. (2007b). *Acta Cryst. E* **63**, o3234–o3235.
- Fakhari, A. R., Khorrami, A. R. & Naeimi, H. (2005). *Talanta*, **66**, 813–817.
- Fun, H.-K., Kargar, H. & Kia, R. (2008). *Acta Cryst. E* **64**, o1308.
- Fun, H.-K. & Kia, R. (2008). *Acta Cryst. E* **64**, o1657–o1658.
- Fun, H.-K., Kia, R. & Kargar, H. (2008). *Acta Cryst. E* **64**, o1335.
- Fun, H.-K., Mirkhani, V., Kia, R. & Rostami Vartooni, A. (2008a). *Acta Cryst. E* **64**, o1374–o1375.
- Fun, H.-K., Mirkhani, V., Kia, R. & Rostami Vartooni, A. (2008b). *Acta Cryst. E* **64**, o1471.
- Granovski, A. D., Nivorozhkin, A. L. & Minkin, V. I. (1993). *Coord. Chem. Rev.* **126**, 1–69.
- Hajioudis, E., Vitterakis, M. & Mustakali-Mavridis, I. (1987). *Tetrahedron*, **43**, 1345–1351.
- Karabiyik, H., Guzel, B., Aygun, M., Boga, G. & Buyukgungor, O. (2007). *Acta Cryst. C* **63**, o215–o218.
- Karthikeyan, M. S., Prasad, D. J., Poojary, B., Bhat, K. S., Holla, B. S. & Kumari, N. S. (2006). *Bioorg. Med. Chem.* **14**, 7482–7489.
- Shahrokhian, S., Amini, M. K., Kia, R. & Tangestaninejad, S. (2000). *Anal. Chem.* **72**, 956–962.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Sriram, D., Yogeewari, P., Myneedu, N. S. & Saraswat, V. (2006). *Bioorg. Med. Chem. Lett.* **16**, 2127–2129.

supporting information

Acta Cryst. (2008). E64, o1790–o1791 [doi:10.1107/S1600536808026305]

(Z)-6-{2-[(E)-2,4-Dihydroxybenzylideneamino]phenylaminomethylene}-3-hydroxycyclohexa-2,4-dienone toluene solvate

Hoong-Kun Fun, Reza Kia, Valiollah Mirkhani and Hasan Zargoshi

S1. Comment

Schiff bases have received much attention because of their potential applications with some of these compounds exhibiting various pharmacological activities, as noted by their anticancer (Dao *et al.*, 2000), anti-HIV (Sriram *et al.*, 2006), antibacterial and antifungal (Karthikeyan *et al.*, 2006) properties. Although numerous transition-metal complexes of Schiff bases have been structurally characterized (Granovski *et al.*, 1993), relatively few free Schiff bases have been similarly characterized. *N*-substituted salicylaldehydes show photochromism and thermochromism in the solid state. These effects are produced by intramolecular proton transfer associated with a change in the π -electron configuration (Hajioudis *et al.* 1987). In addition, some of them may be used as analytical reagents for the determination of trace elements (Eltayeb & Ahmed, 2005*a,b*) such as nickel in some natural food products (Fakhari *et al.*, 2005) or biologically important species (Shahrokhian *et al.*, 2000). As part of a general study of tetradenate and bidentate Schiff bases (Fun, Kargar & Kia 2008; Fun, Kia & Kargar 2008; Fun, Mirkhani *et al.*, 2008*a,b*), we determined the structure of the title compound.

The title compound was synthesized from *o*-phenylenediamine by reaction with two equivalents of 2,4-dihydroxybenzaldehyde, and the expected reaction product would thus have been the bis-Schiff base 4-((*E*)-(2-((*E*)-2-hydroxybenzylideneamino)phenylimino)methyl) benzene-1,3-diol. The actual molecule obtained in the solid state is however its prototropic tautomer formed by transfer of one of the ortho-hydroxyl protons onto the adjacent imine unit. The proton transfer is accompanied by a shift of electron pairs as is evident from the observed C20–O2 and C14–N2 bond distances of 1.305 (2) and 1.315 (2) Å, which are consistent with C=O and C–N distances (Allen *et al.*, 1987), respectively. The formation of a C=O keto group rather than a C–O phenolate is also obvious by comparison with the other three phenol C–OH groups in the structure, which are about 0.05 Å longer than C20–O2. The actual molecule present in the solid state is thus the charge neutral β -keto amine (*Z*)-3-hydroxy-6-((2-((*E*)-2-hydroxybenzylideneamino)phenylamino)-methylene)cyclohexa-2,4-dienone (top isomer in Fig. 4), with a small contribution of its zwitter-ionic valence tautomer via partial delocalization of electron pairs along the atom chain N2–C14–C15–C20–O2 (bottom tautomer in Fig. 4). The other imine group did not undergo proton transfer and is present in its original Schiff base state. Both the imine as well as the amine units are stabilized by strong O–H \cdots N and N–H \cdots O hydrogen bonds (Table 2) that generate *S*(6) ring motifs whereas the intramolecular N–H \cdots N hydrogen bond between the amine and imine (Table 2) exhibits an *S*(5) ring motif (Bernstein *et al.*, 1995). Bond lengths and angles are in normal ranges (Allen *et al.*, 1987) and comparable to those in related structures (Eltayeb *et al.*, 2007*a,b*; Cakir *et al.* 2002; Karabiyik *et al.*, 2007). The C8–C13 phenyl ring makes a dihedral angle of 51.99 (8)° with the dihydroxyphenyl ring (C1–C6/O1/O3) and 12.95 (9)° with the keto-hydroxyphenyl ring (C15–C20/O2/O4). In the crystal packing (Fig. 2), additional O–H \cdots O hydrogen bonds and weak C–H \cdots O interactions (Table 2) link the molecules into one dimensional zigzag extended chains along the *b* axis and these chains


are further stacked (Fig. 2 & 3) along the *c* axis thus forming two-dimensional extended networks parallel to the *bc* plane. The crystal is further stabilized by weak C—H $\cdots\pi$ interactions (Table 2). The short distance between the centroids of the six-membered rings prove an existence of $\pi\cdots\pi$ interactions (Table 1).

S2. Experimental


The title compound was synthesized by adding 2,4-dihydroxybenzaldehyde (0.552 g, 4 mmol) to a solution of *o*-phenylenediamine (0.216 g, 2 mmol) in ethanol (20 ml). The mixture was refluxed with stirring for half an hour. The resultant yellow solution was filtered. Yellow single crystals of the title compound suitable for *X*-ray structure determination were recrystallized from a mixture of THF/toluene (2/1) by slow evaporation of the solvents at room temperature over several days.

S3. Refinement

Hydroxyl and amine/imine H atoms were located from the difference Fourier map and refined as riding on the parent atoms with isotropic refinement of the displacement parameters. The remaining H atoms were geometrically located and refined as riding model. A rotating group model was used for the methyl groups.

**Figure 1**

The asymmetric unit of (I), showing 50% probability displacement ellipsoids and the atomic numbering. Intramolecular hydrogen bonds are drawn as dashed lines.

**Figure 2**

The crystal packing of (I), viewed down the *c* axis, showing the molecular chains along the *b* axis and stacking of these chains along the *c*-axis. Hydrogen bonds are drawn as dashed lines. The toluene molecules were omitted for clarity.





Figure 3

The crystal packing of (I), showing 1-D extended chains along the *c* axis. The toluene molecules were omitted for clarity.

**Figure 4**

The charge neutral β -keto amine (main component) form and the valence tautomer via partial delocalization of electron pairs along the N—C—C—C—O atom chain (small contribution) in the title compound.

(Z)-6-{2-[(E)-2,4-Dihydroxybenzylideneamino]phenylaminomethylene}- 3-hydroxycyclohexa-2,4-dienone toluene solvate

Crystal data

$C_{20}H_{16}N_2O_4 \cdot C_7H_8$

$M_r = 440.48$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2ybc$

$a = 11.9753$ (3) Å

$b = 18.8539$ (5) Å

$c = 9.9240$ (2) Å

$\beta = 108.819$ (1)°

$V = 2120.87$ (9) Å³

$Z = 4$

$F(000) = 928$

$D_x = 1.380$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 5653 reflections

$\theta = 2.4\text{--}29.6^\circ$
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 100\text{ K}$

Plate, yellow
 $0.25 \times 0.13 \times 0.02\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.954$, $T_{\max} = 0.994$

24830 measured reflections
 6233 independent reflections
 4023 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$
 $\theta_{\max} = 30.2^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -14 \rightarrow 16$
 $k = -20 \rightarrow 26$
 $l = -13 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.058$
 $wR(F^2) = 0.163$
 $S = 1.11$
 6233 reflections
 299 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0708P)^2 + 0.4015P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.76\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32\text{ e \AA}^{-3}$

Special details

Experimental. The low-temperature data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.18268 (12)	0.42566 (7)	0.14400 (14)	0.0168 (3)
O3	0.50774 (11)	0.63459 (6)	0.61784 (12)	0.0214 (3)
H1O3	0.5724	0.6608	0.6044	0.032*
O1	0.23309 (11)	0.45373 (6)	0.41957 (12)	0.0225 (3)
H1O1	0.2053	0.4270	0.3349	0.034*
O4	0.48126 (11)	0.07727 (6)	0.62444 (12)	0.0217 (3)
H1O4	0.4999	0.0995	0.7096	0.033*
O2	0.31282 (10)	0.29118 (6)	0.39178 (12)	0.0187 (3)
N2	0.17826 (12)	0.28136 (7)	0.12862 (14)	0.0170 (3)
H1N2	0.2152	0.3043	0.2079	0.025*
C8	0.11363 (14)	0.39346 (9)	0.01527 (17)	0.0165 (3)
C6	0.31241 (14)	0.51993 (9)	0.26313 (17)	0.0161 (3)

C15	0.27803 (14)	0.18211 (9)	0.26242 (17)	0.0165 (3)
C13	0.10818 (14)	0.31889 (9)	0.00863 (17)	0.0164 (3)
C16	0.30131 (15)	0.10819 (9)	0.26262 (18)	0.0194 (4)
H16A	0.2684	0.0818	0.1775	0.023*
C7	0.24537 (14)	0.48061 (9)	0.13868 (17)	0.0165 (3)
H7A	0.2475	0.4959	0.0483	0.020*
C9	0.04597 (15)	0.43207 (9)	-0.10182 (17)	0.0193 (4)
H9A	0.0487	0.4824	-0.0988	0.023*
C19	0.39752 (15)	0.18616 (9)	0.51268 (17)	0.0177 (3)
H19A	0.4305	0.2113	0.5994	0.021*
C5	0.38949 (15)	0.57342 (8)	0.25065 (17)	0.0176 (4)
H5A	0.3961	0.5834	0.1597	0.021*
C1	0.30495 (14)	0.50578 (8)	0.39969 (17)	0.0163 (3)
C17	0.36973 (15)	0.07431 (9)	0.38208 (18)	0.0198 (4)
H17A	0.3853	0.0250	0.3804	0.024*
C18	0.41709 (15)	0.11407 (9)	0.50847 (17)	0.0176 (3)
C2	0.36978 (14)	0.54490 (9)	0.51645 (17)	0.0175 (4)
H2A	0.3627	0.5358	0.6075	0.021*
C12	0.03656 (15)	0.28533 (9)	-0.11326 (17)	0.0204 (4)
H12A	0.0333	0.2350	-0.1173	0.024*
C20	0.32949 (14)	0.22283 (9)	0.39033 (17)	0.0164 (3)
C4	0.45602 (15)	0.61205 (9)	0.36648 (17)	0.0185 (4)
H4A	0.5083	0.6479	0.3559	0.022*
C14	0.20597 (15)	0.21362 (9)	0.13726 (17)	0.0176 (4)
H14A	0.1756	0.1849	0.0548	0.021*
C10	-0.02551 (15)	0.39818 (10)	-0.22315 (18)	0.0225 (4)
H10A	-0.0714	0.4253	-0.3024	0.027*
C3	0.44526 (14)	0.59760 (9)	0.49988 (17)	0.0168 (3)
C11	-0.02984 (15)	0.32460 (10)	-0.22838 (18)	0.0231 (4)
H11A	-0.0786	0.3013	-0.3114	0.028*
C23	0.30310 (18)	0.71109 (11)	0.7808 (2)	0.0329 (5)
H23A	0.3583	0.7384	0.7523	0.040*
C25	0.2145 (2)	0.67854 (11)	0.9559 (2)	0.0352 (5)
H25A	0.2093	0.6830	1.0490	0.042*
C26	0.13983 (19)	0.63297 (10)	0.8595 (2)	0.0314 (5)
H26A	0.0822	0.6073	0.8867	0.038*
C22	0.22868 (18)	0.66449 (10)	0.6864 (2)	0.0293 (4)
H22A	0.2339	0.6604	0.5932	0.035*
C21	0.14696 (17)	0.62366 (10)	0.7224 (2)	0.0292 (4)
C24	0.29687 (18)	0.71778 (11)	0.9173 (2)	0.0350 (5)
H24A	0.3487	0.7490	0.9837	0.042*
C27	0.0688 (2)	0.57353 (12)	0.6196 (2)	0.0402 (5)
H27A	0.1166	0.5365	0.5954	0.060*
H27B	0.0242	0.5992	0.5331	0.060*
H27C	0.0139	0.5517	0.6622	0.060*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0194 (7)	0.0153 (7)	0.0144 (7)	0.0009 (6)	0.0036 (6)	-0.0015 (5)
O3	0.0251 (7)	0.0191 (6)	0.0189 (6)	-0.0058 (5)	0.0056 (5)	-0.0025 (5)
O1	0.0294 (7)	0.0215 (7)	0.0167 (6)	-0.0103 (5)	0.0076 (5)	-0.0025 (5)
O4	0.0273 (7)	0.0183 (6)	0.0174 (6)	0.0051 (5)	0.0043 (5)	0.0029 (5)
O2	0.0224 (6)	0.0131 (6)	0.0187 (6)	-0.0011 (5)	0.0039 (5)	-0.0009 (5)
N2	0.0183 (7)	0.0169 (7)	0.0137 (7)	-0.0029 (6)	0.0025 (6)	-0.0007 (5)
C8	0.0163 (8)	0.0201 (9)	0.0131 (8)	-0.0010 (7)	0.0048 (6)	-0.0024 (6)
C6	0.0180 (8)	0.0144 (8)	0.0150 (8)	0.0017 (6)	0.0040 (6)	-0.0004 (6)
C15	0.0189 (8)	0.0162 (8)	0.0163 (8)	-0.0011 (7)	0.0082 (7)	-0.0006 (6)
C13	0.0152 (8)	0.0170 (8)	0.0172 (8)	0.0002 (6)	0.0053 (7)	0.0015 (6)
C16	0.0261 (9)	0.0173 (9)	0.0168 (8)	-0.0008 (7)	0.0099 (7)	-0.0025 (7)
C7	0.0188 (8)	0.0165 (8)	0.0139 (8)	0.0033 (7)	0.0047 (7)	0.0005 (6)
C9	0.0218 (9)	0.0181 (9)	0.0169 (8)	0.0033 (7)	0.0047 (7)	0.0013 (7)
C19	0.0196 (8)	0.0176 (9)	0.0151 (8)	-0.0014 (7)	0.0044 (7)	-0.0009 (6)
C5	0.0212 (9)	0.0157 (8)	0.0168 (8)	0.0027 (7)	0.0073 (7)	0.0031 (6)
C1	0.0180 (8)	0.0133 (8)	0.0174 (8)	0.0012 (6)	0.0055 (7)	0.0008 (6)
C17	0.0259 (9)	0.0148 (8)	0.0214 (9)	0.0022 (7)	0.0115 (7)	0.0005 (7)
C18	0.0179 (8)	0.0186 (9)	0.0177 (8)	0.0022 (7)	0.0077 (7)	0.0034 (7)
C2	0.0206 (9)	0.0162 (8)	0.0156 (8)	0.0005 (7)	0.0059 (7)	-0.0002 (6)
C12	0.0195 (9)	0.0194 (9)	0.0205 (9)	-0.0045 (7)	0.0041 (7)	-0.0035 (7)
C20	0.0163 (8)	0.0161 (8)	0.0182 (8)	-0.0010 (6)	0.0074 (7)	0.0003 (6)
C4	0.0202 (9)	0.0143 (8)	0.0207 (9)	-0.0003 (7)	0.0062 (7)	0.0011 (7)
C14	0.0214 (9)	0.0156 (8)	0.0167 (8)	-0.0025 (7)	0.0074 (7)	-0.0007 (6)
C10	0.0197 (9)	0.0285 (10)	0.0159 (8)	0.0037 (7)	0.0012 (7)	0.0014 (7)
C3	0.0179 (8)	0.0142 (8)	0.0164 (8)	0.0008 (6)	0.0028 (7)	-0.0034 (6)
C11	0.0197 (9)	0.0291 (10)	0.0168 (9)	-0.0014 (8)	0.0009 (7)	-0.0037 (7)
C23	0.0296 (11)	0.0343 (12)	0.0362 (12)	0.0008 (9)	0.0124 (9)	0.0017 (9)
C25	0.0458 (13)	0.0309 (11)	0.0313 (11)	0.0098 (10)	0.0158 (10)	0.0044 (9)
C26	0.0341 (11)	0.0258 (11)	0.0402 (12)	0.0053 (9)	0.0201 (10)	0.0106 (9)
C22	0.0307 (11)	0.0280 (10)	0.0316 (11)	0.0033 (9)	0.0134 (9)	0.0029 (8)
C21	0.0292 (10)	0.0275 (10)	0.0311 (11)	0.0057 (8)	0.0101 (9)	0.0055 (8)
C24	0.0330 (11)	0.0363 (12)	0.0336 (11)	0.0063 (9)	0.0076 (9)	-0.0001 (9)
C27	0.0429 (13)	0.0380 (13)	0.0415 (13)	-0.0042 (10)	0.0161 (11)	0.0008 (10)

Geometric parameters (\AA , $^\circ$)

N1—C7	1.290 (2)	C5—H5A	0.9500
N1—C8	1.415 (2)	C1—C2	1.382 (2)
O3—C3	1.3608 (19)	C17—C18	1.413 (2)
O3—H1O3	0.9628	C17—H17A	0.9500
O1—C1	1.3608 (19)	C2—C3	1.388 (2)
O1—H1O1	0.9431	C2—H2A	0.9500
O4—C18	1.3513 (19)	C12—C11	1.378 (2)
O4—H1O4	0.9048	C12—H12A	0.9500
O2—C20	1.3048 (19)	C4—C3	1.398 (2)

N2—C14	1.315 (2)	C4—H4A	0.9500
N2—C13	1.406 (2)	C14—H14A	0.9500
N2—H1N2	0.8816	C10—C11	1.389 (3)
C8—C9	1.389 (2)	C10—H10A	0.9500
C8—C13	1.408 (2)	C11—H11A	0.9500
C6—C5	1.399 (2)	C23—C22	1.380 (3)
C6—C1	1.412 (2)	C23—C24	1.387 (3)
C6—C7	1.442 (2)	C23—H23A	0.9500
C15—C14	1.396 (2)	C25—C26	1.379 (3)
C15—C16	1.421 (2)	C25—C24	1.382 (3)
C15—C20	1.441 (2)	C25—H25A	0.9500
C13—C12	1.389 (2)	C26—C21	1.402 (3)
C16—C17	1.363 (2)	C26—H26A	0.9500
C16—H16A	0.9500	C22—C21	1.380 (3)
C7—H7A	0.9500	C22—H22A	0.9500
C9—C10	1.388 (2)	C21—C27	1.482 (3)
C9—H9A	0.9500	C24—H24A	0.9500
C19—C18	1.382 (2)	C27—H27A	0.9800
C19—C20	1.406 (2)	C27—H27B	0.9800
C19—H19A	0.9500	C27—H27C	0.9800
C5—C4	1.377 (2)		
Cg1...Cg1 ⁱ	3.7867 (1)	Cg2...Cg3 ⁱⁱ	4.5626 (3)
C7—N1—C8	119.05 (14)	C11—C12—C13	120.40 (16)
C3—O3—H1O3	112.8	C11—C12—H12A	119.8
C1—O1—H1O1	108.3	C13—C12—H12A	119.8
C18—O4—H1O4	117.3	O2—C20—C19	121.60 (15)
C14—N2—C13	127.78 (14)	O2—C20—C15	120.84 (15)
C14—N2—H1N2	112.0	C19—C20—C15	117.56 (15)
C13—N2—H1N2	120.0	C5—C4—C3	118.85 (15)
C9—C8—C13	118.59 (15)	C5—C4—H4A	120.6
C9—C8—N1	122.94 (15)	C3—C4—H4A	120.6
C13—C8—N1	118.38 (14)	N2—C14—C15	122.80 (15)
C5—C6—C1	117.97 (15)	N2—C14—H14A	118.6
C5—C6—C7	119.71 (14)	C15—C14—H14A	118.6
C1—C6—C7	122.32 (15)	C9—C10—C11	119.84 (16)
C14—C15—C16	118.90 (15)	C9—C10—H10A	120.1
C14—C15—C20	121.53 (15)	C11—C10—H10A	120.1
C16—C15—C20	119.57 (15)	O3—C3—C2	117.60 (14)
C12—C13—N2	122.70 (15)	O3—C3—C4	121.45 (15)
C12—C13—C8	120.12 (15)	C2—C3—C4	120.95 (15)
N2—C13—C8	117.18 (14)	C12—C11—C10	120.08 (16)
C17—C16—C15	121.56 (15)	C12—C11—H11A	120.0
C17—C16—H16A	119.2	C10—C11—H11A	120.0
C15—C16—H16A	119.2	C22—C23—C24	119.6 (2)
N1—C7—C6	123.24 (15)	C22—C23—H23A	120.2
N1—C7—H7A	118.4	C24—C23—H23A	120.2

C6—C7—H7A	118.4	C26—C25—C24	120.16 (19)
C10—C9—C8	120.97 (16)	C26—C25—H25A	119.9
C10—C9—H9A	119.5	C24—C25—H25A	119.9
C8—C9—H9A	119.5	C25—C26—C21	121.50 (19)
C18—C19—C20	120.93 (15)	C25—C26—H26A	119.3
C18—C19—H19A	119.5	C21—C26—H26A	119.3
C20—C19—H19A	119.5	C21—C22—C23	122.41 (19)
C4—C5—C6	121.86 (15)	C21—C22—H22A	118.8
C4—C5—H5A	119.1	C23—C22—H22A	118.8
C6—C5—H5A	119.1	C22—C21—C26	116.88 (19)
O1—C1—C2	118.28 (14)	C22—C21—C27	121.39 (18)
O1—C1—C6	120.95 (14)	C26—C21—C27	121.72 (19)
C2—C1—C6	120.77 (15)	C25—C24—C23	119.4 (2)
C16—C17—C18	118.67 (16)	C25—C24—H24A	120.3
C16—C17—H17A	120.7	C23—C24—H24A	120.3
C18—C17—H17A	120.7	C21—C27—H27A	109.5
O4—C18—C19	122.31 (15)	C21—C27—H27B	109.5
O4—C18—C17	116.02 (15)	H27A—C27—H27B	109.5
C19—C18—C17	121.66 (15)	C21—C27—H27C	109.5
C1—C2—C3	119.59 (15)	H27A—C27—H27C	109.5
C1—C2—H2A	120.2	H27B—C27—H27C	109.5
C3—C2—H2A	120.2		
C7—N1—C8—C9	-43.9 (2)	N2—C13—C12—C11	-179.57 (15)
C7—N1—C8—C13	139.83 (16)	C8—C13—C12—C11	-0.2 (2)
C14—N2—C13—C12	14.2 (3)	C18—C19—C20—O2	-178.20 (15)
C14—N2—C13—C8	-165.19 (16)	C18—C19—C20—C15	2.0 (2)
C9—C8—C13—C12	0.2 (2)	C14—C15—C20—O2	-2.0 (2)
N1—C8—C13—C12	176.67 (14)	C16—C15—C20—O2	177.67 (15)
C9—C8—C13—N2	179.64 (14)	C14—C15—C20—C19	177.82 (15)
N1—C8—C13—N2	-3.9 (2)	C16—C15—C20—C19	-2.5 (2)
C14—C15—C16—C17	-179.13 (15)	C6—C5—C4—C3	-0.5 (2)
C20—C15—C16—C17	1.2 (2)	C13—N2—C14—C15	179.10 (15)
C8—N1—C7—C6	175.76 (15)	C16—C15—C14—N2	178.73 (15)
C5—C6—C7—N1	172.13 (15)	C20—C15—C14—N2	-1.6 (2)
C1—C6—C7—N1	-6.8 (3)	C8—C9—C10—C11	-0.1 (3)
C13—C8—C9—C10	-0.1 (2)	C1—C2—C3—O3	-179.73 (14)
N1—C8—C9—C10	-176.39 (15)	C1—C2—C3—C4	0.5 (2)
C1—C6—C5—C4	-0.5 (2)	C5—C4—C3—O3	-179.24 (15)
C7—C6—C5—C4	-179.50 (15)	C5—C4—C3—C2	0.5 (2)
C5—C6—C1—O1	-178.99 (14)	C13—C12—C11—C10	0.0 (3)
C7—C6—C1—O1	0.0 (2)	C9—C10—C11—C12	0.1 (3)
C5—C6—C1—C2	1.6 (2)	C24—C25—C26—C21	1.7 (3)
C7—C6—C1—C2	-179.48 (15)	C24—C23—C22—C21	0.1 (3)
C15—C16—C17—C18	0.7 (2)	C23—C22—C21—C26	1.9 (3)
C20—C19—C18—O4	-179.47 (15)	C23—C22—C21—C27	-179.43 (19)
C20—C19—C18—C17	-0.1 (2)	C25—C26—C21—C22	-2.7 (3)
C16—C17—C18—O4	178.10 (14)	C25—C26—C21—C27	178.57 (19)

C16—C17—C18—C19	-1.3 (2)	C26—C25—C24—C23	0.4 (3)
O1—C1—C2—C3	178.99 (15)	C22—C23—C24—C25	-1.2 (3)
C6—C1—C2—C3	-1.6 (2)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, -y-1/2, z-3/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1—H1O1...N1	0.94	1.83	2.6568 (17)	145
O3—H1O3...O2 ⁱ	0.96	1.64	2.5919 (18)	171
O4—H1O4...O3 ⁱⁱⁱ	0.90	1.87	2.7403 (16)	162
N2—H1N2...O2	0.88	1.84	2.5954 (18)	143
N2—H1N2...N1	0.88	2.37	2.7245 (19)	104
C16—H16A...O1 ^{iv}	0.95	2.55	3.439 (2)	157
C17—H17A...O4 ^v	0.95	2.51	3.381 (2)	152
C11—H11A...Cg4 ^{vi}	0.95	2.97	3.619 (2)	126

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (iii) $-x+1, y-1/2, -z+3/2$; (iv) $x, -y+1/2, z-1/2$; (v) $-x+1, -y, -z+1$; (vi) $-x, y-1/2, -z+1/2$.