

catena-Poly[[bis(1-methylimidazole- κN^3)-zinc(II)]- μ -isophthalato- $\kappa^2 O^1 : O^3$]

Juan Zhao

College of Mechanical Engineering, Qingdao Technological University, Qingdao 266033, People's Republic of China
Correspondence e-mail: zhaojuanqd@163.com


Received 17 September 2008; accepted 24 September 2008

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.009$ Å; R factor = 0.068; wR factor = 0.207; data-to-parameter ratio = 13.5.

In the solid state, the title compound, $[Zn(C_8H_4O_4)(C_4H_6N_2)_2]_n$, exhibits the existence of polymeric zigzag chains extending along the a axis. Each Zn^{II} ion is coordinated by two N atoms [$Zn-N = 1.996$ (6) and 2.032 (5) Å] and two O atoms [$Zn-O = 1.930$ (4) and 1.976 (4) Å] in a distorted tetrahedral geometry. Weak C–H···O interactions contribute to the crystal packing stability.

Related literature

In the related zinc compound $[Zn(isophthalato)(1-H-imidazole)_2]$ (Yang *et al.*, 2002), the Zn^{II} ions also have a distorted tetrahedral environment.



Experimental

Crystal data

$[Zn(C_8H_4O_4)(C_4H_6N_2)_2]$	$V = 3454.8$ (12) Å ³
$M_r = 393.72$	$Z = 8$
Orthorhombic, $Pbca$	Mo $K\alpha$ radiation
$a = 9.6820$ (19) Å	$\mu = 1.45$ mm ⁻¹
$b = 13.224$ (3) Å	$T = 293$ (2) K
$c = 26.983$ (5) Å	$0.20 \times 0.10 \times 0.10$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer	2041 reflections with $I > 2\sigma(I)$
Absorption correction: none	$R_{int} = 0.035$
3091 measured reflections	3 standard reflections
2967 independent reflections	every 100 reflections
220 parameters	intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.068$	40 restraints
$wR(F^2) = 0.207$	H-atom parameters constrained
$S = 1.05$	$\Delta\rho_{\max} = 0.50$ e Å ⁻³
2967 reflections	$\Delta\rho_{\min} = -0.61$ e Å ⁻³
220 parameters	

Table 1
Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
C1–H1A···O3 ⁱ	0.96	2.55	3.423 (10)	150
C4–H4B···O3 ⁱ	0.93	2.31	3.150 (9)	150
C11–H11A···O2 ⁱⁱ	0.93	2.54	3.457 (8)	171

Symmetry codes: (i) $-x + \frac{3}{2}, y + \frac{1}{2}, z$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported by the National Natural Science Foundation of China (grant No. 20601015) and the Natural Science Foundation of Shandong Province (grant No. Y2006B12).


Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2451).

References

- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Yang, J., Zheng, S., Tao, J., Liu, G. & Chen, X. (2002). *Aust. J. Chem.* **55**, 741–744.

supporting information

Acta Cryst. (2008). E64, m1335 [doi:10.1107/S160053680803078X]



Juan Zhao

S1. Comment

In the title compound, (I) (Fig. 1), the zinc(II) centers are bridged by carboxylate groups of isophthalate ligands. Each Zn^{II} ion is coordinated by two N [Zn—N2 = 1.996 (6) Å, Zn—N4 = 2.032 (5) Å] and two O [Zn—O1 = 1.930 (4) Å, Zn—O4 = 1.976 (4) Å] atoms in a distorted tetrahedral geometry. All these values agree well with those observed in [Zn(isophthalato)(1-H-imidazole)₂] (Yang *et al.*, 2002). Each isophthalate dianion in (I) acts as a bidentate ligand to bridge two Zn^{II} atoms through two monodentate carboxylate groups, building a zigzag polymeric chain along the *a* axis. The metal–metal distance across each polymer backbone is 9.682 (7) Å.


In the crystal, weak C—H···O interactions contribute to the crystal packing stability. In the corresponding zinc compound [Zn(isophthalato)(1-H-imidazole)₂] (Yang *et al.*, 2002), the Zn^{II} ions have a distorted tetrahedral environment.

S2. Experimental

The reaction of ZnCl₂(0.68 g, 5 mmol) with isophthalic acid (0.83 g, 5 mmol) in an aqueous-alcohol (3:1) solution (40 ml) at 363 K for 30 minutes produced a blue solution, to which 1-methylimidazole (0.82 g, 10 mmol) was added. The reaction solution was kept at room temperature after stirring for an hour at 333 K. Colourless crystals were obtained after a few days.

S3. Refinement


H atoms were positioned geometrically (C—H = 0.93 or 0.96 Å) and allowed to ride on their parent atoms with $U_{\text{iso}}(\text{H})$ = 1.2 or 1.5 times $U_{\text{eq}}(\text{C})$.

**Figure 1**

A portion of the polymeric chain in (I) showing the atomic numbering and 30% probability displacement ellipsoids.

catena-Poly[[bis(1-methylimidazole- κ N³)zinc(II)]- μ -isophthalato- κ^2 O¹:O³]

Crystal data



$M_r = 393.72$

Orthorhombic, $Pbca$

Hall symbol: -P 2ac 2ab

$a = 9.6820$ (19) Å

$b = 13.224$ (3) Å

$c = 26.983$ (5) Å

$V = 3454.8$ (12) Å³

$Z = 8$

$F(000) = 1616$

$D_x = 1.514$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 10\text{--}14^\circ$

$\mu = 1.45$ mm⁻¹

$T = 293$ K

Block, colourless

0.20 × 0.10 × 0.10 mm

Data collection

Enraf–Nonius CAD-4
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
3091 measured reflections
2967 independent reflections
2041 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$
 $\theta_{\text{max}} = 25.2^\circ, \theta_{\text{min}} = 1.5^\circ$
 $h = 0 \rightarrow 11$
 $k = 0 \rightarrow 15$
 $l = 0 \rightarrow 32$
3 standard reflections every 100 reflections
intensity decay: none

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.068$
 $wR(F^2) = 0.207$
 $S = 1.05$
2967 reflections
220 parameters
40 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1255P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.50 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.61 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn	0.18405 (6)	0.12667 (5)	0.12048 (3)	0.0338 (3)
N1	0.1883 (6)	0.3348 (5)	0.2326 (2)	0.0594 (17)
N2	0.1689 (5)	0.2071 (4)	0.1829 (2)	0.0434 (13)
N3	0.2381 (6)	0.3608 (4)	0.0199 (2)	0.0461 (14)
N4	0.1948 (5)	0.2244 (4)	0.06251 (18)	0.0358 (12)
O1	0.3326 (4)	0.0294 (4)	0.11870 (16)	0.0460 (12)
O2	0.4748 (4)	0.1603 (3)	0.12273 (18)	0.0510 (13)
O3	1.0442 (4)	-0.0567 (4)	0.1520 (2)	0.0680 (16)
O4	-0.0041 (4)	0.0763 (3)	0.10656 (16)	0.0440 (11)
C1	0.2355 (10)	0.4326 (6)	0.2511 (3)	0.081
H1A	0.3017	0.4605	0.2284	0.122*
H1B	0.2777	0.4239	0.2830	0.122*
H1C	0.1581	0.4776	0.2539	0.122*
C2	0.0944 (8)	0.2711 (7)	0.2534 (3)	0.079 (3)
H2A	0.0483	0.2798	0.2833	0.094*
C3	0.0821 (8)	0.1945 (7)	0.2223 (3)	0.071 (2)

H3A	0.0229	0.1399	0.2267	0.086*
C4	0.2306 (8)	0.2910 (6)	0.1911 (2)	0.0523 (18)
H4B	0.2976	0.3181	0.1703	0.063*
C5	0.2982 (8)	0.4589 (5)	0.0065 (3)	0.060 (2)
H5A	0.3727	0.4745	0.0288	0.089*
H5B	0.2287	0.5104	0.0089	0.089*
H5C	0.3326	0.4560	-0.0268	0.089*
C6	0.2723 (6)	0.3054 (5)	0.0588 (2)	0.0408 (15)
H6A	0.3427	0.3218	0.0809	0.049*
C7	0.1029 (8)	0.2310 (5)	0.0239 (3)	0.0568 (19)
H7A	0.0326	0.1849	0.0175	0.068*
C8	0.1294 (8)	0.3137 (5)	-0.0030 (3)	0.059 (2)
H8A	0.0835	0.3349	-0.0315	0.070*
C9	0.9641 (6)	-0.0064 (5)	0.1281 (2)	0.0361 (14)
C10	0.8172 (5)	-0.0403 (5)	0.1219 (2)	0.0328 (14)
C11	0.7891 (7)	-0.1438 (5)	0.1187 (2)	0.0436 (16)
H11A	0.8608	-0.1907	0.1190	0.052*
C12	0.6544 (7)	-0.1755 (5)	0.1151 (3)	0.060 (2)
H12A	0.6356	-0.2443	0.1126	0.072*
C13	0.5473 (7)	-0.1080 (5)	0.1151 (3)	0.0510 (18)
H13A	0.4572	-0.1318	0.1121	0.061*
C14	0.5702 (5)	-0.0054 (4)	0.1195 (2)	0.0301 (12)
C15	0.7082 (5)	0.0286 (4)	0.1222 (2)	0.0321 (13)
H15A	0.7265	0.0975	0.1242	0.039*
C16	0.4545 (6)	0.0702 (5)	0.1210 (2)	0.0340 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn	0.0166 (4)	0.0401 (4)	0.0446 (5)	0.0005 (3)	-0.0007 (3)	0.0023 (3)
N1	0.067 (4)	0.058 (3)	0.053 (3)	-0.018 (3)	0.016 (3)	-0.013 (3)
N2	0.032 (3)	0.056 (3)	0.043 (3)	-0.008 (2)	0.006 (2)	0.001 (2)
N3	0.051 (3)	0.052 (3)	0.035 (3)	-0.008 (3)	0.003 (3)	0.003 (3)
N4	0.024 (2)	0.043 (3)	0.041 (3)	-0.009 (2)	-0.009 (2)	0.006 (2)
O1	0.019 (2)	0.057 (3)	0.062 (3)	0.004 (2)	0.0020 (19)	0.003 (2)
O2	0.021 (2)	0.036 (2)	0.096 (4)	-0.0011 (19)	-0.001 (2)	0.004 (2)
O3	0.023 (2)	0.075 (3)	0.106 (4)	-0.001 (2)	-0.013 (3)	0.041 (3)
O4	0.020 (2)	0.049 (3)	0.063 (3)	-0.0021 (19)	-0.004 (2)	0.008 (2)
C1	0.081	0.081	0.081	0.000	0.000	0.000
C2	0.068 (5)	0.100 (6)	0.068 (5)	-0.024 (4)	0.038 (4)	-0.019 (4)
C3	0.064 (5)	0.081 (5)	0.069 (5)	-0.030 (4)	0.031 (4)	-0.006 (4)
C4	0.052 (4)	0.071 (4)	0.033 (3)	-0.010 (3)	0.015 (3)	-0.005 (3)
C5	0.069 (5)	0.054 (4)	0.056 (5)	-0.010 (4)	0.008 (4)	0.004 (4)
C6	0.034 (3)	0.051 (4)	0.037 (4)	-0.006 (3)	0.001 (3)	0.003 (3)
C7	0.053 (4)	0.054 (4)	0.063 (5)	-0.009 (4)	-0.017 (4)	0.005 (4)
C8	0.061 (5)	0.060 (5)	0.054 (5)	-0.004 (4)	-0.017 (4)	0.017 (4)
C9	0.017 (3)	0.046 (3)	0.045 (4)	0.006 (3)	0.000 (3)	-0.007 (3)
C10	0.011 (3)	0.054 (4)	0.033 (3)	-0.002 (2)	0.002 (2)	0.005 (3)

C11	0.026 (3)	0.041 (4)	0.064 (4)	0.005 (3)	-0.006 (3)	-0.007 (3)
C12	0.029 (4)	0.031 (3)	0.119 (7)	-0.006 (3)	-0.005 (4)	-0.006 (4)
C13	0.024 (3)	0.042 (4)	0.087 (5)	-0.005 (3)	-0.002 (3)	-0.007 (3)
C14	0.015 (3)	0.037 (3)	0.038 (3)	0.002 (2)	-0.004 (2)	0.001 (3)
C15	0.017 (3)	0.034 (3)	0.046 (4)	-0.001 (2)	-0.002 (2)	0.005 (3)
C16	0.012 (3)	0.049 (4)	0.041 (3)	0.001 (3)	0.001 (2)	0.008 (3)

Geometric parameters (\AA , $^\circ$)

Zn—O1	1.930 (4)	C3—H3A	0.9300
Zn—O4	1.976 (4)	C4—H4B	0.9300
Zn—N2	1.996 (6)	C5—H5A	0.9600
Zn—N4	2.032 (5)	C5—H5B	0.9600
N1—C4	1.327 (8)	C5—H5C	0.9600
N1—C2	1.360 (9)	C6—H6A	0.9300
N1—C1	1.459 (10)	C7—C8	1.339 (9)
N2—C4	1.279 (8)	C7—H7A	0.9300
N2—C3	1.367 (8)	C8—H8A	0.9300
N3—C6	1.322 (8)	C9—O4 ⁱⁱ	1.277 (8)
N3—C8	1.371 (9)	C9—C10	1.501 (7)
N3—C5	1.468 (8)	C10—C15	1.395 (8)
N4—C6	1.311 (7)	C10—C11	1.399 (9)
N4—C7	1.372 (8)	C11—C12	1.373 (8)
O1—C16	1.299 (7)	C11—H11A	0.9300
O2—C16	1.209 (8)	C12—C13	1.368 (9)
O3—C9	1.207 (7)	C12—H12A	0.9300
O4—C9 ⁱ	1.277 (8)	C13—C14	1.381 (8)
C1—H1A	0.9600	C13—H13A	0.9300
C1—H1B	0.9600	C14—C15	1.411 (7)
C1—H1C	0.9600	C14—C16	1.502 (8)
C2—C3	1.319 (11)	C15—H15A	0.9300
C2—H2A	0.9300		
O1—Zn—O4	117.23 (19)	H5A—C5—H5B	109.5
O1—Zn—N2	115.5 (2)	N3—C5—H5C	109.5
O4—Zn—N2	105.80 (19)	H5A—C5—H5C	109.5
O1—Zn—N4	111.51 (19)	H5B—C5—H5C	109.5
O4—Zn—N4	96.63 (18)	N4—C6—N3	111.7 (6)
N2—Zn—N4	108.3 (2)	N4—C6—H6A	124.2
C4—N1—C2	106.5 (6)	N3—C6—H6A	124.2
C4—N1—C1	125.3 (7)	C8—C7—N4	109.9 (6)
C2—N1—C1	128.2 (7)	C8—C7—H7A	125.1
C4—N2—C3	104.9 (6)	N4—C7—H7A	125.1
C4—N2—Zn	125.0 (5)	C7—C8—N3	105.8 (6)
C3—N2—Zn	129.6 (5)	C7—C8—H8A	127.1
C6—N3—C8	107.4 (5)	N3—C8—H8A	127.1
C6—N3—C5	125.9 (6)	O3—C9—O4 ⁱⁱ	124.1 (6)
C8—N3—C5	126.5 (6)	O3—C9—C10	120.2 (6)

C6—N4—C7	105.2 (5)	O4 ⁱⁱ —C9—C10	115.7 (5)
C6—N4—Zn	127.4 (4)	C15—C10—C11	119.5 (5)
C7—N4—Zn	126.2 (4)	C15—C10—C9	121.5 (6)
C16—O1—Zn	113.5 (4)	C11—C10—C9	118.9 (5)
C9 ⁱ —O4—Zn	115.1 (4)	C12—C11—C10	119.2 (6)
N1—C1—H1A	109.5	C12—C11—H11A	120.4
N1—C1—H1B	109.5	C10—C11—H11A	120.4
H1A—C1—H1B	109.5	C13—C12—C11	121.4 (6)
N1—C1—H1C	109.5	C13—C12—H12A	119.3
H1A—C1—H1C	109.5	C11—C12—H12A	119.3
H1B—C1—H1C	109.5	C12—C13—C14	121.2 (6)
C3—C2—N1	105.9 (7)	C12—C13—H13A	119.4
C3—C2—H2A	127.0	C14—C13—H13A	119.4
N1—C2—H2A	127.0	C13—C14—C15	118.0 (5)
C2—C3—N2	110.2 (7)	C13—C14—C16	122.4 (5)
C2—C3—H3A	124.9	C15—C14—C16	119.5 (5)
N2—C3—H3A	124.9	C10—C15—C14	120.5 (5)
N2—C4—N1	112.4 (6)	C10—C15—H15A	119.7
N2—C4—H4B	123.8	C14—C15—H15A	119.7
N1—C4—H4B	123.8	O2—C16—O1	124.0 (6)
N3—C5—H5A	109.5	O2—C16—C14	122.4 (5)
N3—C5—H5B	109.5	O1—C16—C14	113.6 (5)
O1—Zn—N2—C4	89.0 (6)	Zn—N4—C6—N3	−169.7 (4)
O4—Zn—N2—C4	−139.6 (6)	C8—N3—C6—N4	0.9 (8)
N4—Zn—N2—C4	−36.9 (6)	C5—N3—C6—N4	176.2 (6)
O1—Zn—N2—C3	−100.1 (7)	C6—N4—C7—C8	1.9 (8)
O4—Zn—N2—C3	31.4 (7)	Zn—N4—C7—C8	170.1 (5)
N4—Zn—N2—C3	134.1 (7)	N4—C7—C8—N3	−1.4 (9)
O1—Zn—N4—C6	−81.7 (5)	C6—N3—C8—C7	0.3 (9)
O4—Zn—N4—C6	155.6 (5)	C5—N3—C8—C7	−175.0 (6)
N2—Zn—N4—C6	46.5 (6)	O3—C9—C10—C15	141.9 (7)
O1—Zn—N4—C7	112.7 (6)	O4 ⁱⁱ —C9—C10—C15	−38.6 (9)
O4—Zn—N4—C7	−10.0 (6)	O3—C9—C10—C11	−34.3 (9)
N2—Zn—N4—C7	−119.1 (6)	O4 ⁱⁱ —C9—C10—C11	145.2 (6)
O4—Zn—O1—C16	170.5 (4)	C15—C10—C11—C12	1.0 (10)
N2—Zn—O1—C16	−63.7 (4)	C9—C10—C11—C12	177.3 (6)
N4—Zn—O1—C16	60.4 (4)	C10—C11—C12—C13	−0.7 (12)
O1—Zn—O4—C9 ⁱ	44.5 (5)	C11—C12—C13—C14	−1.0 (12)
N2—Zn—O4—C9 ⁱ	−86.0 (4)	C12—C13—C14—C15	2.4 (11)
N4—Zn—O4—C9 ⁱ	162.9 (4)	C12—C13—C14—C16	−178.4 (6)
C4—N1—C2—C3	−2.1 (10)	C11—C10—C15—C14	0.3 (9)
C1—N1—C2—C3	177.6 (8)	C9—C10—C15—C14	−175.9 (5)
N1—C2—C3—N2	1.5 (11)	C13—C14—C15—C10	−2.0 (10)
C4—N2—C3—C2	−0.2 (10)	C16—C14—C15—C10	178.7 (5)
Zn—N2—C3—C2	−172.6 (6)	Zn—O1—C16—O2	−1.8 (8)
C3—N2—C4—N1	−1.2 (9)	Zn—O1—C16—C14	−180.0 (4)
Zn—N2—C4—N1	171.6 (5)	C13—C14—C16—O2	−176.8 (7)

C2—N1—C4—N2	2.1 (9)	C15—C14—C16—O2	2.4 (9)
C1—N1—C4—N2	−177.6 (7)	C13—C14—C16—O1	1.4 (8)
C7—N4—C6—N3	−1.7 (7)	C15—C14—C16—O1	−179.4 (5)

Symmetry codes: (i) $x-1, y, z$; (ii) $x+1, y, z$.

Hydrogen-bond geometry (Å, °)

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C1—H1A···O3 ⁱⁱⁱ	0.96	2.55	3.423 (10)	150
C4—H4B···O3 ⁱⁱⁱ	0.93	2.31	3.150 (9)	150
C11—H11A···O2 ^{iv}	0.93	2.54	3.457 (8)	171

Symmetry codes: (iii) $-x+3/2, y+1/2, z$; (iv) $-x+3/2, y-1/2, z$.