

## Tetrakis( $\mu$ -2-chlorobenzoato- $\kappa^2$ O:O')-bis[(4-vinylpyridine- $\kappa$ N)copper(II)]

Juan Zhao

College of Mechanical Engineering, Qingdao Technological University, Qingdao 266033, People's Republic of China  
Correspondence e-mail: zhaojuanqd@163.com


Received 19 September 2008; accepted 24 September 2008

Key indicators: single-crystal X-ray study;  $T = 297$  K; mean  $\sigma(C-C) = 0.009$  Å;  
 $R$  factor = 0.059;  $wR$  factor = 0.166; data-to-parameter ratio = 14.1.

The title compound,  $[Cu_2(C_7H_4ClO_2)_4(C_7H_7N)_2]$ , consists of centrosymmetric dinuclear molecules with a Cu···Cu separation of 2.6676 (12) Å. In the molecule, four 2-chlorobenzoate anions bridge two Cu<sup>II</sup> ions, while two neutral 4-vinylpyridine ligands coordinate them in axial positions. The Cu<sup>II</sup> ion has a distorted square-planar pyramidal coordination, with four O atoms from the chlorobenzoate anions at the base. The N pyridine atom completes the coordination environment in the apical position.

### Related literature

In the corresponding dinuclear compound [tetrakis( $\mu$ -acetato)bis(2-anilinopyridine)dicopper(II)] (Seco *et al.*, 2002), the Cu<sup>II</sup> has a distorted square-planar pyramidal coordination environment.



### Experimental

#### Crystal data

$[Cu_2(C_7H_4ClO_2)_4(C_7H_7N)_2]$	$V = 2069.2$ (8) Å <sup>3</sup>
$M_r = 959.58$	$Z = 2$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 10.251$ (2) Å	$\mu = 1.34$ mm <sup>-1</sup>
$b = 20.412$ (4) Å	$T = 297$ (2) K
$c = 10.665$ (2) Å	$0.30 \times 0.30 \times 0.20$ mm
$\beta = 111.99$ (3)°	

#### Data collection

Enraf–Nonius CAD-4	3689 independent reflections
diffractometer	2587 reflections with $I > 2\sigma(I)$
Absorption correction: $\psi$ scan	$R_{\text{int}} = 0.029$
(North <i>et al.</i> , 1968)	3 standard reflections
$T_{\min} = 0.677$ , $T_{\max} = 0.767$	every 100 reflections
3708 measured reflections	intensity decay: none

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$	262 parameters
$wR(F^2) = 0.166$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.37$ e Å <sup>-3</sup>
3689 reflections	$\Delta\rho_{\min} = -0.56$ e Å <sup>-3</sup>

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

This work was supported by the National Natural Science Foundation of China (grant No. 20601015) and the Natural Science Foundation of Shandong Province (grant No. Y2006B12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2452).

### References

- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Seco, J. M., González Garmendia, M. J., Pinilla, E. & Torres, M. R. (2002). *Polyhedron*, **21**, 457–464.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

# supporting information

*Acta Cryst.* (2008). E64, m1336 [doi:10.1107/S1600536808030766]

## **Tetrakis( $\mu$ -2-chlorobenzoato- $\kappa^2 O:O'$ )bis[(4-vinylpyridine- $\kappa N$ )copper(II)]**

**Juan Zhao**

### **S1. Comment**


The title compound, (I) (Fig. 1), consists of centrosymmetric dinuclear units, in which four 2-chlorobenzoato groups bridge the two copper ions and a 4-vinylpyridine neutral ligand occupies the axis position of each copper atom, coordinating them through the nitrogen atom. Each copper ion has a distorted square-planar pyramidal coordination, with four oxygen atoms in a plane. The distances for Cu—O1,O2,O3 and O4 are 1.975 (4), 1.957 (4), 1.969 (3) and 1.985 (3) Å, respectively. The fifth coordination position is occupied by the pyridine nitrogen, N, of a ligand molecule at 2.134 (4) Å. All these values agree well with those observed in  $[\text{Cu}_2(v\text{-OOCCH}_3)_4(\text{PhNHpy})_2]$  (PhNHpy is 2-anilinopyridine) (Seco *et al.*, 2002). The Cu…Cu separation in (I) is 2.6473 (12) Å.

### **S2. Experimental**

A solution of 4-vinylpyridine (1.05 g, 10 mmol) in alcohol (10 ml) was added to a solution of  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  (1.70 g, 10 mmol) and 2-chlorobenzoic acid (1.56 g, 10 mmol) and KOH (0.56 g, 10 mmol) in alcohol (40 ml). The solution was stirred during 2 h and a precipitate was formed. The blue precipitate was filtered off, washed with alcohol and dried *in vacuo* over  $\text{CaCO}_3$ . Blue crystals were obtained from recrystallization in alcohol after a few days.

### **S3. Refinement**

H atoms were positioned geometrically (C—H = 0.93 Å) and allowed to ride on their parent atoms with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

The molecular structure of (I), showing 30% probability displacement ellipsoids and the atom-numbering scheme. Unlabelled atoms are related with the labelled ones by symmetry operation ( $-x, -y, 3-z$ ).

### Tetrakis( $\mu$ -2-chlorobenzoato- $\kappa^2$ O:O')bis[(4-vinylpyridine- $\kappa$ N)copper(II)]

#### Crystal data



$M_r = 959.58$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 10.251 (2)$  Å

$b = 20.412 (4)$  Å

$c = 10.665 (2)$  Å

$\beta = 111.99 (3)^\circ$

$V = 2069.2 (8)$  Å<sup>3</sup>

$Z = 2$

$F(000) = 972$

$D_x = 1.540 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 25 reflections

$\theta = 10\text{--}14^\circ$

$\mu = 1.34 \text{ mm}^{-1}$

$T = 297$  K

Block, blue

$0.30 \times 0.30 \times 0.20$  mm

#### Data collection

Enraf–Nonius CAD-4  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction:  $\psi$  scan  
(North et al., 1968)

$T_{\min} = 0.677$ ,  $T_{\max} = 0.767$

3708 measured reflections

3689 independent reflections

2587 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 25.2^\circ$ ,  $\theta_{\min} = 2.0^\circ$

$h = -12 \rightarrow 11$

$k = 0 \rightarrow 24$

$l = 0 \rightarrow 12$

3 standard reflections every 100 reflections

intensity decay: none

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.059$  $wR(F^2) = 0.166$  $S = 1.00$ 

3689 reflections

262 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.5P]$   
where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\text{max}} < 0.001$  $\Delta\rho_{\text{max}} = 0.37 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\text{min}} = -0.57 \text{ e } \text{\AA}^{-3}$ *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu	-0.05229 (6)	0.05461 (3)	1.53164 (6)	0.0465 (2)
N1	-0.1437 (4)	0.1341 (2)	1.5999 (4)	0.0483 (9)
Cl1	0.52811 (19)	-0.03630 (12)	1.6765 (3)	0.1233 (9)
Cl2	0.2716 (2)	0.17779 (10)	1.41176 (18)	0.0975 (6)
O1	-0.2292 (4)	0.0228 (2)	1.3923 (4)	0.0691 (10)
O2	0.1417 (4)	0.0684 (2)	1.6572 (4)	0.0707 (11)
O3	-0.0022 (4)	0.09641 (17)	1.3891 (4)	0.0610 (9)
O4	-0.0849 (4)	-0.00570 (19)	1.6630 (3)	0.0634 (10)
C1	-0.4145 (9)	0.2796 (5)	1.8136 (7)	0.119 (3)
H1A	-0.4628	0.2401	1.8018	0.142*
H1B	-0.4424	0.3151	1.8522	0.142*
C2	-0.3097 (7)	0.2851 (3)	1.7770 (5)	0.0787 (18)
H2A	-0.2638	0.3253	1.7903	0.094*
C3	-0.2568 (6)	0.2323 (3)	1.7152 (5)	0.0568 (13)
C4	-0.3184 (6)	0.1719 (3)	1.6807 (5)	0.0604 (13)
H4A	-0.4007	0.1627	1.6947	0.072*
C5	-0.2598 (5)	0.1244 (3)	1.6253 (5)	0.0561 (12)
H5A	-0.3036	0.0837	1.6050	0.067*
C6	-0.0865 (5)	0.1938 (3)	1.6306 (5)	0.0604 (13)
H6A	-0.0062	0.2027	1.6126	0.072*
C7	-0.1379 (6)	0.2424 (3)	1.6864 (5)	0.0693 (15)
H7A	-0.0924	0.2827	1.7054	0.083*
C8	0.5064 (9)	0.1132 (5)	1.9717 (8)	0.125 (3)
H8A	0.5043	0.1444	2.0344	0.151*
C9	0.3825 (7)	0.0961 (4)	1.8658 (6)	0.094 (2)

H9A	0.2991	0.1173	1.8564	0.113*
C10	0.3818 (5)	0.0475 (3)	1.7737 (5)	0.0560 (13)
C11	0.5093 (6)	0.0209 (3)	1.7889 (6)	0.0678 (15)
C12	0.6323 (7)	0.0384 (4)	1.8922 (8)	0.095 (2)
H12A	0.7168	0.0189	1.8996	0.114*
C13	0.6297 (8)	0.0848 (5)	1.9844 (8)	0.114 (3)
H13A	0.7122	0.0967	2.0550	0.137*
C14	0.2409 (5)	0.0288 (3)	1.6714 (5)	0.0540 (12)
C15	0.1325 (8)	0.1677 (4)	1.0103 (8)	0.091 (2)
H15A	0.1470	0.1902	0.9408	0.109*
C16	0.2011 (7)	0.1873 (3)	1.1424 (7)	0.0791 (18)
H16A	0.2622	0.2229	1.1625	0.095*
C17	0.1783 (6)	0.1538 (3)	1.2444 (6)	0.0643 (14)
C18	0.0894 (5)	0.1012 (2)	1.2185 (5)	0.0495 (11)
C19	0.0226 (6)	0.0818 (3)	1.0849 (5)	0.0713 (15)
H19A	-0.0374	0.0458	1.0640	0.086*
C20	0.0454 (7)	0.1163 (4)	0.9819 (6)	0.091 (2)
H20A	-0.0006	0.1034	0.8924	0.109*
C21	0.0558 (5)	0.0649 (3)	1.3239 (5)	0.0488 (12)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu	0.0445 (3)	0.0547 (4)	0.0436 (3)	0.0030 (3)	0.0201 (2)	-0.0043 (3)
N1	0.045 (2)	0.059 (2)	0.045 (2)	0.0024 (18)	0.0206 (18)	0.0000 (19)
Cl1	0.0601 (10)	0.1388 (18)	0.160 (2)	0.0017 (10)	0.0292 (12)	-0.0630 (16)
Cl2	0.0979 (12)	0.1124 (14)	0.0906 (12)	-0.0378 (11)	0.0451 (10)	-0.0404 (11)
O1	0.049 (2)	0.093 (3)	0.057 (2)	0.000 (2)	0.0117 (17)	-0.010 (2)
O2	0.053 (2)	0.076 (3)	0.075 (3)	0.0039 (19)	0.0150 (19)	-0.024 (2)
O3	0.073 (2)	0.056 (2)	0.069 (2)	0.0110 (18)	0.044 (2)	0.0070 (18)
O4	0.084 (3)	0.064 (2)	0.056 (2)	0.0137 (19)	0.041 (2)	0.0100 (18)
C1	0.120 (7)	0.134 (7)	0.104 (6)	0.044 (6)	0.044 (5)	-0.036 (5)
C2	0.094 (5)	0.082 (4)	0.049 (3)	0.026 (4)	0.014 (3)	-0.011 (3)
C3	0.063 (3)	0.063 (3)	0.040 (3)	0.013 (3)	0.014 (2)	-0.001 (2)
C4	0.056 (3)	0.069 (4)	0.062 (3)	0.008 (3)	0.030 (3)	-0.002 (3)
C5	0.060 (3)	0.052 (3)	0.060 (3)	-0.002 (2)	0.028 (3)	-0.008 (2)
C6	0.056 (3)	0.054 (3)	0.073 (4)	-0.006 (2)	0.026 (3)	-0.004 (3)
C7	0.077 (4)	0.058 (3)	0.063 (4)	-0.002 (3)	0.015 (3)	-0.011 (3)
C8	0.089 (5)	0.182 (10)	0.089 (5)	-0.022 (6)	0.015 (4)	-0.063 (6)
C9	0.061 (4)	0.140 (7)	0.073 (4)	-0.009 (4)	0.015 (3)	-0.037 (4)
C10	0.051 (3)	0.069 (4)	0.045 (3)	-0.005 (2)	0.014 (2)	0.004 (2)
C11	0.058 (3)	0.071 (4)	0.068 (4)	-0.005 (3)	0.017 (3)	0.000 (3)
C12	0.051 (3)	0.128 (6)	0.090 (5)	-0.007 (4)	0.007 (3)	-0.013 (5)
C13	0.075 (5)	0.161 (8)	0.083 (5)	-0.017 (5)	0.002 (4)	-0.030 (5)
C14	0.053 (3)	0.074 (3)	0.040 (3)	-0.007 (3)	0.023 (2)	0.000 (3)
C15	0.081 (5)	0.114 (6)	0.092 (5)	0.017 (4)	0.048 (4)	0.039 (5)
C16	0.081 (4)	0.075 (4)	0.097 (5)	-0.001 (3)	0.051 (4)	0.017 (4)
C17	0.061 (3)	0.074 (4)	0.068 (4)	0.004 (3)	0.037 (3)	0.001 (3)

C18	0.048 (3)	0.056 (3)	0.049 (3)	0.007 (2)	0.023 (2)	0.006 (2)
C19	0.066 (3)	0.096 (4)	0.047 (3)	-0.002 (3)	0.015 (3)	0.002 (3)
C20	0.085 (5)	0.131 (6)	0.051 (4)	0.010 (5)	0.017 (3)	0.021 (4)
C21	0.038 (2)	0.066 (4)	0.043 (3)	0.003 (2)	0.015 (2)	0.004 (2)

Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )

Cu—O2	1.958 (4)	C7—H7A	0.9300
Cu—O3	1.971 (3)	C8—C13	1.350 (12)
Cu—O1	1.975 (4)	C8—C9	1.392 (10)
Cu—O4	1.985 (3)	C8—H8A	0.9300
Cu—N1	2.134 (4)	C9—C10	1.393 (8)
Cu—Cu <sup>i</sup>	2.6676 (12)	C9—H9A	0.9300
N1—C5	1.331 (6)	C10—C11	1.368 (8)
N1—C6	1.339 (6)	C10—C14	1.497 (7)
Cl1—C11	1.735 (6)	C11—C12	1.375 (8)
Cl2—C17	1.750 (6)	C12—C13	1.372 (11)
O1—C14 <sup>i</sup>	1.236 (6)	C12—H12A	0.9300
O2—C14	1.262 (6)	C13—H13A	0.9300
O3—C21	1.250 (6)	C14—O1 <sup>i</sup>	1.236 (6)
O4—C21 <sup>i</sup>	1.240 (6)	C15—C20	1.336 (10)
C1—C2	1.279 (10)	C15—C16	1.378 (10)
C1—H1A	0.9300	C15—H15A	0.9300
C1—H1B	0.9300	C16—C17	1.377 (8)
C2—C3	1.469 (7)	C16—H16A	0.9300
C2—H2A	0.9300	C17—C18	1.369 (7)
C3—C4	1.372 (8)	C18—C19	1.388 (7)
C3—C7	1.379 (8)	C18—C21	1.489 (6)
C4—C5	1.383 (7)	C19—C20	1.397 (8)
C4—H4A	0.9300	C19—H19A	0.9300
C5—H5A	0.9300	C20—H20A	0.9300
C6—C7	1.359 (7)	C21—O4 <sup>i</sup>	1.240 (6)
C6—H6A	0.9300		
O2—Cu—O3	88.52 (17)	C13—C8—C9	120.6 (8)
O2—Cu—O1	166.73 (16)	C13—C8—H8A	119.7
O3—Cu—O1	89.60 (16)	C9—C8—H8A	119.7
O2—Cu—O4	90.15 (17)	C8—C9—C10	120.9 (7)
O3—Cu—O4	166.87 (14)	C8—C9—H9A	119.6
O1—Cu—O4	88.70 (17)	C10—C9—H9A	119.6
O2—Cu—N1	96.94 (15)	C11—C10—C9	116.5 (5)
O3—Cu—N1	102.02 (14)	C11—C10—C14	127.1 (5)
O1—Cu—N1	96.30 (16)	C9—C10—C14	116.4 (5)
O4—Cu—N1	91.12 (15)	C10—C11—C12	122.7 (6)
O2—Cu—Cu <sup>i</sup>	83.76 (11)	C10—C11—Cl1	122.3 (4)
O3—Cu—Cu <sup>i</sup>	85.49 (10)	C12—C11—Cl1	115.0 (5)
O1—Cu—Cu <sup>i</sup>	83.00 (12)	C13—C12—C11	119.7 (7)
O4—Cu—Cu <sup>i</sup>	81.38 (11)	C13—C12—H12A	120.1

N1—Cu—Cu <sup>i</sup>	172.47 (11)	C11—C12—H12A	120.1
C5—N1—C6	115.5 (4)	C8—C13—C12	119.5 (7)
C5—N1—Cu	119.7 (3)	C8—C13—H13A	120.2
C6—N1—Cu	124.6 (3)	C12—C13—H13A	120.2
C14 <sup>i</sup> —O1—Cu	124.5 (4)	O1 <sup>i</sup> —C14—O2	124.9 (5)
C14—O2—Cu	123.8 (4)	O1 <sup>i</sup> —C14—C10	119.1 (5)
C21—O3—Cu	121.5 (3)	O2—C14—C10	116.0 (5)
C21 <sup>i</sup> —O4—Cu	126.0 (3)	C20—C15—C16	120.2 (6)
C2—C1—H1A	120.0	C20—C15—H15A	119.9
C2—C1—H1B	120.0	C16—C15—H15A	119.9
H1A—C1—H1B	120.0	C17—C16—C15	119.4 (6)
C1—C2—C3	124.5 (8)	C17—C16—H16A	120.3
C1—C2—H2A	117.8	C15—C16—H16A	120.3
C3—C2—H2A	117.8	C18—C17—C16	121.9 (6)
C4—C3—C7	115.5 (5)	C18—C17—Cl2	119.6 (4)
C4—C3—C2	124.8 (5)	C16—C17—Cl2	118.5 (5)
C7—C3—C2	119.7 (6)	C17—C18—C19	117.8 (5)
C3—C4—C5	121.0 (5)	C17—C18—C21	124.3 (5)
C3—C4—H4A	119.5	C19—C18—C21	117.9 (5)
C5—C4—H4A	119.5	C18—C19—C20	120.0 (6)
N1—C5—C4	123.0 (5)	C18—C19—H19A	120.0
N1—C5—H5A	118.5	C20—C19—H19A	120.0
C4—C5—H5A	118.5	C15—C20—C19	120.8 (7)
N1—C6—C7	124.4 (5)	C15—C20—H20A	119.6
N1—C6—H6A	117.8	C19—C20—H20A	119.6
C7—C6—H6A	117.8	O4 <sup>i</sup> —C21—O3	125.6 (4)
C6—C7—C3	120.5 (5)	O4 <sup>i</sup> —C21—C18	117.1 (4)
C6—C7—H7A	119.7	O3—C21—C18	117.2 (4)
C3—C7—H7A	119.7		
O2—Cu—N1—C5	-132.2 (4)	C4—C3—C7—C6	-1.5 (8)
O3—Cu—N1—C5	137.8 (4)	C2—C3—C7—C6	179.3 (5)
O1—Cu—N1—C5	46.9 (4)	C13—C8—C9—C10	2.6 (15)
O4—Cu—N1—C5	-41.9 (4)	C8—C9—C10—C11	-3.5 (11)
O3—Cu—N1—C6	-47.9 (4)	C8—C9—C10—C14	174.7 (7)
O1—Cu—N1—C6	-138.8 (4)	C9—C10—C11—C12	2.6 (9)
O4—Cu—N1—C6	132.4 (4)	C14—C10—C11—C12	-175.4 (6)
O2—Cu—O1—C14 <sup>i</sup>	4.6 (10)	C9—C10—C11—Cl1	-175.9 (5)
O3—Cu—O1—C14 <sup>i</sup>	86.4 (4)	C14—C10—C11—Cl1	6.1 (8)
O4—Cu—O1—C14 <sup>i</sup>	-80.5 (4)	C10—C11—C12—C13	-0.5 (12)
N1—Cu—O1—C14 <sup>i</sup>	-171.5 (4)	Cl1—C11—C12—C13	178.0 (7)
Cu <sup>i</sup> —Cu—O1—C14 <sup>i</sup>	0.9 (4)	C9—C8—C13—C12	-0.4 (16)
O3—Cu—O2—C14	-86.8 (4)	C11—C12—C13—C8	-0.6 (14)
O1—Cu—O2—C14	-4.8 (10)	Cu—O2—C14—O1 <sup>i</sup>	0.8 (8)
O4—Cu—O2—C14	80.2 (4)	Cu—O2—C14—C10	-179.3 (3)
N1—Cu—O2—C14	171.3 (4)	C11—C10—C14—O1 <sup>i</sup>	14.1 (8)
Cu <sup>i</sup> —Cu—O2—C14	-1.1 (4)	C9—C10—C14—O1 <sup>i</sup>	-163.9 (6)
O2—Cu—O3—C21	82.9 (4)	C11—C10—C14—O2	-165.8 (6)

O1—Cu—O3—C21	−84.0 (4)	C9—C10—C14—O2	16.2 (7)
O4—Cu—O3—C21	−1.4 (9)	C20—C15—C16—C17	−0.3 (11)
N1—Cu—O3—C21	179.7 (4)	C15—C16—C17—C18	0.1 (9)
Cu <sup>i</sup> —Cu—O3—C21	−1.0 (4)	C15—C16—C17—Cl2	177.6 (5)
O2—Cu—O4—C21 <sup>i</sup>	−84.4 (4)	C16—C17—C18—C19	0.6 (8)
O3—Cu—O4—C21 <sup>i</sup>	−0.3 (10)	Cl2—C17—C18—C19	−176.9 (4)
O1—Cu—O4—C21 <sup>i</sup>	82.4 (4)	C16—C17—C18—C21	−177.2 (5)
N1—Cu—O4—C21 <sup>i</sup>	178.6 (4)	Cl2—C17—C18—C21	5.3 (7)
Cu <sup>i</sup> —Cu—O4—C21 <sup>i</sup>	−0.8 (4)	C17—C18—C19—C20	−1.0 (8)
C1—C2—C3—C4	4.6 (10)	C21—C18—C19—C20	176.9 (5)
C1—C2—C3—C7	−176.3 (7)	C16—C15—C20—C19	−0.1 (11)
C7—C3—C4—C5	2.2 (8)	C18—C19—C20—C15	0.8 (10)
C2—C3—C4—C5	−178.6 (5)	Cu—O3—C21—O4 <sup>i</sup>	1.9 (7)
C6—N1—C5—C4	−0.5 (7)	Cu—O3—C21—C18	−179.9 (3)
Cu—N1—C5—C4	174.3 (4)	C17—C18—C21—O4 <sup>i</sup>	−120.3 (6)
C3—C4—C5—N1	−1.3 (8)	C19—C18—C21—O4 <sup>i</sup>	61.9 (6)
C5—N1—C6—C7	1.2 (8)	C17—C18—C21—O3	61.3 (7)
Cu—N1—C6—C7	−173.3 (4)	C19—C18—C21—O3	−116.5 (6)
N1—C6—C7—C3	−0.2 (9)		

Symmetry code: (i)  $-x, -y, -z+3$ .