

7-Dimethylamino-2-phenyl-1,2,4-triazolo[1,5-a][1,3,5]triazin-5-amine methanol solvate¹

Anton V. Dolzhenko,^{a,*} Geok Kheng Tan,^b Lip Lin Koh,^b Su Fen Woo^b and Wai Keung Chui^a

^aDepartment of Pharmacy, Faculty of Science, National University of Singapore, 18 Science Drive 4, Singapore 117543, Singapore, and ^bDepartment of Chemistry, Faculty of Science, National University of Singapore, 3 Science Drive 3, Singapore 117543, Singapore

Correspondence e-mail: phada@nus.edu.sg


Received 7 August 2008; accepted 22 September 2008

Key indicators: single-crystal X-ray study; $T = 223\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.047; wR factor = 0.139; data-to-parameter ratio = 15.9.

7-Dimethylamino-2-phenyl-1,2,4-triazolo[1,5-a][1,3,5]triazin-5-amine crystallized with one molecule of methanol to give the title compound, $\text{C}_{12}\text{H}_{13}\text{N}_7\cdot\text{CH}_3\text{OH}$. The triazolo[1,5-a]-[1,3,5]triazine heterocyclic core is essentially planar as are both amino groups that are involved in π -electron delocalization with the triazolo[1,5-a][1,3,5]triazine nucleus. The methyl groups of the dimethylamino fragment are involved in the formation of weak intramolecular C—H···N hydrogen bonds with the N atoms of the heterocyclic system. The crystal packing is stabilized by intermolecular N—H···N hydrogen bonds between the triazolo[1,5-a][1,3,5]triazine molecules. The methanol solvent molecule also participates in the formation of the crystal structure via intermolecular O—H···N, N—H···O and weak C—H···O hydrogen bonds, linking the layers of triazolo[1,5-a][1,3,5]triazine molecules.

Related literature

The 1,2,4-triazolo[1,5-a][1,3,5]triazine (5-azapurine) heterocyclic system has been reviewed by Dolzhenko *et al.* (2006). For investigations on 5,7-diamino-1,2,4-triazolo[1,5-a][1,3,5]-triazines, see Dolzhenko *et al.* (2007). For a similar structure, see: Gilardi (1973). For related literature, see: Dolzhenko *et al.* (2008)



¹ Part 11 in the series ‘Fused heterocyclic systems with an *s*-triazine ring’. For Part 10, see Dolzhenko *et al.* (2008).

Experimental

Crystal data

$\text{C}_{12}\text{H}_{13}\text{N}_7\cdot\text{CH}_3\text{OH}$	$\gamma = 102.883(1)^{\circ}$
$M_r = 287.34$	$V = 714.39(8)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.9963(5)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.0435(5)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$c = 13.0942(9)\text{ \AA}$	$T = 223(2)\text{ K}$
$\alpha = 93.493(1)^{\circ}$	$0.74 \times 0.68 \times 0.40\text{ mm}$
$\beta = 93.972(1)^{\circ}$	

Data collection

Bruker SMART APEX CCD diffractometer	9183 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2001)	3256 independent reflections
$T_{\min} = 0.935$, $T_{\max} = 0.964$	2870 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.139$	$\Delta\rho_{\text{max}} = 0.26\text{ e \AA}^{-3}$
$S = 1.07$	$\Delta\rho_{\text{min}} = -0.23\text{ e \AA}^{-3}$
3256 reflections	
205 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^{\circ}$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1S—H1S···N1	0.92 (2)	1.97 (2)	2.8861 (16)	172.8 (18)
N6—H6NB···N4 ⁱ	0.86 (2)	2.11 (2)	2.9679 (17)	178.8 (18)
N6—H6NA···O1S ⁱ	0.89 (2)	2.398 (19)	3.0280 (18)	128.3 (16)
C6—H6C···N2	0.97	2.08	2.8753 (18)	138
C6—H6C···N3	0.97	2.54	2.9484 (17)	105
C7—H7A···N5	0.97	2.22	2.6788 (18)	108
C7—H7C···O1S ⁱⁱ	0.97	2.48	3.4438 (19)	176

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by the Academic Research Fund of the National University of Singapore (WBS R-148-000-069-112) and the National Medical Research Council, Singapore (NMRC/NIG/0019/2008 and NMRC/NIG/0020/2008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FB2109).

References

- Bruker (2001). *SMART* and *SAINT*. Bruker AXS GmbH, Karlsruhe, Germany.
- Dolzhenko, A. V., Dolzhenko, A. V. & Chui, W. K. (2006). *Heterocycles*, **68**, 1723–1759.
- Dolzhenko, A. V., Dolzhenko, A. V. & Chui, W. K. (2007). *Heterocycles*, **71**, 429–436.
- Dolzhenko, A. V., Tan, B. J., Dolzhenko, A. V., Chiu, G. N. C. & Chui, W. K. (2008). *J. Fluorine Chem.*, **129**, 429–434.
- Gilardi, R. D. (1973). *Acta Cryst. B29*, 2089–2095.
- Sheldrick, G. M. (2001). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supporting information

Acta Cryst. (2008). E64, o2021 [doi:10.1107/S1600536808030481]

7-Dimethylamino-2-phenyl-1,2,4-triazolo[1,5-a][1,3,5]triazin-5-amine methanol solvate

Anton V. Dolzhenko, Geok Kheng Tan, Lip Lin Koh, Su Fen Woo and Wai Keung Chui

S1. Comment

1,2,4-Triazolo[1,5-*a*][1,3,5]triazine system is known as 5-aza-isostere of the purine core. Compounds based on this skeleton have been shown to possess a wide range of biological activities (Dolzhenko *et al.*, 2006). In continuation of our investigations on 5,7-diamino-1,2,4-triazolo[1,5-*a*][1,3,5]triazines (Dolzhenko *et al.*, 2007), we report herein the structural study of 7-dimethylamino-2-phenyl-1,2,4-triazolo[1,5-*a*][1,3,5]triazin-5-amine (Fig. 1).

The fused triazine and triazole rings are located practically in the same plane (the angle between the mean planes of these rings makes 1.66 (4)°). The phenyl ring makes a dihedral angle of 22.70 (5)° with the mean plane of the 1,2,4-triazolo[1,5-*a*][1,3,5]triazine core. Similarity of the lengths of C3—N4, C3—N5, C3—N6, C4—N3, C4—N5 and C4—N7 makes evidence for π -electron delocalization of the amino groups with the 1,2,4-triazolo[1,5-*a*][1,3,5]triazine core.

The dimethylamino group (C6—N7—C7) has an out-of-plane twist (4.3 (8)°). The nitrogen atom of dimethylamino group (N7) has a slightly pyramidal stereochemistry [C6—N7—C7 = 115.0 (1)°] and it is located 0.039 (1) Å above the C4/C6/C7 plane. These data are in good agreement with previously reported results on the similar structure of 5,7-bis(dimethylamino)-2-methylthio-1,2,4-triazolo[1,5-*a*][1,3,5]triazine (Gilardi, 1973).

The methyl groups of dimethylamino fragment are involved in the formation of weak C-H···N intramolecular hydrogen-bonds with the nitrogen atoms of the heterocyclic system (Tab. 1).


7-Dimethylamino-2-phenyl-1,2,4-triazolo[1,5-*a*][1,3,5]triazin-5-amine crystallizes together with one molecule of methanol (Fig. 1). The methanol molecule participates in the formation of the crystalline structure *via* intermolecular O—H···N, N—H···O and weak C—H···O hydrogen-bonds linking the layers of the molecules of 7-dimethylamino-2-phenyl-1,2,4-triazolo[1,5-*a*][1,3,5]triazin-5-amine (Tab. 1, Fig. 2).

S2. Experimental


2-Phenyl-7-trichloromethyl-1,2,4-triazolo[1,5-*a*][1,3,5]triazin-5-amine (0.66 g, 2.0 mmol) was added to cold (0–5 °C) dimethylamine (5 ml). The mixture was stirred first in ice-bath for 20 min and then for another 60 min at room temperature. Cold water (20 ml) was added and the product was filtered and recrystallized from methanol (m.p. 521 K).

S3. Refinement

All the hydrogen atoms could have been discerned in the difference electron density map, nevertheless, all the H atoms attached to the carbon atoms were constrained in a riding motion approximation [0.94 Å for C_{aryl}-H and 0.97 Å for methyl groups; U_{iso}(H) = 1.2U_{eq}(C_{aryl}) and U_{iso}(H) = 1.5U_{eq}(C_{methyl})] while the hydroxyl and the amino H atoms were refined freely.

**Figure 1**

The molecular structure of the title molecules with the atomic numbering scheme. The displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**Molecular packing viewed along the axis *c*.**7-Dimethylamino-2-phenyl-1,2,4-triazolo[1,5-a][1,3,5]triazin-5-amine methanol solvate***Crystal data*

$M_r = 287.34$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.9963 (5) \text{ \AA}$

$b = 8.0435 (5) \text{ \AA}$

$c = 13.0942 (9) \text{ \AA}$

$\alpha = 93.493 (1)^\circ$

$\beta = 93.972 (1)^\circ$

$\gamma = 102.883 (1)^\circ$

$V = 714.39 (8) \text{ \AA}^3$

$Z = 2$

$F(000) = 304$

$D_x = 1.336 \text{ Mg m}^{-3}$

Melting point: 521 K

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5236 reflections

$\theta = 2.6\text{--}27.5^\circ$

$\mu = 0.09 \text{ mm}^{-1}$

$T = 223 \text{ K}$

Block, colourless

$0.74 \times 0.68 \times 0.40 \text{ mm}$

*Data collection*Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scansAbsorption correction: multi-scan
(*SADABS*; Sheldrick, 2001)

$T_{\min} = 0.935, T_{\max} = 0.964$

9183 measured reflections

3256 independent reflections

2870 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 2.6^\circ$

$h = -9 \rightarrow 9$

$k = -10 \rightarrow 10$

$l = -16 \rightarrow 16$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.047$$

$$wR(F^2) = 0.139$$

$$S = 1.07$$

3256 reflections

205 parameters

0 restraints

53 constraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
map

Hydrogen site location: difference Fourier map

H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0789P)^2 + 0.1582P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.26 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.81509 (16)	0.63570 (13)	0.67882 (8)	0.0325 (3)
N2	0.66405 (16)	0.37327 (13)	0.60113 (8)	0.0322 (3)
N3	0.73812 (15)	0.48432 (13)	0.52921 (8)	0.0296 (2)
N4	0.90611 (16)	0.77645 (13)	0.52784 (9)	0.0343 (3)
N5	0.81734 (16)	0.59004 (13)	0.37296 (8)	0.0327 (3)
N6	0.9623 (2)	0.87296 (16)	0.36975 (11)	0.0432 (3)
H6NA	0.954 (3)	0.859 (2)	0.3018 (16)	0.053 (5)*
H6NB	1.001 (3)	0.975 (3)	0.3987 (15)	0.055 (5)*
N7	0.65525 (16)	0.30863 (13)	0.37239 (8)	0.0343 (3)
C1	0.71566 (17)	0.47092 (16)	0.68750 (10)	0.0309 (3)
C2	0.82710 (17)	0.64251 (15)	0.57841 (10)	0.0304 (3)
C3	0.89340 (18)	0.74312 (16)	0.42567 (10)	0.0330 (3)
C4	0.73645 (17)	0.45873 (15)	0.42375 (10)	0.0291 (3)
C6	0.5786 (2)	0.14747 (17)	0.41673 (12)	0.0439 (3)
H6A	0.6609	0.0681	0.4018	0.066*
H6B	0.4452	0.0991	0.3874	0.066*
H6C	0.5789	0.1681	0.4905	0.066*
C7	0.6562 (2)	0.29631 (19)	0.26060 (11)	0.0428 (3)
H7A	0.6950	0.4101	0.2371	0.064*
H7B	0.5255	0.2413	0.2299	0.064*
H7C	0.7489	0.2293	0.2406	0.064*
C8	0.66609 (18)	0.40434 (16)	0.78684 (10)	0.0333 (3)
C9	0.7706 (2)	0.4838 (2)	0.87779 (11)	0.0431 (3)
H9	0.8728	0.5815	0.8760	0.052*

C10	0.7244 (3)	0.4193 (2)	0.97062 (12)	0.0517 (4)
H10	0.7960	0.4728	1.0318	0.062*
C11	0.5733 (3)	0.2762 (2)	0.97403 (12)	0.0512 (4)
H11	0.5433	0.2323	1.0374	0.061*
C12	0.4663 (2)	0.1978 (2)	0.88451 (12)	0.0456 (4)
H12	0.3624	0.1016	0.8870	0.055*
C13	0.5127 (2)	0.26140 (17)	0.79114 (11)	0.0386 (3)
H13	0.4402	0.2078	0.7302	0.046*
O1S	1.00567 (17)	0.92566 (15)	0.81876 (10)	0.0530 (3)
H1S	0.936 (3)	0.838 (3)	0.7737 (16)	0.061 (6)*
C1S	0.8695 (3)	0.9935 (3)	0.86937 (19)	0.0739 (6)
H1S1	0.8051	0.9119	0.9148	0.111*
H1S2	0.7720	1.0174	0.8195	0.111*
H1S3	0.9360	1.0986	0.9092	0.111*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0333 (5)	0.0275 (5)	0.0349 (6)	0.0040 (4)	0.0007 (4)	0.0013 (4)
N2	0.0341 (5)	0.0271 (5)	0.0341 (5)	0.0038 (4)	0.0024 (4)	0.0047 (4)
N3	0.0296 (5)	0.0238 (5)	0.0336 (5)	0.0028 (4)	0.0011 (4)	0.0019 (4)
N4	0.0364 (6)	0.0256 (5)	0.0383 (6)	0.0017 (4)	0.0027 (4)	0.0021 (4)
N5	0.0333 (5)	0.0287 (5)	0.0353 (6)	0.0054 (4)	0.0039 (4)	0.0019 (4)
N6	0.0550 (8)	0.0304 (6)	0.0409 (7)	0.0007 (5)	0.0081 (5)	0.0054 (5)
N7	0.0370 (6)	0.0277 (5)	0.0355 (6)	0.0036 (4)	0.0005 (4)	-0.0014 (4)
C1	0.0279 (6)	0.0288 (6)	0.0356 (6)	0.0056 (4)	0.0009 (5)	0.0032 (5)
C2	0.0279 (6)	0.0249 (6)	0.0370 (6)	0.0048 (4)	-0.0005 (5)	-0.0006 (5)
C3	0.0302 (6)	0.0286 (6)	0.0402 (7)	0.0059 (5)	0.0042 (5)	0.0041 (5)
C4	0.0253 (5)	0.0280 (6)	0.0339 (6)	0.0068 (4)	0.0009 (4)	0.0008 (5)
C6	0.0527 (8)	0.0260 (6)	0.0473 (8)	-0.0013 (6)	0.0026 (6)	-0.0023 (5)
C7	0.0503 (8)	0.0391 (7)	0.0357 (7)	0.0062 (6)	0.0021 (6)	-0.0066 (5)
C8	0.0330 (6)	0.0326 (6)	0.0348 (7)	0.0081 (5)	0.0020 (5)	0.0042 (5)
C9	0.0414 (7)	0.0436 (7)	0.0394 (7)	0.0010 (6)	-0.0008 (6)	0.0031 (6)
C10	0.0558 (9)	0.0599 (10)	0.0340 (7)	0.0036 (7)	-0.0015 (6)	0.0021 (6)
C11	0.0560 (9)	0.0599 (10)	0.0368 (7)	0.0074 (7)	0.0106 (6)	0.0109 (7)
C12	0.0446 (8)	0.0442 (8)	0.0462 (8)	0.0029 (6)	0.0098 (6)	0.0095 (6)
C13	0.0394 (7)	0.0355 (7)	0.0388 (7)	0.0046 (5)	0.0017 (5)	0.0034 (5)
O1S	0.0485 (6)	0.0470 (6)	0.0569 (7)	-0.0004 (5)	0.0067 (5)	-0.0090 (5)
C1S	0.0679 (12)	0.0663 (12)	0.0864 (14)	0.0126 (10)	0.0233 (11)	-0.0111 (11)

Geometric parameters (\AA , $^\circ$)

N1—C2	1.3269 (17)	C7—H7A	0.9700
N1—C1	1.3681 (16)	C7—H7B	0.9700
N2—C1	1.3172 (17)	C7—H7C	0.9700
N2—N3	1.3846 (14)	C8—C9	1.393 (2)
N3—C4	1.3826 (16)	C8—C13	1.3939 (19)
N3—C2	1.3830 (15)	C9—C10	1.381 (2)

N4—C2	1.3330 (16)	C9—H9	0.9400
N4—C3	1.3412 (18)	C10—C11	1.384 (2)
N5—C4	1.3226 (16)	C10—H10	0.9400
N5—C3	1.3524 (16)	C11—C12	1.382 (2)
N6—C3	1.3330 (17)	C11—H11	0.9400
N6—H6NA	0.89 (2)	C12—C13	1.385 (2)
N6—H6NB	0.86 (2)	C12—H12	0.9400
N7—C4	1.3323 (16)	C13—H13	0.9400
N7—C6	1.4594 (17)	O1S—C1S	1.386 (2)
N7—C7	1.4617 (18)	O1S—H1S	0.92 (2)
C1—C8	1.4709 (18)	C1S—H1S1	0.9700
C6—H6A	0.9700	C1S—H1S2	0.9700
C6—H6B	0.9700	C1S—H1S3	0.9700
C6—H6C	0.9700		
C2—N1—C1	103.31 (10)	N7—C7—H7A	109.5
C1—N2—N3	101.79 (10)	N7—C7—H7B	109.5
C4—N3—C2	119.85 (11)	H7A—C7—H7B	109.5
C4—N3—N2	130.72 (10)	N7—C7—H7C	109.5
C2—N3—N2	109.43 (10)	H7A—C7—H7C	109.5
C2—N4—C3	113.98 (11)	H7B—C7—H7C	109.5
C4—N5—C3	118.92 (11)	C9—C8—C13	119.13 (13)
C3—N6—H6NA	121.0 (13)	C9—C8—C1	120.54 (12)
C3—N6—H6NB	120.1 (13)	C13—C8—C1	120.33 (12)
H6NA—N6—H6NB	118.3 (18)	C10—C9—C8	120.14 (14)
C4—N7—C6	126.56 (12)	C10—C9—H9	119.9
C4—N7—C7	118.21 (11)	C8—C9—H9	119.9
C6—N7—C7	115.01 (11)	C9—C10—C11	120.31 (15)
N2—C1—N1	116.27 (11)	C9—C10—H10	119.8
N2—C1—C8	121.01 (11)	C11—C10—H10	119.8
N1—C1—C8	122.72 (12)	C12—C11—C10	120.14 (14)
N1—C2—N4	128.12 (11)	C12—C11—H11	119.9
N1—C2—N3	109.21 (11)	C10—C11—H11	119.9
N4—C2—N3	122.66 (12)	C11—C12—C13	119.79 (14)
N6—C3—N4	117.28 (12)	C11—C12—H12	120.1
N6—C3—N5	116.20 (13)	C13—C12—H12	120.1
N4—C3—N5	126.52 (12)	C12—C13—C8	120.48 (13)
N5—C4—N7	119.54 (12)	C12—C13—H13	119.8
N5—C4—N3	117.96 (11)	C8—C13—H13	119.8
N7—C4—N3	122.50 (12)	C1S—O1S—H1S	107.0 (13)
N7—C6—H6A	109.5	O1S—C1S—H1S1	109.5
N7—C6—H6B	109.5	O1S—C1S—H1S2	109.5
H6A—C6—H6B	109.5	H1S1—C1S—H1S2	109.5
N7—C6—H6C	109.5	O1S—C1S—H1S3	109.5
H6A—C6—H6C	109.5	H1S1—C1S—H1S3	109.5
H6B—C6—H6C	109.5	H1S2—C1S—H1S3	109.5
C1—N2—N3—C4	178.41 (12)	C6—N7—C4—N5	173.18 (12)

C1—N2—N3—C2	−0.83 (12)	C7—N7—C4—N5	−1.02 (18)
N3—N2—C1—N1	0.51 (14)	C6—N7—C4—N3	−7.5 (2)
N3—N2—C1—C8	179.98 (10)	C7—N7—C4—N3	178.25 (11)
C2—N1—C1—N2	0.03 (14)	C2—N3—C4—N5	−1.60 (17)
C2—N1—C1—C8	−179.44 (11)	N2—N3—C4—N5	179.22 (11)
C1—N1—C2—N4	178.09 (12)	C2—N3—C4—N7	179.11 (11)
C1—N1—C2—N3	−0.57 (13)	N2—N3—C4—N7	−0.1 (2)
C3—N4—C2—N1	−179.24 (12)	N2—C1—C8—C9	159.09 (13)
C3—N4—C2—N3	−0.74 (18)	N1—C1—C8—C9	−21.47 (19)
C4—N3—C2—N1	−178.42 (10)	N2—C1—C8—C13	−21.37 (18)
N2—N3—C2—N1	0.92 (13)	N1—C1—C8—C13	158.07 (12)
C4—N3—C2—N4	2.83 (18)	C13—C8—C9—C10	1.2 (2)
N2—N3—C2—N4	−177.83 (11)	C1—C8—C9—C10	−179.30 (14)
C2—N4—C3—N6	177.38 (11)	C8—C9—C10—C11	−0.5 (3)
C2—N4—C3—N5	−2.68 (19)	C9—C10—C11—C12	−0.6 (3)
C4—N5—C3—N6	−176.15 (11)	C10—C11—C12—C13	0.9 (3)
C4—N5—C3—N4	3.9 (2)	C11—C12—C13—C8	−0.2 (2)
C3—N5—C4—N7	177.83 (11)	C9—C8—C13—C12	−0.8 (2)
C3—N5—C4—N3	−1.48 (17)	C1—C8—C13—C12	179.66 (12)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1S—H1S···N1	0.92 (2)	1.97 (2)	2.8861 (16)	172.8 (18)
N6—H6NB···N4 ⁱ	0.86 (2)	2.11 (2)	2.9679 (17)	178.8 (18)
N6—H6NA···O1S ⁱ	0.89 (2)	2.398 (19)	3.0280 (18)	128.3 (16)
C6—H6C···N2	0.97	2.08	2.8753 (18)	138
C6—H6C···N3	0.97	2.54	2.9484 (17)	105
C7—H7A···N5	0.97	2.22	2.6788 (18)	108
C7—H7C···O1S ⁱⁱ	0.97	2.48	3.4438 (19)	176

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x+2, -y+1, -z+1$.